

First Observation of a Doubly Charged Tetraquark and Its Neutral Partner

R. Aaij *et al.**
(LHCb Collaboration)

 (Received 7 December 2022; accepted 23 January 2023; published 27 July 2023)

A combined amplitude analysis is performed for the decays $B^0 \rightarrow \bar{D}^0 D_s^+ \pi^-$ and $B^+ \rightarrow D^- D_s^+ \pi^+$, which are related by isospin symmetry. The analysis is based on data collected by the LHCb detector in proton-proton collisions at center-of-mass energies of 7, 8, and 13 TeV. The full data sample corresponds to an integrated luminosity of 9 fb^{-1} . Two new resonant states with masses of $2.908 \pm 0.011 \pm 0.020 \text{ GeV}$ and widths of $0.136 \pm 0.023 \pm 0.013 \text{ GeV}$ are observed, which decay to $D_s^+ \pi^+$ and $D_s^+ \pi^-$ respectively. The former state indicates the first observation of a doubly charged open-charm tetraquark state with minimal quark content $[c\bar{s}u\bar{d}]$, and the latter state is a neutral tetraquark composed of $[c\bar{s}\bar{u}d]$ quarks. Both states are found to have spin-parity of 0^+ , and their resonant parameters are consistent with each other, which suggests that they belong to an isospin triplet.

DOI: [10.1103/PhysRevLett.131.041902](https://doi.org/10.1103/PhysRevLett.131.041902)

In the quark model, tetraquark hadrons are predicted in addition to the conventional mesons and baryons [1]. States composed of four different quark types present unambiguous evidence for tetraquarks [2] and have attracted great interest in studies of hadron spectroscopy [3]. The discoveries of the $D_{s0}^*(2317)^+$ [4] and $D_{s1}(2460)^+$ [5] states prompted speculation that they may have a tetraquark component [6,7], but no evidence for isospin partners was found in explicit searches [8,9]. The D0 Collaboration claimed evidence for an $X(5568)$ state with quark configuration of $[b\bar{s}u\bar{d}]$ in decays to $B_s^0 \pi^\pm$ [10,11]. Its existence, however, has not been confirmed by other experiments [12–15].

In the open charm sector, it is natural to search for $D_s^\pm \pi^\pm$ resonances, which would be candidate tetraquark states with minimal quark content of $[c\bar{s}u\bar{d}]$ and $[c\bar{s}\bar{u}d]$ as predicted in the diquark-antidiquark model [16,17]. In particular, an attractive potential in the $D_s^+ \pi^+$ channel facilitates searches for an open-charm tetraquark state according to lattice QCD calculation [18].

In 2020, two new resonant structures, $X_0(2900)$ and $X_1(2900)$, were observed by the LHCb Collaboration in the $D^- K^+$ mass spectrum of the $B^+ \rightarrow D^+ D^- K^+$ decay [19,20], motivating theoretical speculation as to their nature [3]. Predictions of a doubly charged tetraquark $[c\bar{s}u\bar{d}]$ and its isospin partner $[c\bar{s}\bar{u}d]$ were made based on those observations [21–26].

The decays $B^0 \rightarrow \bar{D}^0 D_s^+ \pi^-$ and $B^+ \rightarrow D^- D_s^+ \pi^+$ are ideal channels in which to search for possible exotic states decaying to $D_s \pi$. The only resonances expected to contribute to the two decays are excited \bar{D}^* resonances decaying to $\bar{D}^0 \pi^-$ and $D^- \pi^+$ final states, which have been extensively studied [6,7,27–29]. Any resonant structure that cannot be described by the known \bar{D}^* excited states ($\bar{D}^0 \pi^-$ or $D^- \pi^+$) implies the existence of potential exotic objects decaying into either $D_s \pi$ or DD_s final states. (Charge conjugation is implied throughout this Letter.)

A combined amplitude analysis of the $B^0 \rightarrow \bar{D}^0 D_s^+ \pi^-$ and $B^+ \rightarrow D^- D_s^+ \pi^+$ decays is performed, where the amplitudes in the two decay modes are related through isospin symmetry. The analysis is based on proton-proton collision data collected with the LHCb detector, corresponding to a total integrated luminosity of 9 fb^{-1} at center-of-mass energies 7, 8, and 13 TeV.

The LHCb detector is a single-arm forward spectrometer covering the pseudorapidity range $2 < \eta < 5$, described in detail in Refs. [30,31]. It is designed specifically for the study of particles containing b or c quarks. Simulation is required to model the effects of the detector acceptance and the imposed selection requirements. The simulated samples are generated with PYTHIA [32,33], EVTGEN [34] and the GEANT4 toolkits [35,36] as described in Ref. [37].

A brief summary of the selections is given here, and further details can be found in Ref. [38]. Charmed mesons are reconstructed using the $\bar{D}^0 \rightarrow K^+ \pi^-$, $\bar{D}^0 \rightarrow K^+ \pi^- \pi^- \pi^+$, $D^- \rightarrow K^+ \pi^- \pi^-$, and $D_s^+ \rightarrow K^+ K^- \pi^+$ decays. The invariant mass M of the $\bar{D}^0 \pi^-$ pair is required to be larger than 2050 MeV (natural units with $\hbar = c = 1$ are used throughout) to veto the majority of the $D^*(2010)^-$ contribution from the $B^0 \rightarrow D^{*-} D_s^+$ decay. A multivariate classifier, based on a boosted decision tree [39,40] algorithm in the

*Full author list given at the end of the Letter.




FIG. 1. Dalitz plots of (a) $B^0 \rightarrow \bar{D}^0 D_s^+ \pi^-$ and (b) $B^+ \rightarrow D^- D_s^+ \pi^+$ decays after the full selection is applied.

TMVA toolkit [41,42], is employed to reduce combinatorial backgrounds. In addition, non-double-charm backgrounds are reduced by requirements on the D meson mass and by requiring a significant flight distance of the reconstructed D candidates from the B decay vertex.

After the above selections, 4009 $B^0 \rightarrow \bar{D}^0 D_s^+ \pi^-$ candidates and 3750 $B^+ \rightarrow D^- D_s^+ \pi^+$ candidates are retained. Extended maximum likelihood fits are performed to the $\bar{D}^0 D_s^+ \pi^-$ and $D^- D_s^+ \pi^+$ invariant-mass distributions separately in order to extract signal yields, all in the range of 5230–5630 MeV. The purities of the $B^0 \rightarrow \bar{D}^0 D_s^+ \pi^-$ and $B^+ \rightarrow D^- D_s^+ \pi^+$ samples are 90.7% and 95.2%, respectively. Dalitz plots for the two decays, shown in Fig. 1, show similar features as a result of the isospin symmetry of the decays, indicating the feasibility of a combined treatment. The vertical band at $M^2(\bar{D}\pi) \approx 6 \text{ GeV}^2$ corresponds to the $J^P = 2^+$ state $D_2^*(2460)$ [$D_2^*(2460)^-$ in $B^0 \rightarrow \bar{D}^0 D_s^+ \pi^-$ and $\bar{D}_2^*(2460)^0$ in $B^+ \rightarrow D^- D_s^+ \pi^+$ decays]. Furthermore, a faint horizontal band at $M^2(D_s^+ \pi) \approx 8.5 \text{ GeV}^2$ is visible in both plots.

The conventional contributions anticipated in the two decays include tails from the $D^*(2010)^-$ and $\bar{D}^*(2007)^0$ resonances in $B^+ \rightarrow D^- D_s^+ \pi^+$ and $B^0 \rightarrow \bar{D}^0 D_s^+ \pi^-$ decays, respectively, in addition to the higher excited $\bar{D}_2^*(2460)$, $\bar{D}_1^*(2600)$, $\bar{D}_3^*(2750)$, $\bar{D}_1^*(2760)$, and $\bar{D}(3000)$ states [6]. In previous measurements, only the neutral $\bar{D}_1^*(2600)^0$, $\bar{D}_1^*(2760)^0$, and $\bar{D}(3000)^0$ mesons were observed, while their charged partners were not. For the $\bar{D}\pi$ S -wave components, a quasi-model-independent (QMI) method [29] is applied where 11 spline points are chosen in the $M(\bar{D}\pi)$ spectrum, at (1.9,2.0,2.1,2.2,2.3,2.4,2.5,2.6,2.7, 2.9,3.4) GeV. As isospin symmetry is imposed; these excited \bar{D}^* states are added in both channels before testing the exotic states decaying either to the $D_s^+ \pi$ or $\bar{D} D_s^+$ final state.

The complex amplitude $\mathcal{A}_i(x; \Theta_i)$ for the decay through each intermediate state i is constructed using the helicity formalism, where x denotes variables calculated from the

four-momenta of the final-state particles in each event, and Θ is a set of model parameters. A relativistic Breit–Wigner function is used to describe the line shape of the resonant states, where the resonant parameters are fixed to their known values [6]. The total amplitude $\mathcal{A}(x; \Theta)$ is the sum of the complex amplitudes for each contribution. The spin parity of the $\bar{D}(3000)^0$ state is currently unknown. The 4^+ spin-parity assignment is found to be the most likely [38] and is used in the default model here. In the QMI component, the amplitudes of the first and last spline points are fixed to 0.

A negative-log-likelihood, $-\sum_j^N \ln P(x_j; \Theta)$, is minimized to determine the fit parameters, where N denotes the number of B meson candidates. The probability density function (PDF) is defined as

$$P(x; \Theta) = f_{\text{sig}} P_{\text{sig}}^{\text{norm}}(x; \Theta) + f_{\text{bkg}} P_{\text{bkg}}^{\text{norm}}(x), \quad (1)$$

where the fractions are determined from the fit to the invariant mass distribution of B candidates. The normalized PDF for the signal is expressed as

$$P_{\text{sig}}^{\text{norm}}(x; \Theta) = \frac{\epsilon(x) |\mathcal{A}(x; \Theta)|^2}{I_{\text{sig}}(\Theta)}. \quad (2)$$

The factor $I_{\text{sig}}(\Theta)$ is the signal PDF normalization, obtained using a simulated sample generated uniformly over the phase space. The efficiency map, $\epsilon(x)$, is the signal efficiency derived from the simulated samples after applying all the selection criteria, and smoothed using a kernel density estimation method [43]. The background PDF is obtained from the Dalitz plot distribution of data in the ranges [5400,5700] MeV for B^+ decays, and [5500,5700] MeV for B^0 decays to avoid tails from $B_s^0 \rightarrow \bar{D}^0 D_s^+ \pi^-$ decays. The extrapolation into the signal region of $|M(B) - m(B)| < 20 \text{ MeV}$ is based on a Gaussian process [44]. Here $M(B)$ and $m(B)$ are the candidate mass and known mass [6] of B^0 or B^+ mesons.




FIG. 2. Distributions of (a) $M(\bar{D}^0\pi^-)$, (b) $M(D_s^+\pi^-)$, and (c) $M(\bar{D}^0D_s^+)$ for the $\bar{D}\pi$ components of the $B^0 \rightarrow \bar{D}^0D_s^+\pi^-$ candidates and (d) $M(D^-\pi^+)$, (e) $M(D_s^+\pi^+)$, and (f) $M(D^-D_s^+)$ of the $B^+ \rightarrow D^-D_s^+\pi^+$ candidates. The data are overlaid with the results of the fits.

A simultaneous fit is performed to the two decay samples, where all parameters are shared except the masses, widths, and complex parameters of different excited $\bar{D}^*(2007)^0$ and $D^*(2010)^-$ states. The projections of the fit results are shown in Fig. 2, only taking $\bar{D}\pi$ components into account. The models describe the data well in the different invariant mass projections, except for the $M(D_s^+\pi)$ distributions, where peaking structures near 2.9 GeV in the data cannot be attributed to any $\bar{D}\pi$ component. Additional fits are attempted where new \bar{D}^* states with different spin parities are added, allowing their masses and widths to vary freely, but no satisfactory description of this region is found.

Two $D_s^+\pi$ states are introduced, and under isospin symmetry, they share the following resonance parameters: the complex amplitude factor, the mass, and the width of the states. The $M(D_s^+\pi)$ distributions of the fit results are

shown in Fig. 3, while the other projections are shown in Fig. S1 of the Supplemental Material [45]. Figure S2 of the Supplemental Material [45] shows the fit projection onto $M(D_s^+\pi)$ with and without the additional $D_s^+\pi$ states in the region of $M(\bar{D}\pi) > 2.7$ GeV, where most of the D^{**} contributions are suppressed. The peaks in the $M(D_s^+\pi)$ distribution near 2.9 GeV, as well as the dip near 3.0 GeV, are better described.

Different spin-parity scenarios are tested. The result with $J^P = 0^+$ $D_s^+\pi$ states produces the best likelihood, and is taken as the default fit result. The mass and width are determined to be 2.909 ± 0.010 GeV and 0.134 ± 0.019 GeV, respectively. The other parameters of the result are given in Table S1 of the Supplemental Material [45]. Following the convention in Ref. [46], the new states are named $T_{cs0}^a(2900)^0$ and $T_{cs0}^a(2900)^{++}$. The $M(D_s\pi)$ mass resolution is estimated to be approximately




FIG. 3. Distributions of (a) $M(D_s^+\pi^-)$ of $B^0 \rightarrow \bar{D}^0D_s^+\pi^-$ decays and (b) $M(D_s^+\pi^+)$ for the $B^+ \rightarrow D^-D_s^+\pi^+$ sample. The data are overlaid with the fit results with the inclusion of the new 0^+ $D_s^+\pi$ resonant states.




FIG. 4. Spin analysis for the $J^P = 0^+$ hypothesis over $J^P = 1^-$. The solid blue and black dashed histograms are the distributions of the $2\Delta LL$ of the pseudoexperiments based on the fit results with the 0^+ and 1^+ hypotheses, respectively. The purple vertical line shows the $2\Delta LL$ of the data fitted with the new $D_s^+\pi$ exotic state under the $J^P = 0^+$ and $J^P = 1^-$ hypotheses. The red curve shows the result of a fit with the Gaussian function to the black dashed histogram.

4 MeV near the $T_{c\bar{s}0}^a(2900)$ measured mass, which is much smaller than the width of the $T_{c\bar{s}0}^a(2900)$ state, and is therefore neglected. When using separate parameters for the $T_{c\bar{s}0}^a(2900)^0$ state in $B^0 \rightarrow \bar{D}^0 D_s^+ \pi^-$ decays and the $T_{c\bar{s}0}^a(2900)^{++}$ state in $B^+ \rightarrow D^- D_s^+ \pi^+$ decays, without changing the treatment of the other states, the masses and widths are found to be 2.894 ± 0.011 GeV and 0.121 ± 0.020 GeV for the $T_{c\bar{s}0}^a(2900)^0$ state, and 2.922 ± 0.012 GeV and 0.138 ± 0.029 GeV for the $T_{c\bar{s}0}^a(2900)^{++}$ state. The fit parameters are consistent with the earlier result, as expected given the isospin symmetry of the decays.

To estimate the significance of the new $T_{c\bar{s}0}^a(2900)$ state, pseudoexperiments are generated without the state, and fitted both with and without the $T_{c\bar{s}0}^a(2900)$ state. The sample size of each pseudoexperiment is Poisson fluctuated around the number of the observed candidates. Events are generated in the six channels separately, modeled by their individual background and efficiency maps. The doubled difference of the log-likelihood ($2\Delta LL$) of the two fit results should follow a χ^2 distribution, where the number of degrees of freedom $N_{\text{d.o.f.}}$ is a fit parameter. Using 500 pseudoexperiments, $N_{\text{d.o.f.}}$ is determined to be 6.99 ± 0.17 . In the obtained χ^2 distribution, the value of $2\Delta LL$ from collision data corresponds to a significance greater than 9 standard deviations (σ).

Among other tested J^P hypotheses beyond the default 0^+ , only the $1^- D_s^+\pi$ state leads to a large significance of 6.3σ ($N_{\text{d.o.f.}} = 6.99$), while the other cases are not significant. A test of the hypothesis with the simultaneous presence of a $0^+ T_{c\bar{s}0}^a(2900)$ state and a $1^- D_s^+\pi$ state yields a significance of 1.3σ ($N_{\text{d.o.f.}} = 6.99$), suggesting there is no need in this analysis to include it. Additional $D\pi$,

$D_s^+\pi$, and DD_s^+ resonances with spin-parity hypotheses up to 4^+ are tested with and without the $T_{c\bar{s}0}^a(2900)$ exotic states. In all cases the additional contributions are disfavored, while the $T_{c\bar{s}0}^a(2900)$ significance remains above 5σ with mass and width parameters consistent with the default results.

Pseudoexperiments are performed to confirm the spin parity of the $T_{c\bar{s}0}^a(2900)$ state. A set of 500 pseudoexperiments is generated from the default fit results; another 500 pseudoexperiments are generated from the fit results with a $1^- D_s^+\pi$ exotic state included. Each pseudoexperiment distribution is fitted both under the $J^P = 0^+$ hypothesis and $J^P = 1^-$ hypothesis, and the $2\Delta LL$ are evaluated, shown in Fig. 4. The $2\Delta LL$ value in data corresponds to a significance of about 7.5σ . This demonstrates that the $J^P = 0^+$ hypothesis for the $T_{c\bar{s}0}^a(2900)$ state is favored with a high significance.

The Argand diagram [6] of the $T_{c\bar{s}0}^a(2900)$ state, which is expected to have a circular form, is shown in Fig. 5. An additional fit, in which the Breit-Wigner line shape of the $T_{c\bar{s}0}^a(2900)$ state is replaced by seven spline points on $M(D_s^+\pi)$, is performed. The complex factor associated to each point is free to vary. The line shape of the $D_s^+\pi$ spline model, also shown in Fig. 5, is consistent with the Breit-Wigner line shape, and further supports the resonant character of the $T_{c\bar{s}0}^a(2900)$ state.

The sources of systematic uncertainty in the amplitude analysis fall into four categories: the B meson signal yields, the background model, the efficiency map, and the fixed parameters in the fit. The dominant systematic uncertainties




FIG. 5. Argand diagram for the $T_{c\bar{s}0}^a(2900)$ exotic state. The green star and black dots show the start point and phase variation for a $T_{c\bar{s}0}^a(2900)$ Breit-Wigner function. The red solid line shows the fitted spline function used to model the $J^P = 0^+ D_s^+\pi$ component in the $T_{c\bar{s}0}^a(2900)$ mass region, and the blue error bars show the uncertainties on the values of the spline at the knots.

arise from the fixed Blatt-Weisskopf radius and D^* resonant parameters. Systematic uncertainties due to the Blatt-Weisskopf radius are estimated by varying the parameter from 3.0 GeV^{-1} to 1.5 GeV^{-1} and 4.5 GeV^{-1} . Systematic uncertainties associated with the fixed D^* parameters include those related to the masses and widths of the D^* resonances, which are allowed to vary but are constrained to their measured values according to their uncertainties. The spline points are shifted by $\pm 10 \text{ MeV}$, and the number of spline points is varied by ± 1 to explore their impact. The J^P hypothesis of the $\overline{D}(3000)$ state is changed to 2^+ , the next most probable J^P hypothesis, in the systematic uncertainty study. The total systematic uncertainty is determined by combining all contributions in quadrature. The minimal significance during the variation of the sources of systematic uncertainties is taken as the final significance of the resonance. Possible biases of fit parameters are studied with pseudoexperiments. A set of 500 pseudoexperiments is generated with the default fit results, and fitted with the same model. The distribution of each fit parameter is fitted with a Gaussian function. Both the residual $(\mu_{\text{pseudo}} - \mu_{\text{default}})$ and pull distributions $(\mu_{\text{pseudo}} - \mu_{\text{default}})/\sigma_{\text{pseudo}}$ are obtained from the pseudoexperiments, and used to correct the mean values and uncertainties of the fit results.

In summary, a combined amplitude analysis of the $B^0 \rightarrow \overline{D}^0 D_s^+ \pi^-$ and $B^+ \rightarrow D^- D_s^+ \pi^+$ decays is performed, under the assumption of isospin symmetry, to determine the contributions of intermediate states in the decays. Considering only resonant contributions of D^* mesons, the $M(D\pi)$ distributions in the two B decay channels are well modeled, but in the $M(D_s\pi)$ projection of each channel, peaking structures near 2.9 GeV are not well described. Two new $D_s\pi$ exotic resonances are included in the fit model. Assuming the neutral $D_s^+ \pi^-$ resonance and the doubly charged $D_s^+ \pi^+$ resonance belong to the same isospin triplet, the common mass and width are determined to be

$$M = 2.908 \pm 0.011 \pm 0.020 \text{ GeV},$$

$$\Gamma = 0.136 \pm 0.023 \pm 0.013 \text{ GeV},$$

where the first and second uncertainties are statistical and systematic, respectively. The spin parity is determined to be 0^+ with a significance of about 7.5σ with respect to the 1^- hypothesis. After considering the systematic uncertainties, the significance of the $T_{c\bar{s}0}^a(2900)$ state is estimated to be greater than 9σ , taking into account the look-elsewhere effect. Meanwhile, amplitude fits with separate $T_{c\bar{s}0}^a(2900)^0$ and $T_{c\bar{s}0}^a(2900)^{++}$ resonance parameters are also performed. The masses and widths of the two resonances are

$$T_{c\bar{s}0}^a(2900)^0: M = 2.892 \pm 0.014 \pm 0.015 \text{ GeV},$$

$$\Gamma = 0.119 \pm 0.026 \pm 0.013 \text{ GeV},$$

$$T_{c\bar{s}0}^a(2900)^{++}: M = 2.921 \pm 0.017 \pm 0.020 \text{ GeV},$$

$$\Gamma = 0.137 \pm 0.032 \pm 0.017 \text{ GeV},$$

which are in good agreement. The significance of the two resonances is found to be $8.0\sigma(N_{\text{d.o.f.}} = 7.29)$ for the $T_{c\bar{s}0}^a(2900)^0$ state and $6.5\sigma(N_{\text{d.o.f.}} = 8.53)$ for the $T_{c\bar{s}0}^a(2900)^{++}$ state, including systematic effects. The additional details are described in Ref. [38]. This is the first observation of a doubly charged tetraquark candidate, $T_{c\bar{s}0}^a(2900)^{++}([c\bar{s}u\bar{d}])$, and of its neutral isospin partner, $T_{c\bar{s}0}^a(2900)^0([c\bar{s}\bar{u}d])$. They belong to a new type of open-charm tetraquark states with c and \bar{s} quarks. The obtained mass of the $T_{c\bar{s}0}^a(2900)$ state is consistent with that of another 0^+ open-charm tetraquark, the $X_0(2900)^0([c\bar{s}\bar{u}\bar{d}])$ state discovered in the D^+K^- final state [19,20], but their widths and flavor contents are different. The observation of the $T_{c\bar{s}0}^a(2900)$ states is a significant step in the study of exotic hadron spectroscopy.

We express our gratitude to our colleagues in the CERN accelerator departments for the excellent performance of the LHC. We thank the technical and administrative staff at the LHCb institutes. We acknowledge support from CERN and from the national agencies: CAPES, CNPq, FAPERJ, and FINEP (Brazil); MOST and NSFC (China); CNRS/IN2P3 (France); BMBF, DFG, and MPG (Germany); INFN (Italy); NWO (Netherlands); MNiSW and NCN (Poland); MEN/IFA (Romania); MICINN (Spain); SNSF and SER (Switzerland); NASU (Ukraine); STFC (United Kingdom); and DOE NP and NSF (USA). We acknowledge the computing resources that are provided by CERN, IN2P3 (France), KIT, and DESY (Germany); INFN (Italy); SURF (Netherlands); PIC (Spain); GridPP (United Kingdom); CSCS (Switzerland); IFIN-HH (Romania); CBPF (Brazil); Polish WLCG (Poland), and NERSC (USA). We are indebted to the communities behind the multiple open-source software packages on which we depend. Individual groups or members have received support from ARC and ARDC (Australia); Minciencias (Colombia); AvH Foundation (Germany); EPLANET, Marie Skłodowska-Curie Actions, and ERC (European Union); A*MIDEX, ANR, IPhU and Labex P2IO, and Région Auvergne-Rhône-Alpes (France); Key Research Program of Frontier Sciences of CAS, CAS PIFI, CAS CCEPP, Fundamental Research Funds for the Central Universities, and Sci. & Tech. Program of Guangzhou (China); GVA, XuntaGal, GENCAT and Prog. Atracción Talento, CM (Spain); SRC (Sweden); and the Leverhulme Trust, the Royal Society, and UKRI (United Kingdom).

- [1] M. Gell-Mann, A schematic model of baryons and mesons, *Phys. Lett.* **8**, 214 (1964).
- [2] R. Molina, T. Branz, and E. Oset, A new interpretation for the $D_{s2}^*(2573)$ and the prediction of novel exotic charmed mesons, *Phys. Rev. D* **82**, 014010 (2010).
- [3] H.-X. Chen, W. Chen, X. Liu, Y.-R. Liu, and S.-L. Zhu, An updated review of the new hadron states, *Rep. Prog. Phys.* **86**, 026201 (2023).
- [4] B. Aubert *et al.* (BABAR Collaboration), Observation of a Narrow Meson Decaying to $D_s^+\pi^0$ at a Mass of 2.32-GeV/ c^2 , *Phys. Rev. Lett.* **90**, 242001 (2003).
- [5] D. Besson *et al.* (CLEO Collaboration), Observation of a narrow resonance of mass 2.46-GeV/ c^2 decaying to $D_s^{*+}\pi^0$ and confirmation of the $D_{sj}^*(2317)$ state, *Phys. Rev. D* **68**, 032002 (2003); **75**, 119908(E) (2007).
- [6] R. L. Workman *et al.* (Particle Data Group), Review of particle physics, *Prog. Theor. Exp. Phys.* **2022**, 083C01 (2022).
- [7] H.-X. Chen, W. Chen, X. Liu, Y.-R. Liu, and S.-L. Zhu, A review of the open charm and open bottom systems, *Rep. Prog. Phys.* **80**, 076201 (2017).
- [8] B. Aubert *et al.* (BABAR Collaboration), A study of the $D_{sj}^*(2317)$ and $D_{sj}(2460)$ mesons in inclusive c anti-c production near $\sqrt{s} = 10.6$ GeV, *Phys. Rev. D* **74**, 032007 (2006).
- [9] S.-K. Choi *et al.* (Belle Collaboration), Measurements of $B \rightarrow \bar{D}D_{s0}^{*+}(2317)$ decay rates and a search for isospin partners of the $D_{s0}^{*+}(2317)$, *Phys. Rev. D* **91**, 092011 (2015).
- [10] V. M. Abazov *et al.* (D0 Collaboration), Evidence for a $B_s^0\pi^\pm$ State, *Phys. Rev. Lett.* **117**, 022003 (2016).
- [11] V. M. Abazov *et al.* (D0 Collaboration), Study of the $X^\pm(5568)$ state with semileptonic decays of the B_s^0 meson, *Phys. Rev. D* **97**, 092004 (2018).
- [12] R. Aaij *et al.* (LHCb Collaboration), Search for Structure in the $B_s^0\pi^\pm$ Invariant Mass Spectrum, *Phys. Rev. Lett.* **117**, 152003 (2016).
- [13] A. M. Sirunyan *et al.* (CMS Collaboration), Search for the $X(5568)$ State Decaying into $B_s^0\pi^\pm$ in Proton-Proton Collisions at $\sqrt{s} = 8$ TeV, *Phys. Rev. Lett.* **120**, 202005 (2018).
- [14] M. Aaboud *et al.* (ATLAS Collaboration), Search for a Structure in the $B_s^0\pi^\pm$ Invariant Mass Spectrum with the ATLAS Experiment, *Phys. Rev. Lett.* **120**, 202007 (2018).
- [15] T. Aaltonen *et al.* (CDF Collaboration), A Search for the Exotic Meson $X(5568)$ with the Collider Detector at Fermilab, *Phys. Rev. Lett.* **120**, 202006 (2018).
- [16] S. S. Agaev, K. Azizi, and H. Sundu, Charmed partner of the exotic $X(5568)$ state and its properties, *Phys. Rev. D* **93**, 094006 (2016).
- [17] W. Chen, H.-X. Chen, X. Liu, T. G. Steele, and S.-L. Zhu, Open-flavor charm and bottom $sq\bar{q}\bar{Q}$ and $qq\bar{q}\bar{Q}$ tetraquark states, *Phys. Rev. D* **95**, 114005 (2017).
- [18] J. Baeza-Ballesteros, P. Hernández, and F. Romero-López, A lattice study of $\pi\pi$ scattering at large N_c , *J. High Energy Phys.* **06** (2022) 049.
- [19] R. Aaij *et al.* (LHCb Collaboration), Model-Independent Study of Structure in $B^+ \rightarrow D^+D^-K^+$ Decays, *Phys. Rev. Lett.* **125**, 242001 (2020).
- [20] R. Aaij *et al.* (LHCb Collaboration), Amplitude analysis of the $B^+ \rightarrow D^+D^-K^+$ decay, *Phys. Rev. D* **102**, 112003 (2020).
- [21] X.-G. He, W. Wang, and R. Zhu, Open-charm tetraquark X_c and open-bottom tetraquark X_b , *Eur. Phys. J. C* **80**, 1026 (2020).
- [22] Q.-F. Lü, D.-Y. Chen, and Y.-B. Dong, Open charm and bottom tetraquarks in an extended relativized quark model, *Phys. Rev. D* **102**, 074021 (2020).
- [23] T. J. Burns and E. S. Swanson, Discriminating among interpretations for the $X(2900)$ states, *Phys. Rev. D* **103**, 014004 (2021).
- [24] S. S. Agaev, K. Azizi, and H. Sundu, Vector resonance $X_1(2900)$ and its structure, *Nucl. Phys.* **A1011**, 122202 (2021).
- [25] S. S. Agaev, K. Azizi, and H. Sundu, Doubly charged vector tetraquark $Z_V^{++} = [cu][\bar{s}\bar{d}]$, *Phys. Lett. B* **820**, 136530 (2021).
- [26] K. Azizi and U. Özdem, Magnetic dipole moments of the T_{cc}^+ and Z_V^{++} tetraquark states, *Phys. Rev. D* **104**, 114002 (2021).
- [27] R. Aaij *et al.* (LHCb Collaboration), Study of D_J meson decays to $D^+\pi^-$, $D^0\pi^+$ and $D^{*+}\pi^-$ final states in p p collisions, *J. High Energy Phys.* **09** (2013) 145.
- [28] R. Aaij *et al.* (LHCb Collaboration), Dalitz plot analysis of $B^0 \rightarrow \bar{D}^0\pi^+\pi^-$ decays, *Phys. Rev. D* **92**, 032002 (2015).
- [29] R. Aaij *et al.* (LHCb Collaboration), Amplitude analysis of $B^- \rightarrow D^+\pi^-\pi^-$ decays, *Phys. Rev. D* **94**, 072001 (2016).
- [30] A. A. Alves Jr. *et al.* (LHCb Collaboration), The LHCb detector at the LHC, *J. Instrum.* **3**, S08005 (2008).
- [31] R. Aaij *et al.* (LHCb Collaboration), LHCb detector performance, *Int. J. Mod. Phys. A* **30**, 1530022 (2015).
- [32] T. Sjöstrand, S. Mrenna, and P. Skands, A brief introduction to PYTHIA8.1, *Comput. Phys. Commun.* **178**, 852 (2008).
- [33] T. Sjöstrand, S. Mrenna, and P. Skands, PYTHIA6.4 physics and manual, *J. High Energy Phys.* **05** (2006) 026.
- [34] D. J. Lange, The EVTGEN particle decay simulation package, *Nucl. Instrum. Methods Phys. Res., Sect. A* **462**, 152 (2001).
- [35] J. Allison *et al.* (GEANT4 Collaboration), GEANT4 developments and applications, *IEEE Trans. Nucl. Sci.* **53**, 270 (2006).
- [36] S. Agostinelli *et al.* (GEANT4 Collaboration), GEANT4: A simulation toolkit, *Nucl. Instrum. Methods Phys. Res., Sect. A* **506**, 250 (2003).
- [37] M. Clemencic, G. Corti, S. Easo, C. R. Jones, S. Miglioranza, M. Pappagallo, and P. Robbe, The LHCb simulation application, Gauss: Design, evolution and experience, *J. Phys. Conf. Ser.* **331**, 032023 (2011).
- [38] R. Aaij *et al.* (LHCb Collaboration), companion paper, Amplitude analysis of $B^0 \rightarrow \bar{D}^0D_s^+\pi^-$ and $B^+ \rightarrow D^-D_s^+\pi^+$ decays, *Phys. Rev. D* **108**, 012017 (2023).
- [39] L. Breiman, J. H. Friedman, R. A. Olshen, and C. J. Stone, *Classification and Regression Trees* (Wadsworth international group, Belmont, California, USA, 1984).
- [40] Y. Freund and R. E. Schapire, A decision-theoretic generalization of on-line learning and an application to boosting, *J. Comput. Syst. Sci.* **55**, 119 (1997).
- [41] H. Voss, A. Hoecker, J. Stelzer, and F. Tegenfeldt, TMVA—Toolkit for multivariate data analysis with ROOT, *Proc. Sci., ACAT2007* (2007) 040.
- [42] A. Hoecker *et al.*, TMVA 4—Toolkit for multivariate data analysis with ROOT. Users guide, [arXiv:physics/0703039](https://arxiv.org/abs/physics/0703039).

- [43] A. Poluektov, Kernel density estimation of a multi-dimensional efficiency profile, *J. Instrum.* **10**, P02011 (2015).
- [44] A. Mathad, D. O'Hanlon, A. Poluektov, and R. Rabadan, Efficient description of experimental effects in amplitude analyses, *J. Instrum.* **16**, P06016 (2021).
- [45] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevLett.131.041902> for the other projections, the slice plots, the fit parameters, and fit fractions of fit results.
- [46] T. Gershon (LHCb Collaboration), Exotic hadron naming convention, [arXiv:2206.15233](https://arxiv.org/abs/2206.15233).

R. Aaij³², A. S. W. Abdelmotteleb⁵⁰, C. Abellan Beteta⁴⁴, F. Abudinén⁵⁰, T. Ackernley⁵⁴, B. Adeva⁴⁰, M. Adinolfi⁴⁸, P. Adlarson⁷⁷, H. Afsharnia⁹, C. Agapopoulou¹³, C. A. Aidala⁷⁸, S. Aiola²⁵, Z. Ajaltouni⁹, S. Akar⁵⁹, K. Akiba³², J. Albrecht¹⁵, F. Alessio⁴², M. Alexander⁵³, A. Alfonso Alberio³⁹, Z. Aliouche⁵⁶, P. Alvarez Cartelle⁴⁹, R. Amalric¹³, S. Amato², J. L. Amey⁴⁸, Y. Amhis^{11,42}, L. An⁴², L. Anderlini²², M. Andersson⁴⁴, A. Andreianov³⁸, M. Andreotti²¹, D. Andreou⁶², D. Ao⁶, F. Archilli¹⁷, A. Artamonov³⁸, M. Artuso⁶², E. Aslanides¹⁰, M. Atzeni⁴⁴, B. Audurier¹², S. Bachmann¹⁷, M. Bachmayer⁴³, J. J. Back⁵⁰, A. Bailly-reyre¹³, P. Baladron Rodriguez⁴⁰, V. Balagura¹², W. Baldini²¹, J. Baptista de Souza Leite¹, M. Barbetti^{22,b}, R. J. Barlow⁵⁶, S. Barsuk¹¹, W. Barter⁵⁵, M. Bartolini⁴⁹, F. Baryshnikov³⁸, J. M. Basels¹⁴, G. Bassi^{29,c}, B. Batsukh⁴, A. Battig¹⁵, A. Bay⁴³, A. Beck⁵⁰, M. Becker¹⁵, F. Bedeschi²⁹, I. B. Bediaga¹, A. Beiter⁶², V. Belavin³⁸, S. Belin⁴⁰, V. Bellee⁴⁴, K. Belous³⁸, I. Belov³⁸, I. Belyaev³⁸, G. Benane¹⁰, G. Bencivenni²³, E. Ben-Haim¹³, A. Berezhnoy³⁸, R. Bernet⁴⁴, S. Bernet Andres⁷⁶, D. Berninghoff¹⁷, H. C. Bernstein⁶², C. Bertella⁵⁶, A. Bertolin²⁸, C. Betancourt⁴⁴, F. Betti⁴², I. A. Bezshyiko⁴⁴, S. Bhasin⁴⁸, J. Bhom³⁵, L. Bian⁶⁸, M. S. Bieker¹⁵, N. V. Biesuz²¹, S. Bifani⁴⁷, P. Billoir¹³, A. Biolchini³², M. Birch⁵⁵, F. C. R. Bishop⁴⁹, A. Bitadze⁵⁶, A. Bizzeti¹⁵, M. P. Blago⁴⁹, T. Blake⁵⁰, F. Blanc⁴³, J. E. Blank¹⁵, S. Blusk⁶², D. Bobulska⁵³, J. A. Boelhauve¹⁵, O. Boente Garcia¹², T. Boettcher⁵⁹, A. Boldyrev³⁸, C. S. Bolognani⁷⁴, R. Bolzonella^{21,d}, N. Bondar^{38,42}, F. Borgato²⁸, S. Borghi⁵⁶, M. Borsato¹⁷, J. T. Borsuk³⁵, S. A. Bouchiba⁴³, T. J. V. Bowcock⁵⁴, A. Boyer⁴², C. Bozzi²¹, M. J. Bradley⁵⁵, S. Braun⁶⁰, A. Brea Rodriguez⁴⁰, J. Brodzicka³⁵, A. Brossa Gonzalo⁴⁰, J. Brown⁵⁴, D. Brundu²⁷, A. Buonaura⁴⁴, L. Buonincontri²⁸, A. T. Burke⁵⁶, C. Burr⁴², A. Bursche⁶⁶, A. Butkevich³⁸, J. S. Butter³², J. Buytaert⁴², W. Byczynski⁴², S. Cadet²⁷, H. Cai⁶⁸, R. Calabrese^{21,d}, L. Calefice¹⁵, S. Cali²³, R. Calladine⁴⁷, M. Calvi^{26,e}, M. Calvo Gomez⁷⁶, P. Campana²³, D. H. Campora Perez⁷⁴, A. F. Campoverde Quezada⁶, S. Capelli^{26,e}, L. Capriotti²⁰, A. Carbone^{20,f}, G. Carboni³¹, R. Cardinale^{24,g}, A. Cardini²⁷, P. Carniti^{26,e}, L. Carus¹⁴, A. Casais Vidal⁴⁰, R. Caspary¹⁷, G. Casse⁵⁴, M. Cattaneo⁴², G. Cavallero⁴², V. Cavallini^{21,d}, S. Celani⁴³, J. Cerasoli¹⁰, D. Cervenkov⁵⁷, A. J. Chadwick⁵⁴, M. G. Chapman⁴⁸, M. Charles¹³, Ph. Charpentier⁴², C. A. Chavez Barajas⁵⁴, M. Chefdeville⁸, C. Chen³, S. Chen⁴, A. Chernov³⁵, S. Chernyshenko⁴⁶, V. Chobanova⁴⁰, S. Cholak⁴³, M. Chruszcz³⁵, A. Chubykin³⁸, V. Chulikov³⁸, P. Ciambone²³, M. F. Cicala⁵⁰, X. Cid Vidal⁴⁰, G. Ciezarek⁴², G. Ciullo^{21,d}, P. E. L. Clarke⁵², M. Clemencic⁴², H. V. Cliff⁴⁹, J. Closier⁴², J. L. Cobbedick⁵⁶, V. Coco⁴², J. A. B. Coelho¹¹, J. Cogan¹⁰, E. Cogneras⁹, L. Cojocariu³⁷, P. Collins⁴², T. Colombo⁴², L. Congedo¹⁹, A. Contu²⁷, N. Cooke⁴⁷, I. Corredoira⁴⁰, G. Corti⁴², B. Couturier⁴², D. C. Craik⁴⁴, M. Cruz Torres^{1,h}, R. Currie⁵², C. L. Da Silva⁶¹, S. Dadabaev³⁸, L. Dai⁶⁵, X. Dai⁵, E. Dall'Occo¹⁵, J. Dalseno⁴⁰, C. D'Ambrosio⁴², J. Daniel⁹, A. Danilina³⁸, P. d'Argent¹⁵, J. E. Davies⁵⁶, A. Davis⁵⁶, O. De Aguiar Francisco⁵⁶, J. de Boer⁴², K. De Bruyn⁷³, S. De Capua⁵⁶, M. De Cian⁴³, U. De Freitas Carneiro Da Graca¹, E. De Lucia²³, J. M. De Miranda¹, L. De Paula², M. De Serio^{19,i}, D. De Simone⁴⁴, P. De Simone²³, F. De Vellis¹⁵, J. A. de Vries⁷⁴, C. T. Dean⁶¹, F. Debernardis^{19,i}, D. Decamp⁸, V. Dedu¹⁰, L. Del Buono¹³, B. Delaney⁵⁸, H.-P. Dembinski¹⁵, V. Denysenko⁴⁴, O. Deschamps⁹, F. Dettori^{27,j}, B. Dey⁷¹, A. Di Cicco²³, P. Di Nezza²³, I. Diachkov³⁸, S. Didenko³⁸, L. Dieste Maronas⁴⁰, S. Ding⁶², V. Dobishuk⁴⁶, A. Dolmatov³⁸, C. Dong³, A. M. Donohoe¹⁸, F. Dordei²⁷, A. C. dos Reis¹, L. Douglas⁵³, A. G. Downes⁸, P. Duda⁷⁵, M. W. Dudek³⁵, L. Dufour⁴², V. Duk⁷², P. Durante⁴², M. M. Duras⁷⁵, J. M. Durham⁶¹, D. Dutta⁵⁶, A. Dziurda³⁵, A. Dzyuba³⁸, S. Easo⁵¹, U. Egede⁶³, V. Egorychev³⁸, S. Eidelman^{38,a}, C. Eirea Orro⁴⁰, S. Eisenhardt⁵², E. Ejopu⁵⁶, S. Ek-In⁴³, L. Eklund⁷⁷, S. Ely⁶², A. Ene³⁷, E. Epple⁵⁹, S. Escher¹⁴, J. Eschle⁴⁴, S. Esen⁴⁴, T. Evans⁵⁶, F. Fabiano^{27,j}, L. N. Falcao¹, Y. Fan⁶, B. Fang⁶⁸, L. Fantini^{72,k}, M. Faria⁴³, S. Farry⁵⁴, D. Fazzini^{26,e}, L. F. Felkowski⁷⁵, M. Feo⁴², M. Fernandez Gomez⁴⁰

A. D. Fernez⁶⁰, F. Ferrari²⁰, L. Ferreira Lopes⁴³, F. Ferreira Rodrigues², S. Ferreres Sole³², M. Ferrillo⁴⁴, M. Ferro-Luzzi⁴², S. Filippov³⁸, R. A. Fini¹⁹, M. Fiorini^{21,d}, M. Firlej³⁴, K. M. Fischer⁵⁷, D. S. Fitzgerald⁷⁸, C. Fitzpatrick⁵⁶, T. Fiutowski³⁴, F. Fleuret¹², M. Fontana¹³, F. Fontanelli^{24,g}, R. Forty⁴², D. Foulds-Holt⁴⁹, V. Franco Lima⁵⁴, M. Franco Sevilla⁶⁰, M. Frank⁴², E. Franzoso^{21,d}, G. Frau¹⁷, C. Frei⁴², D. A. Friday⁵³, J. Fu⁶, Q. Fuehring¹⁵, T. Fulghesu¹³, E. Gabriel³², G. Galati^{19,i}, M. D. Galati³², A. Gallas Torreira⁴⁰, D. Galli^{20,f}, S. Gambetta^{52,42}, Y. Gan³, M. Gandelman², P. Gandini²⁵, Y. Gao⁷, Y. Gao⁵, M. Garau^{27,j}, L. M. Garcia Martin⁵⁰, P. Garcia Moreno³⁹, J. García Pardiñas^{26,e}, B. Garcia Plana⁴⁰, F. A. Garcia Rosales¹², L. Garrido³⁹, C. Gaspar⁴², R. E. Geertsema³², D. Gerick¹⁷, L. L. Gerken¹⁵, E. Gersabeck⁵⁶, M. Gersabeck⁵⁶, T. Gershon⁵⁰, L. Giambastiani²⁸, V. Gibson⁴⁹, H. K. Giemza³⁶, A. L. Gilman⁵⁷, M. Giovannetti^{23,l}, A. Gioventù⁴⁰, P. Gironella Gironell³⁹, C. Giugliano^{21,d}, M. A. Giza³⁵, K. Gizdov⁵², E. L. Gkougkousis⁴², V. V. Gligorov^{13,42}, C. Göbel⁶⁴, E. Golobardes⁷⁶, D. Golubkov³⁸, A. Golutvin^{55,38}, A. Gomes^{1,m}, S. Gomez Fernandez³⁹, F. Goncalves Abrantes⁵⁷, M. Goncerz³⁵, G. Gong³, I. V. Gorelov³⁸, C. Gotti²⁶, J. P. Grabowski⁷⁰, T. Grammatico¹³, L. A. Granado Cardoso⁴², E. Graugés³⁹, E. Graverini⁴³, G. Graziani¹⁰, A. T. Grecu³⁷, L. M. Greeven³², N. A. Grieser⁴, L. Grillo⁵³, S. Gromov³⁸, B. R. Gruberg Cazon⁵⁷, C. Gu³, M. Guarise^{21,d}, M. Guittiere¹¹, P. A. Günther¹⁷, E. Gushchin³⁸, A. Guth¹⁴, Y. Guz³⁸, T. Gys⁴², T. Hadavizadeh⁶³, G. Haefeli⁴³, C. Haen⁴², J. Haimberger⁴², S. C. Haines⁴⁹, T. Halewood-leagas⁵⁴, M. M. Halvorsen⁴², P. M. Hamilton⁶⁰, J. Hammerich⁵⁴, Q. Han⁷, X. Han¹⁷, E. B. Hansen⁵⁶, S. Hansmann-Menzemer¹⁷, L. Hao⁶, N. Harnew⁵⁷, T. Harrison⁵⁴, C. Hasse⁴², M. Hatch⁴², J. He^{6,n}, K. Heijhoff³², C. Henderson⁵⁹, R. D. L. Henderson^{63,50}, A. M. Hennequin⁵⁸, K. Hennessy⁵⁴, L. Henry⁴², J. Herd⁵⁵, J. Heuel¹⁴, A. Hicheur², D. Hill⁴³, M. Hilton⁵⁶, S. E. Hollitt¹⁵, J. Horswill⁵⁶, R. Hou⁷, Y. Hou⁸, J. Hu¹⁷, J. Hu⁶⁶, W. Hu⁵, X. Hu³, W. Huang⁶, X. Huang⁶⁸, W. Hulsbergen³², R. J. Hunter⁵⁰, M. Hushchyn³⁸, D. Hutchcroft⁵⁴, P. Ibis¹⁵, M. Idzik³⁴, D. Ilin³⁸, P. Ilten⁵⁹, A. Inglessi³⁸, A. Iniukhin³⁸, A. Ishteev³⁸, K. Ivshin³⁸, R. Jacobsson⁴², H. Jage¹⁴, S. J. Jaimes Elles⁴¹, S. Jakobsen⁴², E. Jans³², B. K. Jashal⁴¹, A. Jawahery⁶⁰, V. Jevtic¹⁵, E. Jiang⁶⁰, X. Jiang^{4,6}, Y. Jiang⁶, M. John⁵⁷, D. Johnson⁵⁸, C. R. Jones⁴⁹, T. P. Jones⁵⁰, B. Jost⁴², N. Jurik⁴², I. Juszcak³⁵, S. Kandybei⁴⁵, Y. Kang³, M. Karacson⁴², D. Karpenkov³⁸, M. Karpov³⁸, J. W. Kautz⁵⁹, F. Keizer⁴², D. M. Keller⁶², M. Kenzie⁵⁰, T. Ketel³², B. Khanji¹⁵, A. Kharisova³⁸, S. Kholodenko³⁸, G. Khreich¹¹, T. Kim¹⁴, V. S. Kirsebom⁴³, O. Kitouni⁵⁸, S. Klaver³³, N. Kleijne^{29,c}, K. Klimaszewski³⁶, M. R. Kmiec³⁶, S. Koliiev⁴⁶, A. Kondybayeva³⁸, A. Konoplyannikov³⁸, P. Kopciewicz³⁴, R. Kopečna¹⁷, P. Koppenburg³², M. Korolev³⁸, I. Kostiuk^{32,46}, O. Kot⁴⁶, S. Kotriakhova³⁸, A. Kozachuk³⁸, P. Kravchenko³⁸, L. Kravchuk³⁸, R. D. Krawczyk⁴², M. Kreps⁵⁰, S. Kretschmar¹⁴, P. Krokovny³⁸, W. Krupa³⁴, W. Krzemien³⁶, J. Kubat¹⁷, S. Kubis⁷⁵, W. Kucewicz^{35,34}, M. Kucharczyk³⁵, V. Kudryavtsev³⁸, A. Kupsc⁷⁷, D. Lacarrere⁴², G. Lafferty⁵⁶, A. Lai²⁷, A. Lampis^{27,j}, D. Lancierini⁴⁴, C. Landesa Gomez⁴⁰, J. J. Lane⁵⁶, R. Lane⁴⁸, G. Lanfranchi²³, C. Langenbruch¹⁴, J. Langer¹⁵, O. Lantwin³⁸, T. Latham⁵⁰, F. Lazzari^{29,o}, M. Lazzaroni^{25,p}, R. Le Gac¹⁰, S. H. Lee⁷⁸, R. Lefèvre⁹, A. Leflat³⁸, S. Legotin³⁸, P. Lenisa^{21,d}, O. Leroy¹⁰, T. Lesiak³⁵, B. Leverington¹⁷, A. Li³, H. Li⁶⁶, K. Li⁷, P. Li¹⁷, P.-R. Li⁶⁷, S. Li⁷, T. Li⁴, T. Li⁶⁶, Y. Li⁴, Z. Li⁶², X. Liang⁶², C. Lin⁶, T. Lin⁵¹, R. Lindner⁴², V. Lisovskyi¹⁵, R. Litvinov^{27,j}, G. Liu⁶⁶, H. Liu⁶, Q. Liu⁶, S. Liu^{4,6}, Y. Liu⁶, A. Lobo Salvia³⁹, A. Loi²⁷, R. Lollini⁷², J. Lomba Castro⁴⁰, I. Longstaff⁵³, J. H. Lopes², A. Lopez Huertas³⁹, S. López Soliño⁴⁰, G. H. Lovell⁴⁹, Y. Lu^{4,q}, C. Lucarelli^{22,b}, D. Lucchesi^{28,r}, S. Luchuk³⁸, M. Lucio Martinez⁷⁴, V. Lukashenko^{32,46}, Y. Luo³, A. Lupato⁵⁶, E. Luppi^{21,d}, A. Lusiani^{29,c}, K. Lynch¹⁸, X.-R. Lyu⁶, L. Ma⁴, R. Ma⁶, S. Maccolini²⁰, F. Machefert¹¹, F. Maciuc³⁷, I. Mackay⁵⁷, V. Macko⁴³, P. Mackowiak¹⁵, L. R. Madhan Mohan⁴⁸, A. Maevskiy³⁸, D. Maisuzenko³⁸, M. W. Majewski³⁴, J. J. Malczewski³⁵, S. Malde⁵⁷, B. Malecki^{35,42}, A. Malinin³⁸, T. Maltsev³⁸, G. Manca^{27,j}, G. Mancinelli¹⁰, C. Mancuso^{11,25,p}, D. Manuzzi²⁰, C. A. Manzari⁴⁴, D. Marangotto^{25,p}, J. F. Marchand⁸, U. Marconi²⁰, S. Mariani^{22,b}, C. Marin Benito³⁹, J. Marks¹⁷, A. M. Marshall⁴⁸, P. J. Marshall⁵⁴, G. Martelli^{72,k}, G. Martellotti³⁰, L. Martinazzoli^{42,e}, M. Martinelli^{26,e}, D. Martinez Santos⁴⁰, F. Martinez Vidal⁴¹, A. Massafferri¹, M. Materok¹⁴, R. Matev⁴², A. Mathad⁴⁴, V. Matiunin³⁸, C. Matteuzzi²⁶, K. R. Mattioli¹², A. Mauri³², E. Maurice¹², J. Mauricio³⁹, M. Mazurek⁴², M. McCann⁵⁵, L. McConnell¹⁸, T. H. McGrath⁵⁶, N. T. McHugh⁵³, A. McNab⁵⁶, R. McNulty¹⁸, J. V. Mead⁵⁴, B. Meadows⁵⁹, G. Meier¹⁵, D. Melnychuk³⁶, S. Meloni^{26,e}, M. Merk^{32,74}, A. Merli^{25,p}, L. Meyer Garcia², D. Miao^{4,6}, M. Mikhasenko^{70,s}, D. A. Milanec⁶⁹, E. Millard⁵⁰, M. Milovanovic⁴²

M.-N. Minard,^{8,a} A. Minotti,^{26,e} T. Miralles,⁹ S. E. Mitchell,⁵² B. Mitreska,⁵⁶ D. S. Mitzel,¹⁵ A. Mödden,¹⁵ R. A. Mohammed,⁵⁷ R. D. Moise,¹⁴ S. Mokhnenko,³⁸ T. Mombächer,⁴⁰ M. Monk,^{50,63} I. A. Monroy,⁶⁹ S. Monteil,⁹ M. Morandin,²⁸ G. Morello,²³ M. J. Morello,^{29,c} J. Moron,³⁴ A. B. Morris,⁷⁰ A. G. Morris,⁵⁰ R. Mountain,⁶² H. Mu,³ E. Muhammad,⁵⁰ F. Muheim,⁵² M. Mulder,⁷³ K. Müller,⁴⁴ C. H. Murphy,⁵⁷ D. Murray,⁵⁶ R. Murta,⁵⁵ P. Muzzetto,^{27,j} P. Naik,⁴⁸ T. Nakada,⁴³ R. Nandakumar,⁵¹ T. Nanut,⁴² I. Nasteva,² M. Needham,⁵² N. Neri,^{25,p} S. Neubert,⁷⁰ N. Neufeld,⁴² P. Neustroev,³⁸ R. Newcombe,⁵⁵ J. Nicolini,^{15,11} E. M. Niel,⁴³ S. Nieswand,¹⁴ N. Nikitin,³⁸ N. S. Nolte,⁵⁸ C. Normand,^{8,27,j} J. Novoa Fernandez,⁴⁰ C. Nunez,⁷⁸ A. Oblakowska-Mucha,³⁴ V. Obraztsov,³⁸ T. Oeser,¹⁴ D. P. O'Hanlon,⁴⁸ S. Okamura,^{21,d} R. Oldeman,^{27,j} F. Oliva,⁵² C. J. G. Onderwater,⁷³ R. H. O'Neil,⁵² J. M. Otalora Goicochea,² T. Ovsianikova,³⁸ P. Owen,⁴⁴ A. Oyanguren,⁴¹ O. Ozcelik,⁵² K. O. Padeken,⁷⁰ B. Pagare,⁵⁰ P. R. Pais,⁴² T. Pajero,⁵⁷ A. Palano,¹⁹ M. Palutan,²³ Y. Pan,⁵⁶ G. Panshin,³⁸ L. Paolucci,⁵⁰ A. Papanestis,⁵¹ M. Pappagallo,^{19,i} L. L. Pappalardo,^{21,d} C. Pappenheimer,⁵⁹ W. Parker,⁶⁰ C. Parkes,⁵⁶ B. Passalacqua,^{21,d} G. Passaleva,²² A. Pastore,¹⁹ M. Patel,⁵⁵ C. Patrignani,^{20,f} C. J. Pawley,⁷⁴ A. Pearce,⁴² A. Pellegrino,³² M. Pepe Altarelli,⁴² S. Perazzini,²⁰ D. Pereima,³⁸ A. Pereiro Castro,⁴⁰ P. Perret,⁹ M. Petric,⁵³ K. Petridis,⁴⁸ A. Petrolini,^{24,g} A. Petrov,³⁸ S. Petrucci,⁵² M. Petruzzo,²⁵ H. Pham,⁶² A. Philippov,³⁸ R. Piandani,⁶ L. Pica,^{29,c} M. Piccini,⁷² B. Pietrzyk,⁸ G. Pietrzyk,¹¹ M. Pili,⁵⁷ D. Pinci,³⁰ F. Pisani,⁴² M. Pizzichemi,^{26,42,e} V. Placinta,³⁷ J. Plews,⁴⁷ M. Plo Casasus,⁴⁰ F. Polci,^{13,42} M. Poli Lener,²³ M. Poliakova,⁶² A. Poluektov,¹⁰ N. Polukhina,³⁸ I. Polyakov,⁴² E. Polycarpo,² S. Ponce,⁴² D. Popov,^{6,42} S. Popov,³⁸ S. Poslavskii,³⁸ K. Prasanth,³⁵ L. Promberger,¹⁷ C. Prouve,⁴⁰ V. Pugatch,⁴⁶ V. Puill,¹¹ G. Punzi,^{29,t} H. R. Qi,³ W. Qian,⁶ N. Qin,³ S. Qu,³ R. Quagliani,⁴³ N. V. Raab,¹⁸ R. I. Rabadan Trejo,⁶ B. Rachwal,³⁴ J. H. Rademacker,⁴⁸ R. Rajagopalan,⁶² M. Rama,²⁹ M. Ramos Pernas,⁵⁰ M. S. Rangel,² F. Ratnikov,³⁸ G. Raven,^{33,42} M. Rebollo De Miguel,⁴¹ F. Redi,⁴² J. Reich,⁴⁸ F. Reiss,⁵⁶ C. Remon Alepuz,⁴¹ Z. Ren,³ P. K. Resmi,¹⁰ R. Ribatti,^{29,c} A. M. Ricci,²⁷ S. Ricciardi,⁵¹ K. Richardson,⁵⁸ M. Richardson-Slipper,⁵² K. Rinnert,⁵⁴ P. Robbe,¹¹ G. Robertson,⁵² A. B. Rodrigues,⁴³ E. Rodrigues,⁵⁴ E. Rodriguez Fernandez,⁴⁰ J. A. Rodriguez Lopez,⁶⁹ E. Rodriguez Rodriguez,⁴⁰ D. L. Rolf,⁴² A. Rollings,⁵⁷ P. Roloff,⁴² V. Romanovskiy,³⁸ M. Romero Lamas,⁴⁰ A. Romero Vidal,⁴⁰ J. D. Roth,^{78,a} M. Rotondo,²³ M. S. Rudolph,⁶² T. Ruf,⁴² R. A. Ruiz Fernandez,⁴⁰ J. Ruiz Vidal,⁴¹ A. Ryzhikov,³⁸ J. Ryzka,³⁴ J. J. Saborido Silva,⁴⁰ N. Sagidova,³⁸ N. Sahoo,⁴⁷ B. Saitta,^{27,j} M. Salomoni,⁴² C. Sanchez Gras,³² I. Sanderswood,⁴¹ R. Santacesaria,³⁰ C. Santamarina Rios,⁴⁰ M. Santimaria,²³ E. Santovetti,^{31,1} D. Saranin,³⁸ G. Sarpis,¹⁴ M. Sarpis,⁷⁰ A. Sarti,³⁰ C. Satriano,^{30,u} A. Satta,³¹ M. Saur,¹⁵ D. Savrina,³⁸ H. Sazak,⁹ L. G. Scantlebury Smead,⁵⁷ A. Scarabotto,¹³ S. Schael,¹⁴ S. Scherl,⁵⁴ M. Schiller,⁵³ H. Schindler,⁴² M. Schmelling,¹⁶ B. Schmidt,⁴² S. Schmitt,¹⁴ O. Schneider,⁴³ A. Schopper,⁴² M. Schubiger,³² S. Schulte,⁴³ M. H. Schune,¹¹ R. Schwemmer,⁴² B. Sciascia,^{23,42} A. Sciucchi,⁴² S. Sellam,⁴⁰ A. Semennikov,³⁸ M. Senghi Soares,³³ A. Sergi,^{24,g} N. Serra,⁴⁴ L. Sestini,²⁸ A. Seuthe,¹⁵ Y. Shang,⁵ D. M. Shangase,⁷⁸ M. Shapkin,³⁸ I. Shchemerov,³⁸ L. Shchutka,⁴³ T. Shears,⁵⁴ L. Shekhtman,³⁸ Z. Shen,⁵ S. Sheng,^{4,6} V. Shevchenko,³⁸ B. Shi,⁶ E. B. Shields,^{26,e} Y. Shimizu,¹¹ E. Shmanin,³⁸ R. Shorkin,³⁸ J. D. Shupperd,⁶² B. G. Siddi,^{21,d} R. Silva Coutinho,⁶² G. Simi,²⁸ S. Simone,^{19,i} M. Singla,⁶³ N. Skidmore,⁵⁶ R. Skuza,¹⁷ T. Skwarnicki,⁶² M. W. Slater,⁴⁷ J. C. Smallwood,⁵⁷ J. G. Smeaton,⁴⁹ E. Smith,⁴⁴ K. Smith,⁶¹ M. Smith,⁵⁵ A. Snoch,³² L. Soares Lavra,⁹ M. D. Sokoloff,⁵⁹ F. J. P. Soler,⁵³ A. Solomin,^{38,48} A. Solovev,³⁸ I. Solovyev,³⁸ R. Song,⁶³ F. L. Souza De Almeida,² B. Souza De Paula,² B. Spaan,^{15,a} E. Spadaro Norella,^{25,p} E. Spedicato,²⁰ E. Spiridenkov,³⁸ P. Spradlin,⁵³ V. Sriskaran,⁴² F. Stagni,⁴² M. Stahl,⁴² S. Stahl,⁴² S. Stanislaus,⁵⁷ E. N. Stein,⁴² O. Steinkamp,⁴⁴ O. Stenyakin,³⁸ H. Stevens,¹⁵ S. Stone,^{62,a} D. Strelakina,³⁸ F. Suljik,⁵⁷ J. Sun,²⁷ L. Sun,⁶⁸ Y. Sun,⁶⁰ P. Svihra,⁵⁶ P. N. Swallow,⁴⁷ K. Swientek,³⁴ A. Szabelski,³⁶ T. Szumlak,³⁴ M. Szymanski,⁴² Y. Tan,³ S. Taneja,⁵⁶ A. R. Tanner,⁴⁸ M. D. Tat,⁵⁷ A. Terentev,³⁸ F. Teubert,⁴² E. Thomas,⁴² D. J. D. Thompson,⁴⁷ K. A. Thomson,⁵⁴ H. Tilquin,⁵⁵ V. Tisserand,⁹ S. T'Jampens,⁸ M. Tobin,⁴ L. Tomassetti,^{21,d} G. Tonani,^{25,p} X. Tong,⁵ D. Torres Machado,¹ D. Y. Tou,³ S. M. Trilov,⁴⁸ C. Trippel,⁴³ G. Tuci,⁶ A. Tully,⁴³ N. Tuning,³² A. Ukleja,³⁶ D. J. Unverzagt,¹⁷ A. Usachov,³² A. Ustyuzhanin,³⁸ U. Uwer,¹⁷ A. Vagner,³⁸ V. Vagnoni,²⁰ A. Valassi,⁴² G. Valenti,²⁰ N. Valls Canudas,⁷⁶ M. van Beuzekom,³² M. Van Dijk,⁴³ H. Van Hecke,⁶¹ E. van Herwijnen,⁵⁵ C. B. Van Hulse,^{40,v} M. van Veghel,⁷³ R. Vazquez Gomez,³⁹ P. Vazquez Regueiro,⁴⁰ C. Vázquez Sierra,⁴² S. Vecchi,²¹ J. J. Velthuis,⁴⁸ M. Veltri,^{22,w} A. Venkateswaran,⁴³ M. Veronesi,³²

M. Vesterinen⁵⁰, D. Vieira⁵⁹, M. Vieites Diaz⁴³, X. Vilasis-Cardona⁷⁶, E. Vilella Figueras⁵⁴, A. Villa²⁰, P. Vincent¹³, F. C. Volle¹¹, D. vom Bruch¹⁰, A. Vorobyev³⁸, V. Vorobyev³⁸, N. Voropaev³⁸, K. Vos⁷⁴, C. Vrahas⁵², R. Waldi¹⁷, J. Walsh²⁹, G. Wan⁵, C. Wang¹⁷, G. Wang⁷, J. Wang⁵, J. Wang⁴, J. Wang³, J. Wang⁶⁸, M. Wang⁵, R. Wang⁴⁸, X. Wang⁶⁶, Y. Wang⁷, Z. Wang⁴⁴, Z. Wang³, Z. Wang⁶, J. A. Ward^{50,63}, N. K. Watson⁴⁷, D. Websdale⁵⁵, Y. Wei⁵, C. Weissler⁵⁸, B. D. C. Westhenry⁴⁸, D. J. White⁵⁶, M. Whitehead⁵³, A. R. Wiederhold⁵⁰, D. Wiedner¹⁵, G. Wilkinson⁵⁷, M. K. Wilkinson⁵⁹, I. Williams⁴⁹, M. Williams⁵⁸, M. R. J. Williams⁵², R. Williams⁴⁹, F. F. Wilson⁵¹, W. Wislicki³⁶, M. Witek³⁵, L. Witola¹⁷, C. P. Wong⁶¹, G. Wormser¹¹, S. A. Wotton⁴⁹, H. Wu⁶², J. Wu⁷, K. Wyllie⁴², Z. Xiang⁶, D. Xiao⁷, Y. Xie⁷, A. Xu⁵, J. Xu⁶, L. Xu³, L. Xu³, M. Xu⁵⁰, Q. Xu⁶, Z. Xu⁹, Z. Xu⁶, D. Yang³, S. Yang⁶, X. Yang⁵, Y. Yang⁶, Z. Yang⁵, Z. Yang⁶⁰, L. E. Yeomans⁵⁴, V. Yeroshenko¹¹, H. Yeung⁵⁶, H. Yin⁷, J. Yu⁶⁵, X. Yuan⁶², E. Zaffaroni⁴³, M. Zavertyaev¹⁶, M. Zdybal³⁵, O. Zenaiev⁴², M. Zeng³, C. Zhang⁵, D. Zhang⁷, L. Zhang³, S. Zhang⁶⁵, S. Zhang⁵, Y. Zhang⁵, Y. Zhang⁵⁷, A. Zharkova³⁸, A. Zhelezov¹⁷, Y. Zheng⁶, T. Zhou⁵, X. Zhou⁶, Y. Zhou⁶, V. Zhovkovska¹¹, X. Zhu³, X. Zhu⁷, Z. Zhu⁶, V. Zhukov^{14,38}, Q. Zou^{4,6}, S. Zucchelli^{20,f}, D. Zuliani²⁸ and G. Zunica⁵⁶

(LHCb Collaboration)

¹Centro Brasileiro de Pesquisas Físicas (CBPF), Rio de Janeiro, Brazil

²Universidade Federal do Rio de Janeiro (UFRJ), Rio de Janeiro, Brazil

³Center for High Energy Physics, Tsinghua University, Beijing, China

⁴Institute Of High Energy Physics (IHEP), Beijing, China

⁵School of Physics State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing, China

⁶University of Chinese Academy of Sciences, Beijing, China

⁷Institute of Particle Physics, Central China Normal University, Wuhan, Hubei, China

⁸Université Savoie Mont Blanc, CNRS, IN2P3-LAPP, Annecy, France

⁹Université Clermont Auvergne, CNRS/IN2P3, LPC, Clermont-Ferrand, France

¹⁰Aix Marseille Univ, CNRS/IN2P3, CPPM, Marseille, France

¹¹Université Paris-Saclay, CNRS/IN2P3, IJCLab, Orsay, France

¹²Laboratoire Leprince-Ringuet, CNRS/IN2P3, Ecole Polytechnique, Institut Polytechnique de Paris, Palaiseau, France

¹³LPNHE, Sorbonne Université, Paris Diderot Sorbonne Paris Cité, CNRS/IN2P3, Paris, France

¹⁴I. Physikalisches Institut, RWTH Aachen University, Aachen, Germany

¹⁵Fakultät Physik, Technische Universität Dortmund, Dortmund, Germany

¹⁶Max-Planck-Institut für Kernphysik (MPIK), Heidelberg, Germany

¹⁷Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany

¹⁸School of Physics, University College Dublin, Dublin, Ireland

¹⁹INFN Sezione di Bari, Bari, Italy

²⁰INFN Sezione di Bologna, Bologna, Italy

²¹INFN Sezione di Ferrara, Ferrara, Italy

²²INFN Sezione di Firenze, Firenze, Italy

²³INFN Laboratori Nazionali di Frascati, Frascati, Italy

²⁴INFN Sezione di Genova, Genova, Italy

²⁵INFN Sezione di Milano, Milano, Italy

²⁶INFN Sezione di Milano-Bicocca, Milano, Italy

²⁷INFN Sezione di Cagliari, Monserrato, Italy

²⁸Università degli Studi di Padova, Università e INFN, Padova, Padova, Italy

²⁹INFN Sezione di Pisa, Pisa, Italy

³⁰INFN Sezione di Roma La Sapienza, Roma, Italy

³¹INFN Sezione di Roma Tor Vergata, Roma, Italy

³²Nikhef National Institute for Subatomic Physics, Amsterdam, Netherlands

³³Nikhef National Institute for Subatomic Physics and VU University Amsterdam, Amsterdam, Netherlands

³⁴AGH - University of Science and Technology, Faculty of Physics and Applied Computer Science, Kraków, Poland

³⁵Henryk Niewodniczanski Institute of Nuclear Physics Polish Academy of Sciences, Kraków, Poland

³⁶National Center for Nuclear Research (NCBJ), Warsaw, Poland

³⁷Horia Hulubei National Institute of Physics and Nuclear Engineering, Bucharest-Magurele, Romania

³⁸Affiliated with an institute covered by a cooperation agreement with CERN

- ³⁹ICCUB, Universitat de Barcelona, Barcelona, Spain
- ⁴⁰Instituto Galego de Física de Altas Enerxías (IGFAE), Universidade de Santiago de Compostela, Santiago de Compostela, Spain
- ⁴¹Instituto de Física Corpuscular, Centro Mixto Universidad de Valencia - CSIC, Valencia, Spain
- ⁴²European Organization for Nuclear Research (CERN), Geneva, Switzerland
- ⁴³Institute of Physics, Ecole Polytechnique Fédérale de Lausanne (EPFL), Lausanne, Switzerland
- ⁴⁴Physik-Institut, Universität Zürich, Zürich, Switzerland
- ⁴⁵NSC Kharkiv Institute of Physics and Technology (NSC KIPT), Kharkiv, Ukraine
- ⁴⁶Institute for Nuclear Research of the National Academy of Sciences (KINR), Kyiv, Ukraine
- ⁴⁷University of Birmingham, Birmingham, United Kingdom
- ⁴⁸H.H. Wills Physics Laboratory, University of Bristol, Bristol, United Kingdom
- ⁴⁹Cavendish Laboratory, University of Cambridge, Cambridge, United Kingdom
- ⁵⁰Department of Physics, University of Warwick, Coventry, United Kingdom
- ⁵¹STFC Rutherford Appleton Laboratory, Didcot, United Kingdom
- ⁵²School of Physics and Astronomy, University of Edinburgh, Edinburgh, United Kingdom
- ⁵³School of Physics and Astronomy, University of Glasgow, Glasgow, United Kingdom
- ⁵⁴Oliver Lodge Laboratory, University of Liverpool, Liverpool, United Kingdom
- ⁵⁵Imperial College London, London, United Kingdom
- ⁵⁶Department of Physics and Astronomy, University of Manchester, Manchester, United Kingdom
- ⁵⁷Department of Physics, University of Oxford, Oxford, United Kingdom
- ⁵⁸Massachusetts Institute of Technology, Cambridge, Massachusetts, USA
- ⁵⁹University of Cincinnati, Cincinnati, Ohio, USA
- ⁶⁰University of Maryland, College Park, Maryland, USA
- ⁶¹Los Alamos National Laboratory (LANL), Los Alamos, New Mexico, USA
- ⁶²Syracuse University, Syracuse, New York, USA
- ⁶³School of Physics and Astronomy, Monash University, Melbourne, Australia
(associated with Institution Department of Physics, University of Warwick, Coventry, United Kingdom)
- ⁶⁴Pontifícia Universidade Católica do Rio de Janeiro (PUC-Rio), Rio de Janeiro, Brazil
(associated with Institution Universidade Federal do Rio de Janeiro (UFRJ), Rio de Janeiro, Brazil)
- ⁶⁵Physics and Micro Electronic College, Hunan University, Changsha City, China
(associated with Institution Institute of Particle Physics, Central China Normal University, Wuhan, Hubei, China)
- ⁶⁶Guangdong Provincial Key Laboratory of Nuclear Science, Guangdong-Hong Kong Joint Laboratory of Quantum Matter, Institute of Quantum Matter, South China Normal University, Guangzhou, China
(associated with Institution Center for High Energy Physics, Tsinghua University, Beijing, China)
- ⁶⁷Lanzhou University, Lanzhou, China
(associated with Institution Institute Of High Energy Physics (IHEP), Beijing, China)
- ⁶⁸School of Physics and Technology, Wuhan University, Wuhan, China
(associated with Institution Center for High Energy Physics, Tsinghua University, Beijing, China)
- ⁶⁹Departamento de Física, Universidad Nacional de Colombia, Bogota, Colombia
(associated with Institution LPNHE, Sorbonne Université, Paris Diderot Sorbonne Paris Cité, CNRS/IN2P3, Paris, France)
- ⁷⁰Universität Bonn - Helmholtz-Institut für Strahlen und Kernphysik, Bonn, Germany
(associated with Institution Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany)
- ⁷¹Eotvos Lorand University, Budapest, Hungary
(associated with Institution European Organization for Nuclear Research (CERN), Geneva, Switzerland)
- ⁷²INFN Sezione di Perugia, Perugia, Italy
(associated with Institution INFN Sezione di Ferrara, Ferrara, Italy)
- ⁷³Van Swinderen Institute, University of Groningen, Groningen, Netherlands
(associated with Institution Nikhef National Institute for Subatomic Physics, Amsterdam, Netherlands)
- ⁷⁴Universiteit Maastricht, Maastricht, Netherlands
(associated with Institution Nikhef National Institute for Subatomic Physics, Amsterdam, Netherlands)
- ⁷⁵Tadeusz Kosciuszko Cracow University of Technology, Cracow, Poland
(associated with Institution Henryk Niewodniczanski Institute of Nuclear Physics Polish Academy of Sciences, Kraków, Poland)
- ⁷⁶DS4DS, La Salle, Universitat Ramon Llull, Barcelona, Spain
(associated with Institution ICCUB, Universitat de Barcelona, Barcelona, Spain)
- ⁷⁷Department of Physics and Astronomy, Uppsala University, Uppsala, Sweden
(associated with Institution School of Physics and Astronomy, University of Glasgow, Glasgow, United Kingdom)
- ⁷⁸University of Michigan, Ann Arbor, Michigan, USA
(associated with Institution Syracuse University, Syracuse, New York, USA)

^aDeceased.

^bAlso at Università di Firenze, Firenze, Italy.

^cAlso at Scuola Normale Superiore, Pisa, Italy.

^dAlso at Università di Ferrara, Ferrara, Italy.

^eAlso at Università di Milano Bicocca, Milano, Italy.

^fAlso at Università di Bologna, Bologna, Italy.

^gAlso at Università di Genova, Genova, Italy.

^hAlso at Universidad Nacional Autónoma de Honduras, Tegucigalpa, Honduras.

ⁱAlso at Università di Bari, Bari, Italy.

^jAlso at Università di Cagliari, Cagliari, Italy.

^kAlso at Università di Perugia, Perugia, Italy.

^lAlso at Università di Roma Tor Vergata, Roma, Italy.

^mAlso at Universidade de Brasília, Brasília, Brazil.

ⁿAlso at Hangzhou Institute for Advanced Study, UCAS, Hangzhou, China.

^oAlso at Università di Siena, Siena, Italy.

^pAlso at Università degli Studi di Milano, Milano, Italy.

^qAlso at Central South U., Changsha, China.

^rAlso at Università di Padova, Padova, Italy.

^sAlso at Excellence Cluster ORIGINS, Munich, Germany.

^tAlso at Università di Pisa, Pisa, Italy.

^uAlso at Università della Basilicata, Potenza, Italy.

^vAlso at Universidad de Alcalá, Alcalá de Henares, Spain.

^wAlso at Università di Urbino, Urbino, Italy.