



Comparison of the Asthma Control Questionnaire and patient diaries in uncontrolled asthma

Copyright ©The authors 2023

This version is distributed under the terms of the Creative Commons Attribution Non-Commercial Licence 4.0. For commercial reproduction rights and permissions contact permissions@ersnet.org

Received: 15 Feb 2023

Accepted: 19 March 2023


To the Editor:

The Asthma Control Questionnaire (ACQ) is a validated tool commonly used to assess therapeutic efficacy in asthma clinical trials, as it is responsive to changes in asthma control [1]. However, it depends on accurate recall since patients are asked about their symptoms, activities and rescue medication use over the previous week. Even if patients can accurately remember this period, this also means that data are averaged. Guidelines recommend that patients with asthma are trained to keep track of their symptoms (e.g. using a diary [2]) and clinical trials often require patients to complete a diary twice daily.

TRIMARAN and TRIGGER were two phase III, double-blind, 52-week studies in patients with uncontrolled asthma (seven-item Asthma Control Questionnaire (ACQ-7) score ≥ 1.5) [3]. Both compared the efficacy and safety of single-inhaler triple therapy containing an extrafine formulation of the inhaled corticosteroid beclomethasone dipropionate (BDP), the long-acting β_2 -agonist formoterol fumarate (FF) and the long-acting muscarinic antagonist glycopyrronium *versus* BDP/FF, with TRIGGER including a third arm in which patients received open-label BDP/FF plus tiotropium. Patients completed an electronic diary (eDiary) twice daily (morning and evening), recording their symptom score and the number of puffs of rescue medication used during the day or night. The four daytime symptoms recorded were cough, wheeze, chest tightness and breathlessness, each rated by the patient from 0 (no symptoms) to 3 (severe symptoms); any night-time awakening due to asthma that resulted in rescue medication use was captured as “yes” or “no”. At baseline and during clinic visits, they also completed the ACQ-7, which includes five questions on symptoms, one question on rescue medication use and forced expiratory volume in 1 s (FEV₁).

We tested the hypothesis that ACQ and eDiary assessments of asthma control may differ. Using TRIMARAN and TRIGGER data, we compared ACQ-6 results (*i.e.* excluding FEV₁) at baseline and weeks 26 and 52 with those from the eDiary (using data collected in the 7 days prior to the clinic visit). The analyses only included patients with complete eDiary data over that period. Data from the eDiary were mapped to the Global Initiative for Asthma (GINA) symptom control assessment [2], allocating a score of 1 to each of the following: 1) mild, moderate or severe daytime asthma symptoms (*i.e.* if at least one of the daytime eDiary symptoms scored 1, 2 or 3) on >2 days; 2) any night-time awakenings due to asthma; 3) rescue medication use on >2 days; and 4) as there is no direct question on activity limitation in the eDiary, patients who had moderate or severe daytime asthma symptoms (*i.e.* at least one of the daytime eDiary symptoms scored 2 or 3, both defined as impacting activity) on >2 days received an additional score of 1. These scores were then summed, consistent with the GINA assessment of asthma control, with a total of 3–4 indicating uncontrolled asthma. An ACQ-6 score ≥ 1.5 indicates uncontrolled asthma [4, 5]. We used ACQ-6 rather than ACQ-7, since GINA does not include FEV₁ in the assessment of asthma control, although low FEV₁ is considered a risk factor for poor outcomes [2].

Overall, a higher proportion of patients in both studies met the ACQ-6 definition of uncontrolled asthma than met the eDiary definition at all three visits, with an improvement (*i.e.* reduction) from baseline in the proportion at weeks 26 and 52 with both measures (figure 1a). Given these results could be influenced by the cut-point selected to define “uncontrolled”, we also conducted a series of analyses on the actual ACQ-6 and eDiary scores, by mapping the two sets of scores to each other (controlled asthma: ACQ-6 scores ≤ 0.75 equalled eDiary total score of 0; poorly controlled: ACQ-6 >0.75 – ≤ 1.5 equalled eDiary 1–2;



Shareable abstract (@ERSpublications)

These *post hoc* analyses suggest that the Asthma Control Questionnaire and eDiary have different measurement properties, and it is therefore potentially valuable to include both in clinical trials
<https://bit.ly/3TIYCX>

Cite this article as: Singh D, Papi A, Brusselle G, *et al.* Comparison of the Asthma Control Questionnaire and patient diaries in uncontrolled asthma. *ERJ Open Res* 2023; 9: 00104-2023 [DOI: 10.1183/23120541.00104-2023].




FIGURE 1 a) Proportion of patients with uncontrolled asthma according to the electronic diary (eDiary) (total score 3 or 4) or six-item Asthma Control Questionnaire (ACQ-6) (score ≥ 1.5). b) Proportion of patients with ACQ-6 score less than, equal to or greater than eDiary total score. W: week.

uncontrolled: ACQ-6 > 1.5 equalled eDiary 3–4). At baseline, 3.3% and 2.3% of patients had a lower ACQ-6 score than eDiary total score, indicating that asthma was rated as more controlled when using ACQ-6 than the eDiary (increasing to 12.4–17.5% during follow-up) (figure 1b). However, although 52.9% and 48.4% had their asthma rated similarly using the two scales at baseline, 43.8% and 49.3% had a higher ACQ-6 score than eDiary total score, indicating that asthma was rated as less controlled when using the ACQ than the eDiary, consistent with the first set of analyses. In terms of patient baseline characteristics, other than ACQ-6 (where the mean baseline value was higher in the groups with ACQ-6 score higher than the eDiary score) there were no consistent differences between these categories that could explain these results.

Both the ACQ and the eDiary combine objective parameters (e.g. rescue medication use) with subjective questions. We therefore conducted a further set of analyses specific to rescue medication use, where the eDiary data over the 7-day period prior to the ACQ assessment were averaged to allow a direct comparison between the two. The data matched for the ACQ and eDiary for 52.0–63.9% of patients (with a better match at weeks 26 and 52 than at baseline) but for approximately one-third of the patients (29.9–39.0%), the ACQ results were again higher than the eDiary.

Taken together, these results suggest that the two tools have different measurement properties and that asthma control tends to be rated worse when using the ACQ than the eDiary. It is possible that they are

measuring different components of asthma (given the questions are phrased differently). However, the main advantage of the eDiary is the twice-daily completion, whereas the ACQ is only completed at study visits and asks patients to recall their asthma symptoms over the previous 7 days. The rescue medication data suggest that the 7-day recall period of the ACQ may result in patients overestimating the impact of asthma on their day-to-day life and that the data collected twice-daily in the eDiary are more precise due to the absence of the recall period. The main limitation of these analyses is that whereas the ACQ is a validated endpoint, having demonstrated good cross-sectional validity to both the Asthma Quality of Life Questionnaire and the Short Form-36 [1], the eDiary was developed specifically for these studies and is not similarly validated. In addition, although the eDiary questions were mapped to the GINA symptom control assessment, the questions are not a perfect match and in future, it would perhaps be useful to include a direct question on activity limitation.

In conclusion, given the differing results from these two tools, it is potentially valuable to include both in clinical trials. Inclusion of the (validated) ACQ facilitates comparisons between trials (including of the patients recruited into different trials). However, the results (particularly the rescue medication data) highlight a potential limitation of the ACQ: the recall period. A daily eDiary, which by its nature, does not require patients to accurately recall their asthma control over a prolonged period (although more burdensome), perhaps more accurately reflects a patient's level of asthma control. Importantly, a twice-daily eDiary is more able to capture day-to-day variability in symptoms, which may be of particular interest in a disease such as asthma that is highly variable.

Dave Singh¹, Alberto Papi², Guy Brusselle³, J. Christian Virchow⁴, Giorgio Walter Canonica⁵, Eva Topole⁵, Andrea Vele⁶ and George Georges⁶

¹Medicines Evaluation Unit, The University of Manchester, Manchester University NHS Foundation Trust, Manchester, UK. ²Respiratory Medicine, Department Translational Medicine, University of Ferrara, Ferrara, Italy. ³Department of Respiratory Medicine, Ghent University Hospital, Ghent, Belgium. ⁴Universitätsmedizin Rostock, Zentrum für Innere Medizin, Abteilung für Pneumologie, Rostock, Germany. ⁵Humanitas University and Research Hospital, Center of Personalized Medicine: Asthma and Allergy, Milan, Italy. ⁶Global Clinical Development, Chiesi Farmaceutici SpA, Parma, Italy.

Corresponding author: Dave Singh (dsingh@meu.org.uk)

Provenance: Submitted article, peer reviewed.

Acknowledgements: D. Singh is supported by the National Institute for Health Research Manchester Biomedical Research Centre.

Conflicts of interest: D. Singh reports consulting fees from Aerogen, AstraZeneca, Boehringer Ingelheim, Chiesi, Cipla, CSL Behring, Epiendo, Genentech, GlaxoSmithKline, Glenmark, Gossamerbio, Kinaset, Menarini, Novartis, Pulmatrix, Sanofi, Synairgen, Teva, Theravance and Verona, all outside the submitted work. A. Papi reports grant payments to his institution from Chiesi, AstraZeneca, GlaxoSmithKline, Boehringer Ingelheim, Pfizer, Teva and Sanofi, consultancy fees from Chiesi, AstraZeneca, GlaxoSmithKline, Novartis, Sanofi, Iqvia, Avillion and Elpen Pharmaceuticals, and payment or honoraria for lectures, presentations, speakers' bureaus, manuscript writing or educational events from Chiesi, AstraZeneca, GlaxoSmithKline, Boehringer Ingelheim, Menarini, Novartis, Zambon, Mundipharma, Teva, Sanofi, Edmond Pharma, Iqvia, MSD, Avillion and Elpen Pharmaceuticals, all outside the submitted work. G. Brusselle reports honoraria for lectures from AstraZeneca, Boehringer Ingelheim, Chiesi, GlaxoSmithKline, Novartis and Teva; he is a member of advisory boards for Amgen, AstraZeneca, Boehringer-Ingelheim, GlaxoSmithKline, Novartis, Sanofi/Regeneron and Teva. All of these declarations are outside the submitted work. J.C. Virchow reports consulting fees and payment or honoraria for lectures, presentations, speakers' bureaus, manuscript writing or educational events from Novartis, Chiesi, GlaxoSmithKline, AstraZeneca, TEVA, Menarini, Berlin-Chemie, Sanofi and Boehringer Ingelheim, support for attending meetings and/or travel from Sanofi and Boehringer Ingelheim, and participation on a data safety monitoring board or advisory board for Chiesi, all outside the submitted work. G.W. Canonica has no other relevant conflicts of interest to disclose. E. Topole is an employee of Chiesi Farmaceutici SpA, the sponsor of the studies. A. Vele is an employee of Chiesi Farmaceutici SpA, the sponsor of the studies. G. Georges is an employee of Chiesi USA, Inc.

Support statement: These studies were funded by Chiesi Farmaceutici SpA. We thank the investigators and patients at the investigative sites for their support of these studies. Writing support was provided by David Young

of Young Medical Communications and Consulting Ltd, funded by Chiesi Farmaceutici SpA. Funding information for this article has been deposited with the Crossref Funder Registry.

References

- 1 Juniper EF, O'Byrne PM, Guyatt GH, *et al.* Development and validation of a questionnaire to measure asthma control. *Eur Respir J* 1999; 14: 902–907.
- 2 Global Initiative for Asthma. Global strategy for asthma management and prevention, 2022. Available from: <https://ginasthma.org/gina-reports/>
- 3 Virchow JC, Kuna P, Paggiaro P, *et al.* Single inhaler extrafine triple therapy in uncontrolled asthma (TRIMARAN and TRIGGER): two double-blind, parallel-group, randomised, controlled phase 3 trials. *Lancet* 2019; 394: 1737–1749.
- 4 Juniper EF, Bousquet J, Abetz L, *et al.* Identifying “well-controlled” and “not well-controlled” asthma using the Asthma Control Questionnaire. *Respir Med* 2006; 100: 616–621.
- 5 Juniper EF, Svensson K, Mörk A-C, *et al.* Measurement properties and interpretation of three shortened versions of the asthma control questionnaire. *Respir Med* 2005; 99: 553–558.