

Article

Different Chromosome Segregation Patterns Coexist in the Tetraploid Adriatic Sturgeon *Acipenser naccarii*

Stefano Dalle Valle, Elisa Boscari, Simone Giulio Bordignon, Víctor Hugo Muñoz-Mora, Giorgio Bertorelle and Leonardo Congiu

Special Issue

Conservation Genetics of Sturgeons

Edited by

Dr. Elisa Boscari, Dr. Peilin Cheng, Dr. Leonardo Congiu, Prof. Dr. Arne Ludwig, Dr. Nikolai Mugue, Dr. Mohammad Pourkazemi and Dr. Kim Scribner



Article

Different Chromosome Segregation Patterns Coexist in the Tetraploid Adriatic Sturgeon *Acipenser naccarii*

Stefano Dalle Palle ^{1,†}, Elisa Boscarì ^{1,†} , Simone Giulio Bordignon ¹, Víctor Hugo Muñoz-Mora ²,
Giorgio Bertorelle ²  and Leonardo Congiu ^{1,*}

¹ Department of Biology, University of Padova, Via Ugo Bassi 58b, 35121 Padova, Italy

² Department of Life Sciences and Biotechnology, University of Ferrara, Via Luigi Borsari 46, 44121 Ferrara, Italy

* Correspondence: leonardo.congiu@unipd.it

† These authors contributed equally to this work.

Abstract: The Adriatic sturgeon, *Acipenser naccarii* (Bonaparte, 1836), is a critically endangered tetraploid endemism of the Adriatic region; it has been targeted, over the last 20 years, by different conservation programs based on controlled reproduction of captive breeders followed by the release of their juvenile offspring; its preservation would greatly benefit from the correct and coordinated management of the residual genetic variability available in the different captive stocks. In this sense, the setup of an efficient parental allocation procedure would allow identifying familiar groups and establishing informed breeding plans, effectively preserving genetic variation. However, being the species tetraploid, the analyses often deal with complex genome architecture and a preliminary evaluation of allele segregation patterns at different chromosomes is necessary to assess whether the species can be considered a pure tetraploid, as previously observed at some loci, or if a more complex situation is present. Here we study the segregation at 14 microsatellites loci in 12 familiar groups. Results support in different families the tetrasomic segregation pattern at 11 markers and the disomic segregation at three markers. The Adriatic sturgeon thus shows a mixed inheritance modality. In this species, and likely in other sturgeons, accurate knowledge of the loci used for paternity analysis is therefore required.

Keywords: acipenseridae; autotertraploid; allopolyploid; Adriatic sturgeon; disomic segregation; inheritance; microsatellites; tetrasomic segregation



Citation: Dalle Palle, S.; Boscarì, E.; Bordignon, S.G.; Muñoz-Mora, V.H.; Bertorelle, G.; Congiu, L. Different Chromosome Segregation Patterns Coexist in the Tetraploid Adriatic Sturgeon *Acipenser naccarii*. *Diversity* **2022**, *14*, 745. <https://doi.org/10.3390/d14090745>

Academic Editors: Michael Wink and Simon Blanchet

Received: 13 August 2022

Accepted: 7 September 2022

Published: 10 September 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Sturgeons are the most endangered group of species, according to the International Union for Conservation of Nature (IUCN, July 2021) (<http://www.iucnredlist.org>, accessed on 8 August 2022). For this reason, sturgeons are targeted by several conservation efforts that often include restocking programs with juveniles produced in captivity; these ex-situ conservation activities must necessarily be supported by studies aimed at preserving the residual genetic diversity through long-term breeding programs [1]. However, genetic analyses on sturgeons deal with complex genomes and various levels of ploidy, due to independent events of whole-genome duplication [2,3]. The first event of duplication took place in the Acipenseriformes' common ancestor starting from sixty chromosomes. Then, secondary events of duplication occurred in the Pacific and Atlantic clades leading to a total of 240 chromosomes [2]. Finally, a third event led to the unique number of 360 chromosomes observable in *Acipenser brevirostrum* (Lesueur, 1818) [4]. The number of chromosomes associated with the distinct levels of ploidy has been the subject of an extensive debate between two main positions. The first argues that, since the condition with 120 chromosomes results from a duplication event in the common ancestor, the species with 120 and 240 chromosomes must be considered tetraploid and octaploid, respectively. The second position, taking into account the functional reduction of ploidy that follows whole

genome duplications, attributes to the two groups a condition of diploidy and tetraploidy, respectively [5]. The two views use different criteria to define the nominal ploidy, the number of duplications and the functional activity of genes, respectively, and are both correct and fully compatible [3].

The Adriatic sturgeon, *Acipenser naccarii* (Bonaparte, 1836), a species with 240 chromosomes, is in general considered a functional tetraploid based on the analyses of 28S and 5S rDNA through in situ hybridization [6]; this is also confirmed by most of the microsatellites analysed in this species which mostly show up to 4 alleles per individuals. However, some loci consistently showed more than 4 alleles in many individuals [7]; this could be due to the duplication of the region of such microsatellites but, a still incomplete process of functional reduction following genome duplication cannot be excluded as already proposed also for other sturgeon species [8,9].

For these reasons, loci to be used for applied purposes, such as parental allocation and kinship analysis, should be carefully selected and their functional ploidy should be preliminarily assessed. Moreover, these applications require a careful preliminary investigation also on the segregation modalities which, in tetraploids, can be of different types. In fact, polyploidization can originate through the fusion of unreduced gametes at intraspecific or interspecific levels, leading to two types of conditions in tetraploids, called autotetraploidy and allotetraploidy, respectively. The origins of polyploidization have important implications on the segregation patterns of the alleles within gametes. In complete autotetraploids, there are always four homologous chromosomes, and random pairs of bivalents and quadrivalents are possible during meiosis (see Figure 1 for graphical support) [10]; this condition leads to tetrasomic inheritance, which means that all allelic combinations within gametes are possible; this is not the case in complete allotetraploids, where each tetrad is composed of two sets of homeologous chromosomes, originating from the two parental species. In this situation, the homeologous chromosomes do not form pairs, leading to disomic inheritance, where only four out of six allelic combinations are possible [10]; these are two extreme cases and intermediate conditions are observable when the chromosomes have different degrees of preferential pairing. Indeed, the inheritance may shift from tetrasomic to disomic or vice versa [10]. For example, in autotetraploids, fertility and karyotype stability can be negatively impacted by imperfect multivalent pairing, thus promoting diploidization and consequent shifting to disomy; on the contrary, in allotetraploids, the homeologous chromosomes from two distinct parental species could maintain some degree of genetic affinity permitting the competition with the homologous pair during meiotic interactions with a certain degree of tetrasomic inheritance [11]. Understanding the mechanisms of chromosomal segregation in a tetraploid species can have important conservation implications, for example when the development of parental allocation methods is required.

In the Adriatic sturgeon, allele segregation was previously investigated at only 7 loci using microsatellite markers [12]. All loci showed a tetrasomic inheritance pattern, pointing at the probable autoployploidization origin of this species. However, as secondary differentiation of some homologous chromosomes cannot be excluded and different segregation patterns can be followed by different chromosomes, we took advantage of the recent availability of new complete family groups and new isolated and tested microsatellite loci not yet explored, to provide a deeper insight into the mode of chromosome segregation in this species.

This is a key step not only to have a better understanding of the karyotype in the Adriatic sturgeon but also for the correct interpretation of the genetic analysis and parental allocation which in this species can have multiple applications. Firstly, the distinction of extremely rare individuals of wild origin from released ones. Secondly, the identification of groups of siblings existing in the different farms by assigning everyone to his pair of parents of the F0 generation. Finally, knowing the segregation patterns is crucial when the generation of virtual genotypes starting from observed genetic profiles is needed [13,14].




Figure 1. Expected segregation patterns under pure disomy (a) or tetrasomy (b), respectively expected in Autotetraploid and Allotetraploid genomes.

Thanks to the possibility of making 12 distinct crosses and raising their progeny separately within the project ENDEMIXIT (<https://endemixit.com/>, accessed on 8 August 2022), many new informative familiar groups became available for the analysis of segregation patterns at a much higher number of loci than those available in the past. The main purpose of the present study is therefore to exhaustively describe the modalities of microsatellite alleles inheritance in the Adriatic sturgeon with the following objectives: (a) verifying if the inferred pure tetrasomy is confirmed on a high number of loci, (b) providing a significant contribution to the management of the residual genetic diversity of this critically endangered species, and (c) shedding light on the functional ploidy level across its genome.

2. Materials and Methods

2.1. Samples and DNA Purification

Most samples analyzed in the present study were obtained from the aquaculture plant Storione Ticino (Cassolnovo, Italy). The reproduction of six females (F2, F4, F5, F6, F7, and F8) and six males (M2, M4, M5, M6, M7, and M8) was performed in different combinations. For each parent, a fin clip was collected for genotyping and, for each cross, the progeny was reared in captivity and spontaneously dead animals were collected and stored in ethanol. Moreover, five familiar groups (Nacc7 ♀ × Nacc5 ♂, Nacc8 ♀ × Nacc31 ♂, Nacc19 ♀ × Nacc17 ♂, Nacc19 ♀ × Nacc30 ♂, Nacc28 ♀ × Nacc23 ♂) derived from crosses performed in the past and for which fingerlings were available were also used. In total, 21 adult parents (12 parent pairs) and 376 fingerlings (with about 30 individuals per family) were collected (Table 1).

Genomic DNA was extracted from breeder’s fin clips (10–100 mg) and from offspring’s muscular tissue, using Euroclone spinNAker Universal Genomic DNA mini kit (Euroclone) and stored at −20 °C till their processing for microsatellite analysis.

2.2. Selection of Loci and Genotyping

Loci analysed in the present study were selected based on the possibility of unambiguously tracing the genetic contribution of at least one of the two parents in at least one of the

available families. For this reason, all loci in which there were no complete heterozygotes or there were individuals with more than four alleles were discarded. In fact, it is known that the Adriatic sturgeon presents a minority of loci at which more than four alleles were can be observed in different individuals; thus, segregation anomalies cannot be detected [7]; these extra numerary alleles could originate from duplication or from a locally unreduced octaploid condition. In fact, we recall that the Adriatic sturgeon is functionally a tetraploid but evolutionarily it is an octaploid [3]. Nevertheless, segregation anomalies were also observed in some individuals of each family that were accordingly excluded from the analysis.

Table 1. Screening of microsatellite loci.

Locus	Ref	Ta	Size	Family	Parents	Genotypes	Genotyped Fingerling (Y/N)	#
LS-39	15	52 °C	116–155	Not informative	-	-	N	-
AfuG113	16	Td	326–364	Not informative	-	-	N	-
AfuG132	16	61 °C	259–346	F5 ♀ × M6 ♂	F5 ♀ M6 ♂	293/309/318/330 313/318/322	Y	32
				F6 ♀ × M7 ♂	F6 ♀ M7 ♂	313/318/342 293/309/318/322	Y	32
				F8 ♀ × M2 ♂	F8 ♀ M2 ♂	304/318/322 313/338/342/346	Y	30
AfuG41	16	58 °C	156–198	F8 ♀ × M2 ♂	F8 ♀ M2 ♂	212/234/293 203/255/259/285	Y	30
				F2 ♀ × M4 ♂	F2 ♀ M4 ♂	247/255/259/285 212/255/293/305	Y	30
				F4 ♀ × M4 ♂	F4 ♀ M4 ♂	212/217/242/259 212/255/293/305	Y	30
				F6 ♀ × M7 ♂	F6 ♀ M7 ♂	212/242/259/278 203/225/229/259	Y	32
				Nacc7 ♀ × Nacc5 ♂	Nacc7 ♀ Nacc5 ♂	225/229/252/259 203/225/247/305	Y	32
				Nacc8 ♀ × Nacc31 ♂	Nacc8 ♀ Nacc31 ♂	212/242/293/305 203/234/305	Y	32
An20 *	17	62 °C	159–213	F7 ♀ × M8 ♂	F7 ♀ M8 ♂	160/164/194 160/172/182/186	Y	30
Anac-15214	7	61 °C	259–285	Not informative	-	-	N	-
Anac-2589	7	63 °C	224–294	Not informative	-	-	N	-
Anac-6784	7	62 °C	311–346	Nacc19 ♀ × Nacc17 ♂	Nacc19 ♀ Nacc17 ♂	311/322/330/334 322/326	Y	32
Anac-3133	7	56 °C	164–178	F7 ♀ × M8 ♂	F7 ♀ M8 ♂	164/174 164/166/170/172	Y	30
AnacA6 *	18	62 °C	289–313	F5 ♀ × M6 ♂	F5 ♀ M6 ♂	307/313 293/297/301/307	Y	32
				F6 ♀ × M7 ♂	F6 ♀ M7 ♂	148/150 138/144/148/162	Y	32
AnacB11	18	60 °C	132–162	Nacc19 ♀ × Nacc17 ♂	Nacc19 ♀ Nacc17 ♂	132/136/138/144 132/150/162	Y	32
AnacB7	18	60 °C	152–198	F6 ♀ × M7 ♂	F6 ♀ M7 ♂	166/172/174/176 154/156/164	Y	32
				Nacc19 ♀ × Nacc30 ♂	Nacc19 ♀ Nacc30 ♂	156/166/174 154/170/174/176	Y	32
AnacC11	18	50 °C	167–193	Not informative	-	-	N	-
AnacE4 *	18	58 °C	326–354	F5 ♀ × M6 ♂	F5 ♀ M6 ♂	332/340/346/354 336/346	Y	32
				F4 ♀ × M5 ♂	F4 ♀ M5 ♂	123/127/131/139 131/135/155	Y	30
AoxD161	19	60 °C	111–155	F8 ♀ × M2 ♂	F8 ♀ M2 ♂	123/131/135/139 127/131	Y	30
				Nacc7 ♀ × Nacc5 ♂	Nacc7 ♀ Nacc5 ♂	123/127/131/143 131/135/139	Y	32

Table 1. Cont.

Locus	Ref	Ta	Size	Family	Parents	Genotypes	Genotyped Fingerling (Y/N)	#
AoxD234 *	19	52 °C	215–275	F2 ♀ × M4 ♂	F2 ♀ M4 ♂	219/223/243/255 227/243	Y	30
				F4 ♀ × M5 ♂	F4 ♀ M5 ♂	227/243/247/251 235/239/255/259	Y	30
				Nacc28 ♀ × Nacc23 ♂	Nacc28 ♀ Nacc23 ♂	219/243/247/263 239/243/251/255	Y	24
				F6 ♀ × M7 ♂	F6 ♀ M7 ♂	227/243/247/255 219/239/247	Y	32
AoxD241 *	19	57 °C	156–196	F7 ♀ × M8 ♂	F7 ♀ M8 ♂	168/176/180 164/172/176/184	Y	30
AoxD64 *	19	60 °C	216–260	Not informative	-	-	N	-
Spl-120	20	55 °C	263–303	Not informative	-	-	N	-
Spl-163	20	63 °C	166–233	F2 ♀ × M4 ♂	F2 ♀ M4 ♂	207/215/220 166/215/224/229	Y	30
				F7 ♀ × M8 ♂	F7 ♀ M8 ♂	232/240/294 218/236/271/273	Y	30
				F4 ♀ × M5 ♂	F4 ♀ M5 ♂	209/232/257/273 218/232/279/282	Y	30
				F6 ♀ × M7 ♂	F6 ♀ M7 ♂	227/236/271/286 240/245/264/269	Y	32
				Nacc19 ♀ × Nacc17 ♂	Nacc19 ♀ Nacc17 ♂	218/227/269/294 227/240/286/314	Y	32
				Nacc7 ♀ × Nacc5 ♂	Nacc7 ♀ Nacc5 ♂	232/245/264/294 214/240/273/279	Y	32
				Nacc8 ♀ × Nacc31 ♂	Nacc8 ♀ Nacc31 ♂	209/236/271/279 218/249/273/279	Y	32
Spl-168	10	63 °C	200–314	F7 ♀ × M8 ♂	F7 ♀ M8 ♂	232/240/294 218/236/271/273	Y	30
				F4 ♀ × M5 ♂	F4 ♀ M5 ♂	209/232/257/273 218/232/279/282	Y	30
				F6 ♀ × M7 ♂	F6 ♀ M7 ♂	227/236/271/286 240/245/264/269	Y	32
				Nacc19 ♀ × Nacc17 ♂	Nacc19 ♀ Nacc17 ♂	218/227/269/294 227/240/286/314	Y	32
				Nacc7 ♀ × Nacc5 ♂	Nacc7 ♀ Nacc5 ♂	232/245/264/294 214/240/273/279	Y	32
				Nacc8 ♀ × Nacc31 ♂	Nacc8 ♀ Nacc31 ♂	209/236/271/279 218/249/273/279	Y	32

Loci were used for genotyping of breeders for the selection of informative family/locus combinations for which progeny was also processed. References, annealing temperatures used, and size are reported. For each informative family/locus combination, the genotypes of each parent at such loci are also reported and the parent's genotypes for which the allele segregation was followed in the progeny is in bold. The total number of fingerlings processed for each informative family is reported in the last column. * Loci for which the segregation inheritance pattern was already explored in other familiar groups of the same species [12]. Y = Yes, N = Not processed. # = number of analyzed individuals.

All breeders were genotyped at 21 microsatellite loci (Table 1) [7,15–20] to select the informative Family/Locus combinations for the assessment of chromosomal segregation. For each selected combination, the progeny was also amplified and genotyped. Tracking segregation in the progeny requires the satisfaction of some features, such as (i) the complete heterozygosity (four different alleles) of at least one parent to ensure that each allele can unambiguously mark the segregation of its own chromosome and (ii) no more than an allele shared by the two parents to avoid ambiguity in following the alleles transmission to the progeny [12].

A total of 33 Family/Locus combinations were finally selected and approximately 30 fingerlings each were genotyped (Table 1). In 22 out of 33 case studies only one breeder of the parent pair was completely heterozygote and therefore informative to follow the segregation of the alleles, while for the remnant 11 Family/Locus combinations the segregation pattern was followed for both breeders of the parent pair as both they showed completely heterozygous genotypes with a single allele shared at most (Table 1).

Microsatellite loci were amplified from genomic DNA in a 10 µL reaction: 1X Master Mix Buffer (QIAGEN), 0.2 µM of each primer and about 50 ng of genomic DNA. Amplifications were performed on SimpliAmp Thermal Cycler (Thermo Fisher Scientific, Bologna, Italy). Amplifications were checked on 1.8% agarose gel in TBE1X stained with GelRed (BIOTIUM, Fremont, Canada, GelRed™ Nucleic Acid Stain). Genotyping was performed on ABI PRISM 310 Genetic Analyzer (external service, BMR Genomics, Padova, Italy). For microsatellite scoring the software GeneMarker Version 1.95 (SoftGenetics LLC, State College, PA, USA) was used.

2.3. Data Analyses

Scoring was performed by two operators, independently, with GeneMarker Version 1.95 (SoftGenetics LLC). The final dataset was carefully checked, and each individual was controlled for perfect Mendelian inheritance within his Family/Locus combination under the assumption that the observed genotypes at a given locus should be compatible with the inheritance of two allele copies from each parent. Individuals that for different possible reasons discussed later did not match this assumption were discarded. Since discordances with the above-accepted criterion can be attributed to anomalies in chromosomal segregation in the gametes of one parent, which are independent of what happened in the mating partner's gametes, it was decided not to discard those animals whose segregation anomaly was due to a problem in the uninformative parent. Limiting our observations to those animals that met standard inheritance of two out of four alleles, we consciously excluded possible genotypes that were not necessarily derived from segregation anomalies such as the case of double reduction, a phenomenon widely observed in autotetraploid plants in which the four homolog chromosomes may form multivalents [21,22] in which identical alleles carried on the sister chromatids may enter the same gamete with consequent segregation of two copies of the same allele even if it was present in a single copy in the parental genome [23]. We have decided to limit ambiguities by disregarding this phenomenon and excluding all the individuals that could be compatible with it focusing on animals in which alleles were unambiguously traceable.

As proposed by Stift et al. (2008) [10], Likelihood Ratio Tests (LRT) with 1 df were applied to compare the null model of tetrasomy with the other alternative models intermediate between disomic and tetrasomic. For each parent-locus combination and each alternative inheritance model, the log-likelihood was estimated from constrained nonlinear regression models, using SPSS syntax as reported in the original reference [10]. The Sequential Bonferroni correction [24] was applied to adjust significance levels ($p < 0.05$) for multiple comparisons across loci and families.

3. Results

From 22 to 32 individuals for each family-locus combination were successfully genotyped. In some cases, a few animals were discarded for unreliable profiles or, after the allele scoring, due to the presence of segregation anomalies of alleles inherited from the informative parent (Table 2).

At 11 (AfuG132, AfuG41, An20, Anac6784, Anac3133, AnacA6, AnacB11, AnacB7, AnacE4, AoxD234, AoxD241) out of 14 informative loci the LRT did not lead to the rejection of the null model of inheritance thus supporting the hypothesis of tetrasomy. At almost all these loci, indeed, all six allele combinations were observed in almost all tested families (Figure 1a, Table 2, Figure A1 of Appendix A). The only exceptions were the loci AoxD234 and Anac-3133 at a single family each showing only five allele combinations. However, in both cases, the strict disomic inheritance was excluded after correction for multiple tests (Table 2). In these two families, the sixth combination is expected to be detectable by increasing the sample size. Five (An20, AnacA6, AnacE4, AoxD234, and AoxD241) of the 11 loci showing tetrasomic inheritance were already tested to assess the inheritance pattern in different familiar groups of Adriatic sturgeon [12] and their tetrasomy has been here confirmed. Specifically, the locus AnacA6 here analysed at a single family showed only four allele combinations but, missing combinations are not compatible with disomic inheritance modality and even in this case the null model was not rejected.

On the contrary, a significant rejection of the null model was observed at three loci, Spl163, AoxD161, and Spl168, analysed respectively at one, three and six families and never analysed before in other studies. In some cases, the analysed families have both parents informative for segregation and agree in suggesting a disomic mode of inheritance (Table 2, Figure 1b, Figure A1 Appendix A). The disomic model fitted the observed allele combination frequencies significantly better than the null model, and the parameter τ equal to zero indicates a full disomic inheritance at almost all Family/Locus combinations. The

only two exceptions were observed in the segregation of the alleles of female F8 at locus AoxD161 and male Nacc31 at locus Spl168 (Table 2). In these two cases, five combinations of alleles were present, the parameter τ assumed a very low value confirming a strong degree of preferential pairing during meiosis but the rejection of the null model was not significant after the Bonferroni correction, thus suggesting possible imperfect preferential pairing.

Table 2. Likelihood Ratio Test.

Locus	Family	Informative Parent	N (Nd)	Null Model (T = 1) Likelihood Obs	Best Intermediate Model			Model Comparison: LRT	p-Values
					Pairing Alleles	T	Likelih-ood Obs		
Afug132	F5 ♀ × M6 ♂	F5 ♀	22 (8)	39.42	AC/BD	0.82	39.23	0.19	0.3322
	F6 ♀ × M7 ♂	M7 ♂	29 (1)	51.96	AD/BC	0.83	51.74	0.22	0.3185
	F8 ♀ × M2 ♂	M2 ♂	29 (1)	51.96	AB/CD	0.83	51.74	0.22	0.3185
Afug41	F8 ♀ × M2 ♂	M2 ♂	27 (3)	48.38	AC/BD	0.89	48.29	0.09	0.3853
	F2 ♀ × M4 ♂	F2 ♀	27 (3)	48.38	AB/CD	0.78	48.03	0.35	0.2776
		M4 ♂	27 (3)	48.38	AB/CD	0.44	45.98	2.39	0.0609
	F4 ♀ × M4 ♂	F4 ♀	29 (1)	51.96	AC/BD and AD/BC	0.93	51.93	0.03	0.4259
		M4 ♂	29 (1)	51.96	AD/BC	0.83	51.74	0.22	0.3185
	F6 ♀ × M7 ♂	F6 ♀	28 (2)	50.17	AD/BC	0.86	50.02	0.15	0.3509
M7 ♂		28 (2)	50.17	AB/CD	0.86	50.02	0.15	0.3509	
Nacc7 ♀ × Nacc5 ♂	Nacc7 ♀	28 (4)	50.17	AD/BC	0.64	49.21	0.96	0.1631	
	Nacc5 ♂	28 (4)	50.17	AD/BC	0.75	49.71	0.46	0.2489	
Nacc8 ♀ × Nacc31 ♂	Nacc8 ♀	31 (1)	55.54	AD/BC	0.77	55.13	0.41	0.2603	
An20 *	F7 ♀ × M8 ♂	M8 ♂	30 (0)	53.75	AD/BC	0.70	53.03	0.72	0.1984
Anac6784	Nacc19 ♀ × Nacc17 ♂	Nacc19 ♀	30 (2)	53.75	AC/BD and AD/BC	0.91	53.68	0.08	0.3912
Anac3133	F7 ♀ × M8 ♂	M8 ♂	29 (1)	51.96	AB/CD	0.41	49.06	2.90	0.0444 ^a
AnacA6 *	F5 ♀ × M6 ♂	M6 ♂	29 (1)	51.96	AB/CD and AC/BD	0.72	51.38	0.58	0.2225
	F6 ♀ × M7 ♂	M7 ♂	26 (4)	46.59	AB/CD	0.58	45.31	1.28	0.1290
AnacB11	Nacc19 ♀ × Nacc17 ♂	Nacc19 ♀	26 (6)	46.59	AB/CD	0.81	46.34	0.25	0.3088
AnacB7	F6 ♀ × M7 ♂	F6 ♀	27 (3)	48.38	AB/CD	0.67	47.57	0.80	0.1849
	Nacc19 ♀ × Nacc30 ♂	Nacc30 ♂	32 (0)	57.34	AD/BC	0.94	57.30	0.03	0.4295
AnacE4 *	F5 ♀ × M6 ♂	F5 ♀	28 (2)	50.17	AB/CD and AD/BC	0.96	50.16	0.01	0.4622
AoxD161	F4 ♀ × M5 ♂	F4 ♀	29 (1)	51.96	AB/CD	0.00	40.20	11.76	0.0003
	F8 ♀ × M2 ♂	F8 ♀	30 (0)	53.75	AC/BD	0.10	45.28	8.47	0.0018 ^a
	Nacc7 ♀ × Nacc5 ♂	Nacc7 ♀	32 (0)	57.34	AD/BC	0.00	44.36	12.97	0.0002
AoxD234 *	F2 ♀ × M4 ♂	F2 ♀	27 (3)	48.38	AD/BC	0.89	48.29	0.09	0.3853
	F4 ♀ × M5 ♂	F4 ♀	28 (2)	50.17	AD/BC	0.86	50.02	0.15	0.3509
		M5 ♂	28 (2)	50.17	AD/BC	0.64	49.21	0.96	0.1631
	Nacc28 ♀ × Nacc23 ♂	Nacc28 ♀	23 (1)	41.21	AB/CD	0.78	40.93	0.28	0.2972
		Nacc23 ♂	23 (1)	41.21	AC/BD	0.26	37.29	3.92	0.0239 ^a
F6 ♀ × M7 ♂	F6 ♀	29 (1)	51.96	AC/BD	0.52	50.07	1.89	0.0844	
AoxD241 *	F7 ♀ × M8 ♂	M8 ♂	29 (1)	51.96	AC/BD	0.72	51.38	0.58	0.2225
Spl163	F2 ♀ × M4 ♂	M4 ♂	28 (2)	50.17	AC/BD	0.00	38.82	11.35	0.0004
	F7 ♀ × M8 ♂	M8 ♂	26 (4)	46.59	AB/CD	0.00	36.04	10.54	0.0006
Spl168	F4 ♀ × M5 ♂	F4 ♀	30 (0)	53.75	AB/CD	0.00	41.59	12.16	0.0002
		M5 ♂	30 (0)	53.75	AB/CD	0.00	41.59	12.16	0.0002
	F6 ♀ × M7 ♂	F6 ♀	29 (1)	51.96	AB/CD	0.00	40.20	11.76	0.0003
		M7 ♂	29 (1)	51.96	AB/CD	0.00	40.20	11.76	0.0003
	Nacc19 ♀ × Nacc17 ♂	Nacc19 ♀	23 (9)	41.21	AB/CD	0.00	31.88	9.33	0.0011
		Nacc17 ♂	23 (9)	41.21	AB/CD	0.00	31.88	9.33	0.0011
Nacc17 ♂		23 (9)	41.21	AB/CD	0.00	31.88	9.33	0.0011	
Nacc7 ♀ × Nacc5 ♂	Nacc7 ♀	32 (0)	57.34	AB/CD	0.00	44.36	12.97	0.0002	
	Nacc5 ♂	32 (0)	57.34	AB/CD	0.00	44.36	12.97	0.0002	

Table 2. Cont.

Locus	Family	Informative Parent	N (Nd)	Null Model (T = 1) Likelihood Obs	Best Intermediate Model			Model Comparison: LRT	p-Values
					Pairing Alleles	T	Likelihood Obs		
	Nacc8 ♀ × Nacc31 ♂	Nacc8 ♀	29 (3)	51.96	AB/CD	0.00	40.20	11.76	0.0003
		Nacc31 ♂	29 (3)	51.96	AB/CD	0.10	43.86	8.10	0.0022 ^a

Results of Likelihood Ratio Test between the null model of tetrasomy (TAU = 1) and the best fitting intermediate one (TAU estimated). Significant *p*-values (after Bonferroni correction) that reject the null hypothesis of tetrasomy are highlighted in grey. ^a not significant values after Bonferroni correction for multiple tests. * Loci for which the segregation inheritance pattern was already explored in other familiar groups of the same species [12].

Expected allele combinations in tetrasomic and disomic mode of inheritance in tetraploids. (a) Observed allele combinations inherited from a complete heterozygote parent in autotetraploids in which tetrads are generated during meiosis and random segregation of allele pairs is expected. In brackets, the number of individuals carrying the relative allele combination is reported. (b) Observed allele combinations inherited from a complete heterozygote parent in allotetraploids in which a preferential pair between homologous chromosomes occurs. Only four possible gametes are expected. Alleles of the parent for which the segregation is reported in each table are marked in bold and by a letter used to indicate the observed combinations. Complete tables for all loci are reported in Appendix A. (Created with BioRender.com, accessed on 8 August 2022).

4. Discussion

The segregation pattern observed in the present study at most loci indicates that the Adriatic sturgeon can be considered predominantly tetraploid with a tetrasomic inheritance pattern. The presence of three disomic loci, however, indicates that the functional diversification process is at different stages in different parts of the genome, with some regions possibly still octaploid, most tetraploid, and some others in which the degree of divergence has gone up to a condition of double diploidy. We also observed the presence of two loci with a marked tendency to disomy with a few unexpected allele combinations, pointing to a possible imperfect preferential pairing expected at intermediate stages of functional diploidization [10,11]; this coexistence of different ploidy levels was previously described in other organisms. In plants, for example, the co-existence of different segregation patterns (e.g., tetrasomic and disomic) with different intermediate degrees of preferential pairing among chromosomes provides evidence of a process of functional reduction of ploidy which reasonably cannot occur simultaneously throughout the genome, but which is the result of a progressive differentiation [11,25].

As for the sturgeons, in other species the presence of different degrees of ploidy has been deduced based on the maximum number of alleles per individual present at the different loci [16] and, in some cases, the Mendelian transmission in the progeny has also been verified [15,26,27]. However, the selection of the cross/locus combinations to allow the traceability of every single allele and consequently to distinguish between the different modes of tetraploid segregation (e.g., disomic, tetrasomic or intermediate) has been conducted to our knowledge only on the Adriatic sturgeon.

The evidence that the level of homology between chromosomes within the Adriatic sturgeon genome is likely to vary from chromosome to chromosome and that different parts of the genome may consequently have different degrees of functional ploidy (2 to 8) should be considered when characterizing the genome of this species and probably of sturgeons in general. Genome assembly procedures must contemplate the possibility that different regions are present with a variable number of copies.

Whatever the evolutionary origin and implications of our results, which can be better investigated only when the genome of this species will be available, the identification of different inheritance pattern at different markers may have practical consequences for the conservation of the Adriatic sturgeon. In fact, our findings suggest that before developing parental allocation methods, a preliminary analysis of the loci used is recommended; this

would ease the reconstruction of individual genealogies of animals kept as captive breeders and the reallocation of any individual recaptured after release. Another interesting and relatively unexplored aspect of sturgeon *ex situ* conservation in which a clear knowledge of the inheritance patterns at different loci could be useful is the monitoring of the genomic impact of breeding protocols. Captive breeding is usually performed following standardized protocols of hormonal induction of egg and milt release and fertilization is done in an excess of sperms. Then, the resulting viable progeny is released without verifying if the procedure used had some effect on their genomic asset, for example, by inducing aneuploidies, which are a common phenomenon in some captive sturgeon stocks [28]. The release in nature of genetically anomalous individuals should be avoided as the consequences that this can have on the following generations and on their reproductive efficiency is unknown and, in the case of animals with long generation times, could be revealed after decades. Random screening of familiar groups to verify the correct parents-to-progeny segregation at both disomic and tetrasomic loci could significantly contribute to reducing the potential impact of genomic anomalies on natural populations.

5. Conclusions

Thanks to the availability of some family groups not previously analyzed, it was possible to better investigate the patterns of chromosomal segregation in the polyploid Adriatic sturgeon. The picture that emerged is that of an extremely dynamic genome in which it is possible to find the co-existence of regions with different degrees of ploidy, some of which retain the legacy of ancient duplications and others show a dynamic reduction of functional ploidy; this pattern is probably shared with other polyploid sturgeons in which different patterns of segregation have been observed [27], but it is not known whether the same genomic regions are involved.

Additional studies are required to better characterize the distribution of ploidy across the genome; this will likely contribute to explaining some specific features of the sturgeons, such as the ability of species with different degrees of ploidy to hybridize and produce viable offspring [29]; this extreme genomic plasticity could somehow be linked to a high degree of genomic redundancy; it would also be very interesting to verify whether the regions with different segregation patterns are somehow related to the size of the chromosomes, which in sturgeons is known to be very variable, with a small number of large chromosomes and a high number of micro-chromosomes. However, until complete genomes of good quality are available, it is difficult to move beyond speculation on these topics.

Author Contributions: Conceptualization, S.D.P., E.B. and L.C.; methodology, S.D.P., E.B. and L.C.; investigation, S.D.P. and S.G.B.; writing—original draft preparation, S.D.P. and L.C.; writing—review and editing, E.B., V.H.M.-M. and G.B.; project administration, G.B. and L.C.; funding acquisition, G.B. and L.C. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by Italian PRIN 2017, grant number 201794ZXTL.

Institutional Review Board Statement: The animal study protocol was approved by the Institutional Ethics Committee of the University of Padova (OPBA)—(protocol code 421922 of 14 October 2020).

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.

Acknowledgments: We would like to thank the Province of Vicenza and the “Bacino Pesca B (VI)” for the hospitality in the “Oasi le Sorgenti” Natural Reserve and for the logistical support and the Storione Ticino aquaculture farm for performing all crosses that allowed this study.

Conflicts of Interest: The authors declare no conflict of interest.

Appendix A

Locus <i>Afug132</i>		Phenotypes						
F5 ♀ x M6 ♂		A	B		C		D	
		293	309		318		330	
Observed combinations				313	318	322		
		AB (2)	293	309				
		AC (3)	293			318		
		AD (6)	293					330
		BC (2)		309		318		
		BD (3)		309				330
		CD (6)				318		330

Locus <i>Afug132</i>		Phenotypes						
F6 ♀ x M7 ♂				313	318		342	
		E	F		G	H		
Observed combinations		293	309		318	322		
		EF (7)	293	309				
		EG (7)	293			318		
		EH (4)	293				322	
		FG (4)		309		318		
		FH (4)		309			322	
		GH (3)				318	322	

Locus <i>Afug132</i>		Phenotypes						
F8 ♀ x M2 ♂		304		318	322			
		E			F	G	H	
Observed combinations		313			338	342	346	
		EF (4)		313		338		
		EG (1)		313			342	
		EH (8)		313				346
		FG (4)				338	342	
		FH (8)				338		346
		GH (4)					342	346

Locus <i>Afug41</i>		Phenotypes						
F8 ♀ x M2 ♂			212	234			293	
		E			F	G	H	
Observed combinations		203			255	259	285	
		EF (3)	203		255			
		EG (4)	203			259		
		EH (4)	203					285
		FG (5)			255	259		
		FH (4)			255		285	
		GH (7)				259	285	

Locus <i>Afug41</i>		Phenotypes						
F2 ♀ x M4 ♂		A	B	C	D			
		247	255	259	285			
Observed combinations		E		F		G	H	
		212		255			293	305
		AB (4)		247	255			
		AC (4)		247		259		
		AD (7)		247			285	
		BC (3)			255	259		
		BD (6)			255		285	
CD (3)				259	285			

Locus <i>Afug41</i>		Phenotypes						
F2 ♀ x M4 ♂		A	B	C	D			
		247	255	259	285			
Observed combinations		E		F		G	H	
		212		255			293	305
		EF (2)	212		255			
		EG (5)	212					293
		EH (7)	212					305
		FG (5)			255			293
		FH (6)			255			305
GH (2)						293	305	

Locus <i>Afug41</i>		Phenotypes						
F4 ♀ x M4 ♂		A	B	C	D			
		212	217	242		259		
Observed combinations		E		F		G	H	
		212		255			293	305
		AB (6)	212	217				
		AC (6)	212		242			
		AD (4)	212				259	
		BC (5)		217	242			
		BD (3)		217			259	
CD (5)			242		259			

Locus <i>Afug41</i>		Phenotypes						
F4 ♀ x M4 ♂		A	B	C	D			
		212	217	242		259		
Observed combinations		E		F		G	H	
		212		255			293	305
		EF (6)	212		255			
		EG (5)	212					293
		EH (3)	212					305
		FG (5)			255			293
		FH (6)			255			305
GH (4)						293	305	

Figure A1. Cont.

Locus <i>Afug132</i>		Phenotypes					
F5 ♀ x M6 ♂		A	B		C		D
		293	309		318		330
Observed combinations				313	318	322	
		AB (2)	293	309			
		AC (3)	293			318	
		AD (6)	293				330
		BC (2)		309		318	
		BD (3)		309			330
		CD (6)				318	330

Locus <i>Afug132</i>		Phenotypes							
F8 ♀ x M2 ♂		304		318	322				
			E			F	G	H	
Observed combinations						338	342	346	
		EF (4)		313			338		
		EG (1)		313				342	
		EH (8)		313				346	
		FG (4)					338	342	
		FH (8)					338	346	
		GH (4)						342	346

Locus <i>Afug132</i>		Phenotypes					
F6 ♀ x M7 ♂				313	318		342
		E	F		G	H	
Observed combinations							
		EF (7)	293	309			
		EG (7)	293			318	
		EH (4)	293				322
		FG (4)		309		318	
		FH (4)		309			322
		GH (3)				318	322

Locus <i>Afug41</i>		Phenotypes						
F8 ♀ x M2 ♂			212	234			293	
		E			F	G	H	
Observed combinations						255	259	285
		EF (3)	203			255		
		EG (4)	203				259	
		EH (4)	203					285
		FG (5)				255	259	
		FH (4)				255		285
		GH (7)					259	285

Locus <i>Afug41</i>		Phenotypes					
F2 ♀ x M4 ♂		A	B	C	D		
		247	255	259	285		
Observed combinations		E		F		G	H
		212		255			293
Observed combinations		AB (4)	247	255			
		AC (4)	247		259		
		AD (7)	247				285
		BC (3)		255	259		
		BD (6)		255		285	
		CD (3)			259	285	

Locus <i>Afug41</i>		Phenotypes						
F2 ♀ x M4 ♂		A	B	C	D			
		247	255	259	285			
Observed combinations		E		F		G	H	
		212		255			293	305
Observed combinations		EF (2)	212	255				
		EG (5)	212				293	
		EH (7)	212					305
		FG (5)		255			293	
		FH (6)		255				305
		GH (2)					293	305

Locus <i>Afug41</i>		Phenotypes					
F4 ♀ x M4 ♂		A	B	C		D	
		212	217	242		259	
Observed combinations		E		F		G	H
		212			255		293
Observed combinations		AB (6)	212	217			
		AC (6)	212		242		
		AD (4)	212				259
		BC (5)		217	242		
		BD (3)		217			259
		CD (5)			242		259

Locus <i>Afug41</i>		Phenotypes						
F4 ♀ x M4 ♂		A	B	C		D		
		212	217	242		259		
Observed combinations		E		F		G	H	
		212			255		293	305
Observed combinations		EF (6)	212			255		
		EG (5)	212				293	
		EH (3)	212					305
		FG (5)				255	293	
		FH (6)				255		305
		GH (4)					293	305

Figure A1. Cont.

Locus <i>A6</i>		Phenotypes				
F5 ♀ x M6 ♂					307	313
		E	F	G	H	
		293	297	301	307	
Observed combinations	EF (1)	293	297			
	EG (7)	293		301		
	EH (0)					
	FG (15)		297	301		
	FH (0)					
	GH (6)			301	307	

Locus <i>B11</i>		Phenotypes				
F6 ♀ x M7 ♂				148	150	
		E	F	G		H
		138	144	148		162
Observed combinations	EF (2)	138	144			
	EG (6)	138		148		
	EH (3)	138				
	FG (4)		144	148		
	FH (8)		144			162
	GH (3)				148	162

Locus <i>B11</i>		Phenotypes				
Nacc19 ♀ x Nacc17 ♂		A	B	C	D	
		132	136	138	144	
		132				150
Observed combinations	AB (4)	132	136			
	AC (7)	132		138		
	AD (3)	132			144	
	BC (5)		136	138		
	BD (4)		136		144	
	CD (3)			138	144	

Locus <i>B7</i>		Phenotypes						
F6 ♀ x M7 ♂					A	B	C	D
					166	172	174	176
		154	156	164				
Observed combinations	AB (3)			166	172			
	AC (8)			166		174		
	AD (6)			166			176	
	BC (5)				172	174		
	BD (2)				172		176	
	CD (3)					174	176	

Locus <i>B7</i>		Phenotypes					
Nacc19 ♀ x Nacc30 ♂			156	166		174	
		E			F	G	H
		154			170	174	176
Observed combinations	EF (8)	154			170		
	EG (8)	154				174	
	EH (4)	154				176	
	FG (6)				170	174	
	FH (3)				170	176	
	GH (3)					174	176

Locus <i>E4</i>		Phenotypes				
F5 ♀ x M6 ♂		A		B	C	D
		332		340	346	354
			336		346	
Observed combinations	AB (3)	332		340		
	AC (5)	332			346	
	AD (3)	332				354
	BC (6)			340		
	BD (5)			340		354
	CD (6)				346	354

Locus <i>Aox161</i>		Phenotypes				
F4 ♀ x M5 ♂		A	B	C		D
		123	127	131		139
				131	135	
Observed combinations	AB (0)					
	AC (10)	123		131		
	AD (7)	123				139
	BC (6)		127	131		
	BD (6)		127			139
	CD (0)					

Locus <i>Aox161</i>		Phenotypes				
F8 ♀ x M2 ♂		A		B	C	D
		123		131	135	139
			127	131		
Observed combinations	AB (7)	123		131		
	AC (1)	123			135	
	AD (6)	123				139
	BC (5)			131	135	
	BD (0)					
	CD (11)				135	139

Figure A1. Cont.

Locus <i>AoxD161</i>		Phenotypes					
Nacc7♀ x Nacc5♂		A	B	C			D
		123	127	131			143
Observed combinations				131	135	139	
		AB (24)	123	127			
		AC (6)	123		131		
		AD (0)					
		BC (0)					
		BD (7)		127			
CD (5)			131			143	

Locus <i>Aox234</i>		Phenotypes					
F2 ♀ x M4 ♂		A	B		C	D	
		219	223		243	255	
Observed combinations				227	243		
		AB (8)	219	223			
		AC (5)	219			243	
		AD (4)	219				255
		BC (4)		223		243	
		BD (4)		223			255
CD (2)				243	255		

Locus <i>Aox234</i>		Phenotypes							
F4 ♀ x M5 ♂		A		B	C	D			
		227		243	247	251			
Observed combinations			E	F			G	H	
			235	239			255	259	
		AB (6)	227		243				
		AC (4)	227			247			
		AD (6)	227				251		
		BC (2)			243	247			
		BD (7)			243		251		
		CD (3)				247	251		

Locus <i>Aox234</i>		Phenotypes							
F4 ♀ x M5 ♂		A		B	C	D			
		227		243	247	251			
Observed combinations			E	F			G	H	
			235	239			255	259	
		EF (8)		235	239				
		EG (4)		235				255	
		EH (4)		235					259
		FG (2)			239			255	
		FH (4)			239				259
		GH (6)						255	259

Locus <i>Aox234</i>		Phenotypes					
Nacc28♀ x Nacc23♂		A		B	C		D
		219		243	247		263
Observed combinations			E	F		G	H
			239	243		251	255
		AB (4)	219	243			
		AC (7)	219		247		
		AD (3)	219				263
		BC (4)		243	247		
BD (3)		243			263		
CD (2)			247		263		

Locus <i>Aox234</i>		Phenotypes					
Nacc28♀ x Nacc23♂		A		B	C		D
		219		243	247		263
Observed combinations			E	F		G	H
			239	243		251	255
		EF (6)		239	243		
		EG (2)		239			251
		EH (6)		239			255
		FG (5)			243	251	
		FH (0)					
		GH (4)				251	255

Locus <i>Aox234</i>		Phenotypes					
F6 ♀ x M7 ♂		A		B	C	D	
		219		227	243	247	255
Observed combinations			219		239	247	
		AB (3)		227	243		
		AC (1)		227		247	
		AD (8)		227			255
		BC (5)			243	247	
		BD (4)			243		255
		CD (8)					255

Locus <i>AoxD241</i>		Phenotypes					
F7 ♀ x M8 ♂			168		176	180	
		E		F	G	H	
Observed combinations			164		172	176	184
		EF (2)	164		172		
		EG (2)	164		176		
		EH (5)	164				184
		FG (9)			172	176	
		FH (5)			172		184
		GH (6)				176	184

Locus <i>Spl163</i>		Phenotypes					
F2 ♀ x M4 ♂			207	215	220		
		E		F		G	H
Observed combinations			166	215		224	229
		EF (11)	166	215			
		EG (0)					
		EH (5)	166				229
		FG (7)			215	224	
		FH (0)					
		GH (5)				224	229

Locus <i>Spl168</i>		Phenotypes					
F7 ♀ x M8 ♂			232		240		294
		E		F	G	H	
Observed combinations			218		236		273
		EF (0)					
		EG (5)	218			271	
		EH (6)	218				273
		FG (7)			236	271	
		FH (8)			236		273
		GH (0)					

Figure A1. Cont.

Locus <i>Spl168</i>		Phenotypes							
F4 ♀ x M5 ♂		A		B	C	D			
		209		232	257	273			
			E	F			G	H	
		218	232				279	282	
Observed combinations	AB (0)								
	AC (10)	209			257				
	AD (6)	209				273			
	BC (7)			232	257				
	BD (7)			232		273			
	CD (0)								

Locus <i>Spl168</i>		Phenotypes							
F4 ♀ x M5 ♂		A		B	C	D			
		209		232	257	273			
			E	F			G	H	
		218	232				279	282	
Observed combinations	EF (0)								
	EG (7)		218					279	
	EH (2)		218						282
	FG (13)			232				279	
	FH (8)			232					282
	GH (0)								

Locus <i>Spl168</i>		Phenotypes							
F6 ♀ x M7 ♂		A	B				C	D	
		227	236					271	286
				E	F	G	H		
			240	245	264	269			
Observed combinations	AB (0)								
	AC (6)	227						271	
	AD (4)	227							286
	BC (9)		236					271	
	BD (10)		236						286
	CD (0)								

Locus <i>Spl168</i>		Phenotypes							
F6 ♀ x M7 ♂		A	B				C	D	
		227	236					271	286
				E	F	G	H		
			240	245	264	269			
Observed combinations	EF (0)								
	EG (3)			240		264			
	EH (15)			240				269	
	FG (6)				245	264			
	FH (5)				245			269	
	GH (0)								

Locus <i>Spl168</i>		Phenotypes							
Nacc19 ♀ x Nacc17 ♂		A	B		C	D			
		218	227		269		294		
				E	F		G	H	
			227	240		286		314	
Observed combinations	AB (0)								
	AC (4)	218			269				
	AD (7)	218					294		
	BC (3)		227		269				
	BD (9)		227				294		
	CD (0)								

Locus <i>Spl168</i>		Phenotypes							
Nacc19 ♀ x Nacc17 ♂		A	B		C	D			
		218	227		269		294		
				E	F		G	H	
			227	240		286		314	
Observed combinations	EF (0)								
	EG (4)		227				286		
	EH (5)		227					314	
	FG (8)			240		286			
	FH (6)			240				314	
	GH (0)								

Locus <i>Spl168</i>		Phenotypes							
Nacc7 ♀ x Nacc5 ♂		A		B	C			D	
			232		245	264			294
		E		F			G	H	
		214		240			273	279	
Observed combinations	AB (0)								
	AC (12)		232			264			
	AD (9)		232					294	
	BC (5)				245	264			
	BD (6)				245			294	
	CD (0)								

Locus <i>Spl168</i>		Phenotypes							
Nacc7 ♀ x Nacc5 ♂		A		B	C			D	
			232		245	264			294
		E		F			G	H	
		214		240			273	279	
Observed combinations	EF (0)								
	EG (8)	214						273	
	EH (7)	214						279	
	FG (7)			240				273	
	FH (10)			240				279	
	GH (0)								

Figure A1. Cont.

Locus <i>Spl168</i>		Phenotypes						
		A		B		C	D	
Nacc8♀ x Nacc 31♂		209		236		271	279	
			E		F		G	H
			218		249		273	279
Observed combinations	AB (0)							
	AC (9)	209				271		
	AD (7)	209					279	
	BC (6)			236		271		
	BD (7)			236			279	
	CD (0)							

Locus <i>Spl168</i>		Phenotypes						
		A		B		C	D	
Nacc8♀ x Nacc 31♂		209		236		271	279	
			E		F		G	H
			218		249		273	279
Observed combinations	EF (1)			218		249		
	EG (8)			218			273	
	EH (4)			218			279	
	FG (11)					249	273	
	FH (5)					249	279	
	GH (0)							

Figure A1. Total segregation results. Single-locus microsatellite inheritance for each studied sturgeon family. Microsatellite alleles of parents are reported as sizes in bp. For families in which both parents are informative, two different schemes are shown. The 4 alleles of the informative parents are highlighted in bold and marked with capital letters A, B, C, D for females and E, F, G, H for males, also used to label allele combinations observed in the progeny. The number of F1 in which each combination has been observed is reported in brackets. Missing combinations are marked in grey.

References

- Ludwig, A. A Sturgeon View on Conservation Genetics. *Eur. J. Wildl. Res.* **2005**, *52*, 3–8. [\[CrossRef\]](#)
- Peng, Z.; Ludwig, A.; Wang, D.; Diogo, R.; Wei, Q.; He, S. Age and Biogeography of Major Clades in Sturgeons and Paddlefishes (Pisces: Acipenseriformes). *Mol. Phylogenetics Evol.* **2007**, *42*, 854–862. [\[CrossRef\]](#) [\[PubMed\]](#)
- Fontana, F.; Congiu, L.; Mudrak, V.A.; Quattro, J.M.; Smith, T.I.; Ware, K.; Doroshov, S.I. Evidence of Hexaploid Karyotype in Shortnose Sturgeon. *Genome* **2008**, *51*, 113–119. [\[CrossRef\]](#) [\[PubMed\]](#)
- Kim, D.S.; Nam, Y.K.; Noh, J.K.; Park, C.H.; Chapman, F.A. Karyotype of North American Shortnose Sturgeon *Acipenser brevirostrum* with the Highest Chromosome Number in the Acipenseriformes. *Ichthyol. Res.* **2005**, *52*, 94–97. [\[CrossRef\]](#)
- Fontana, F.; Zane, L.; Pepe, A.; Congiu, L. Polyploidy in Acipenseriformes: Cytogenetic and Molecular Approaches. In *Fish Cytogenetics*; Pisano, E., Ozouf-Costaz, C., Foresti, F., Kapoor, B.G., Eds.; Science Publisher, Inc.: Enfield, NH, USA, 2007; pp. 385–403.
- Fontana, F.; Lanfredi, M.; Chicca, M.; Congiu, L.; Tagliavini, J.; Rossi, R. Fluorescent in Situ Hybridization with rDNA Probes on Chromosomes of *Acipenser ruthenus* and *Acipenser naccarii* (Osteichthyes, Acipenseriformes). *Genome* **1999**, *42*, 1008–1012. [\[CrossRef\]](#)
- Boscari, E.; Vidotto, M.; Martini, D.; Papetti, C.; Ogden, R.; Congiu, L. Microsatellites from the Genome and the Transcriptome of the Tetraploid Adriatic Sturgeon, *Acipenser naccarii* (Bonaparte, 1836) and Cross-Species Applicability to the Diploid Beluga Sturgeon, *Huso huso* (Linnaeus, 1758). *J. Appl. Ichth.* **2015**, *31*, 977–983. [\[CrossRef\]](#)
- Ludwig, A.; Belfiore, N.M.; Pitra, C.; Svirsky, V.; Jenneckens, I. Genome Duplication Events and Functional Reduction of Ploidy Levels in Sturgeon (*Acipenser*, *Huso* and *Scaphirhynchus*). *Genetics* **2001**, *158*, 1203–1215. [\[CrossRef\]](#)
- Havelka, M.; Hulák, M.; Bailie, D.A.; Prodöhl, P.A.; Flajšhans, M. Extensive Genome Duplications in Sturgeons: New Evidence from Microsatellite Data. *J. Appl. Ichthyol.* **2013**, *29*, 704–708. [\[CrossRef\]](#)
- Stift, M.; Berenos, C.; Kuperus, P.; van Tienderen, P.H. Segregation Models for Disomic, Tetrasomic and Intermediate Inheritance in Tetraploids: A General Procedure Applied to Rorippa (Yellow Cress) Microsatellite Data. *Genetics* **2008**, *179*, 2113–2123. [\[CrossRef\]](#)
- Soares, N.R.; Mollinari, M.; Oliveira, G.K.; Pereira, G.S.; Vieira, M.L.C. Meiosis in Polyploids and Implications for Genetic Mapping: A Review. *Genes* **2021**, *12*, 1517. [\[CrossRef\]](#)
- Boscari, E.; Barbisan, F.; Congiu, L. Inheritance pattern of Microsatellite Loci in the Polyploid Adriatic Sturgeon (*Acipenser naccarii*). *Aquaculture* **2011**, *321*, 223–229. [\[CrossRef\]](#)
- Congiu, L.; Pujolar, J.M.; Forlani, A.; Cenadelli, S.; Dupanloup, I.; Barbisan, F.; Galli, A.; Fontana, F. Managing Polyploidy in Ex Situ Conservation Genetics: The Case of the Critically Endangered Adriatic Sturgeon (*Acipenser naccarii*). *PLoS ONE* **2011**, *6*, e18249. [\[CrossRef\]](#) [\[PubMed\]](#)
- Boscari, E.; Pujolar, J.M.; Dupanloup, I.; Corradin, R.; Congiu, L. Captive Breeding Programs Based on Family Groups in Polyploid Sturgeons. *PLoS ONE* **2014**, *9*, e110951. [\[CrossRef\]](#) [\[PubMed\]](#)
- May, B.; Krueger, C.C.; Kincaid, H.L. Genetic Variation at Microsatellite Loci in Sturgeon: Primer Sequence Homology in *Acipenser* and *Scaphirhynchus*. *Can. J. Fish. Aquat. Sci.* **1997**, *54*, 1542–1547. [\[CrossRef\]](#)
- Welsh, A.B.; Blumberg, M.; May, B. Identification of Microsatellite Loci in Lake Sturgeon, *Acipenser fulvescens*, and Their Variability in Green Sturgeon, *A. medirostris*. *Mol. Ecol. Notes* **2002**, *3*, 47–55. [\[CrossRef\]](#)
- Zane, L.; Patarnello, T.; Ludwig, A.; Fontana, F.; Congiu, L. Isolation and Characterization of Microsatellites in the Adriatic Sturgeon (*Acipenser naccarii*). *Mol. Ecol. Notes* **2002**, *2*, 586–588. [\[CrossRef\]](#)

18. Forlani, A.; Fontana, F.; Congiu, L. Isolation of Microsatellite Loci from the Endemic and Endangered Adriatic Sturgeon (*Acipenser naccarii*). *Conserv. Genet.* **2007**, *9*, 461–463. [[CrossRef](#)]
19. Henderson-Arzapalo, A.; King, T.L. Novel Microsatellite Markers for Atlantic Sturgeon (*Acipenser oxyrinchus*) Population De-Lineation and Broodstock Management. *Mol. Ecol. Notes* **2002**, *2*, 437–439. [[CrossRef](#)]
20. McQuown, E.C.; Sloss, B.L.; Sheehan, R.J.; Rodzen, J.; Tranah, G.J.; May, B. Microsatellite Analysis of Genetic Variation in Sturgeon: New Primer Sequences for *Scaphirhynchus* and *Acipenser*. *Trans. Am. Fish. Soc.* **2000**, *129*, 1380–1388. [[CrossRef](#)]
21. Sybenga, J. Chromosome Pairing Affinity and Quadrivalent Formation in Polyploids: Do Segmental Allopolyploids Exist? *Genome* **1996**, *39*, 1176–1184. [[CrossRef](#)]
22. Lloyd, A.; Bomblies, K. Meiosis in Autopolyploid and Allopolyploid Arabidopsis. *Curr. Opin. Plant Biol.* **2016**, *30*, 116–122. [[CrossRef](#)] [[PubMed](#)]
23. Choudhary, A.; Wright, L.; Ponce, O.; Chen, J.; Prashar, A.; Sanchez-Moran, E.; Luo, Z.; Compton, L. Varietal Variation and Chromosome Behaviour During Meiosis in *Solanum Tuberosum*. *Heredity* **2020**, *125*, 212–226. [[CrossRef](#)] [[PubMed](#)]
24. Rice, W.R. Analyzing tables of statistical tests. *Evolution* **1989**, *43*, 223–225. [[CrossRef](#)] [[PubMed](#)]
25. Scott, A.D.; Van de Velde, J.D.; Novikova, P.Y. Inference of Polyploidy Origin and Inheritance Mode from Population Genomic Data. *BioRxiv*. [[CrossRef](#)]
26. Pyatskowitz, J.D.; Krueger, C.C.; Kincaid, H.L.; May, B. Inheritance of Microsatellite Loci in the Polyploid Lake Sturgeon (*Acipenser fulvescens*). *Genome* **2001**, *44*, 185–191. [[CrossRef](#)]
27. Rodzen, J.A.; May, B. Inheritance of Microsatellite Loci in the White Sturgeon (*Acipenser transmontanus*). *Genome* **2002**, *45*, 1064–1076. [[CrossRef](#)]
28. Schreier, A.D.; Van Eenennaam, J.P.; Anders, P.; Young, S.; Crossman, J. Spontaneous Autopolyploidy in the Acipenseriformes, with Recommendations for Management. *Rev. Fish Biol. Fish.* **2021**, *31*, 159–180. [[CrossRef](#)]
29. Birstein, V.J. Sturgeon Species and Hybrids: Can Hybrids Produce Caviar. *Environ. Policy Law* **2002**, *32*, 210–214.