

Probing ionizing radiation of $L \lesssim 0.1L^*$ star-forming galaxies at $z \gtrsim 3$ with strong lensing

Eros Vanzella,^{1*} Mario Nonino,¹ Stefano Cristiani,¹ Piero Rosati,² Adi Zitrin,³ Matthias Bartelmann,³ Andrea Grazian,⁴ Tom Broadhurst,^{5,6} Massimo Meneghetti⁷ and Claudio Grillo⁸

¹INAF – Osservatorio Astronomico di Trieste, Via G. B. Tiepolo 11, 34131 Trieste, Italy

²European Southern Observatory, Karl-Schwarzschild-Str. 2, D-85748 Garching bei Munchen, Germany

³Institut für Theoretische Astrophysik, Zentrum für Astronomie der Universität Heidelberg, Albert-Überle-Str. 2, D-69120 Heidelberg, Germany

⁴INAF – Osservatorio Astronomico di Roma, Via Frascati 33, 00040 Monteporzio (RM), Italy

⁵Theoretical Physics, University of the Basque Country, Bilbao 48080, Spain

⁶Ikerbasque, Basque Foundation for Science, Alameda Urquijo, 36-5 Plaza Bizkaia, 48011 Bilbao, Spain

⁷INAF – Osservatorio Astronomico di Bologna, via Ranzani 1, 40127 Bologna, Italy

⁸Dark Cosmology Centre, Niels Bohr Institute, University of Copenhagen, Juliane Mariesvej 30, DK-2100 Copenhagen, Denmark

Accepted 2012 May 17. Received 2012 May 9; in original form 2012 April 8

ABSTRACT

We show the effectiveness of strong lensing in the characterization of Lyman continuum emission from faint $L \lesssim 0.1L^*$ star-forming galaxies at redshift $\gtrsim 3$. Past observations of $L \gtrsim L^*$ galaxies at redshift $\gtrsim 3$ have provided upper limits of the average escape fraction of ionizing radiation of $f_{\text{esc}} \sim 5$ per cent. Galaxies with relatively high f_{esc} (> 10 per cent) seem to be particularly rare at these luminosities; there is therefore the need to explore fainter limits. Before the advent of giant ground-based telescopes, one viable way to probe f_{esc} down to $0.05\text{--}0.15L^*$ was to exploit strong lensing magnification. This is investigated with Monte Carlo simulations that take into account the current observational capabilities. Adopting a lensing cross-section of 10 arcmin^2 within which the magnification is higher than 1 (achievable with about four to five galaxy clusters), with a U -band survey depth of 30 (30.5) (AB , 1σ), it is possible to constrain f_{esc} for $z \simeq 3$ star-forming galaxies down to 15 (10) per cent at 3σ for $L < 0.15L^*$ luminosities. This is particularly interesting if f_{esc} increases at fainter luminosities, as predicted from various H I reionization scenarios and radiation transfer modelling. Ongoing observational programmes on galaxy clusters are discussed and positive prospects for the future are offered, even though from space the *Hubble Space Telescope* (*HST*)/Wide Field Camera 3 (WFC3) instrument represents the only option we have to investigate details of the spatial distribution of the Lyman continuum emission arising from $z \sim 2\text{--}4$ galaxies.

Key words: gravitational lensing: strong – galaxies: distances and redshifts – galaxies: high-redshift.

1 INTRODUCTION

Recent works suggest that at $z > 3$ star-forming galaxies are the leading candidates for the production of ionizing photons (e.g. Haardt & Madau 2011; Ciardi et al. 2012; Kuhlen & Faucher-Giguère 2012). However, the mechanisms regulating the escape fraction of ionizing radiation (f_{esc}) from galaxies are still unknown. In particular, it is not clear whether f_{esc} evolves with cosmic time or whether it is luminosity-dependent at fixed redshift. The complexity in mod-

elling of galaxy evolution and the inclusion of radiative transfer prescriptions make the predictions on f_{esc} very uncertain, and opposite results are often obtained from simulations. Moreover, the predictions on f_{esc} show a large variation from galaxy to galaxy, between 0.01 and nearly 1, as a result of differences in the hydrogen distribution (e.g. Gnedin, Kravtsov & Chen 2008; Fernandez & Shull 2011; Yajima, Choi & Nagamine 2011, and references therein). From an observational point of view, the current situation is also far from clear. Various observations have provided significant upper limits on Lyman continuum (LyC) emission in the redshift range 1–4, i.e. f_{esc} smaller than 5–10 per cent (Vanzella et al. 2012, hereafter V12, and references therein). In particular, Vanzella et al.

*E-mail: vanzella@oats.inaf.it

(2010b, hereafter V10b) and Boutsia et al. (2011) reported upper limits on f_{esc} of 5 per cent for $L \gtrsim L^*$ LBGs at redshift of 3–4. Very few galaxies with possible LyC detection have been reported in the literature (Shapley et al. 2006; Iwata et al. 2009; Nestor et al. 2011). The fact that bright sources with high f_{esc} (>10 per cent) are rare could support the interpretation that f_{esc} increases at lower luminosities. Indeed, recent radiative transfer calculations coupled with cosmological simulations show that low-luminosity and dwarf galaxies have high f_{esc} (>20–40 per cent) and are the major contributors to the ionizing background at redshift $3 < z < 6$ (e.g. Wise & Cen 2009; Razoumov & Sommer-Larsen 2010; Fernandez & Shull 2011; Yajima et al. 2011; but see Gnedin et al. 2008, for different results). The investigation of faint (and/or low-mass) star-forming galaxies at moderate redshift ($z \sim 3$ –4) is therefore crucial to probe the ionization regimes and conditions that would be in place during the reionization epoch ($z > 7$). It is worth noting that the reionization at $z = 7$ –10 requires an average f_{esc} much higher than what is observed for relatively bright and lower redshift galaxies ($z < 4$) (Bouwens et al. 2010; Robertson et al. 2010; Fernandez & Shull 2011).

While the direct measurements of escaping LyC photons are prohibitive during the epoch of reionization, they are still accessible at $z \lesssim 4$. The investigation of sub- L^* galaxies requires extremely deep U -band surveys ($U > 30$). However, even if they were available, spectroscopic redshift measurements for $L \ll L^*$ galaxies (e.g. r band > 27) would be very challenging with current facilities. In this respect, before the advent of the ELT and JWST telescopes, strong lensing magnification offered a viable way to explore faint limits in luminosity (and mass).

2 THE f_{esc} AND STRONG LENSING

The *relative* fraction of escaping LyC photons (at $\lambda < 912 \text{ \AA}$) relative to the fraction of escaping non-ionizing ultraviolet (1500 \AA) photons is defined as (Steidel, Pettini & Adelberger 2001)

$$f_{\text{esc,rel}} \equiv \frac{(L_{1500}/L_{\text{LyC}})_{\text{int}}}{\left(\frac{\mu F_{1500}}{\mu F_{\text{LyC}}}\right)_{\text{obs}}} \exp\left(\tau_{\text{LyC}}^{\text{IGM}}\right), \quad (1)$$

where $(F_{1500}/F_{\text{LyC}})_{\text{obs}}$, $(L_{1500}/L_{\text{LyC}})_{\text{int}}$ and $\tau_{\text{LyC}}^{\text{IGM}}$ represent the observed $1500 \text{ \AA}/\text{LyC}$ flux density ratio, the intrinsic $1500 \text{ \AA}/\text{LyC}$ luminosity density ratio and the line-of-sight opacity of the intergalactic medium (IGM) for LyC photons, respectively. It is worth noting that the two quantities $(F_{1500})_{\text{obs}}$ and $(F_{\text{LyC}})_{\text{obs}}$ are measured in the same spatial (i.e. physical) region, where the ionizing and non-ionizing radiations arise (V12). If the dust attenuation A_{1500} is known, $f_{\text{esc,rel}}$ can be converted to the absolute f_{esc} as $f_{\text{esc}} = 10^{-0.4A_{1500}} f_{\text{esc,rel}}$ (e.g. Siana et al. 2007). The term μ is the magnification factor provided by a lens (e.g. a cluster of galaxies) and applies to both flux densities observed at two wavelengths (we leave the parameter explicitly in the expression). We assume for simplicity that it does not vary substantially within the isophote of the sources. Therefore, given the achromatic nature of lensing, f_{esc} is independent from μ . The two quantities f_{esc} and $f_{\text{esc,rel}}$ are equal if $A_{1500} = 0$. Conservatively, in the following we perform calculations assuming $A_{1500} = 0$; the limits probed on f_{esc} are deeper if $A_{1500} > 0$, e.g. they are halved if $A_{1500} = 0.6$ (V10b). If not specified, we assume an intrinsic luminosity density ratio $(L_{1500}/L_{\text{LyC}})_{\text{int}} = 7$ (Siana et al. 2007; V10b). In Section 3.2, the IGM transmission of Inoue et al. (2011) is convolved with the adopted U -band filter. We perform calculations for three U -band filters: *F336W*, *u*-band Strömberg (*U-strom*) and *U-special* available at the Large Binocular

Telescope (*LBC-U*) (in general the results do not change if other filters with similar widths and wavelength coverages are considered). It is clear from equation (1) that high magnitude contrasts between LyC and the 1500 \AA rest frame provide strong constraints on f_{esc} . We can rearrange equation (1) to give an estimate of the expected magnified flux at wavelengths smaller than the Lyman limit as a function of f_{esc} and magnification factor μ (i.e. $F_{\text{LyC}}^{\text{Lens}} = \mu F_{\text{LyC}}$):

$$F_{\text{LyC}}^{\text{Lens}} = \left(\frac{L\lambda_{\text{rest}}}{L_{1500}}\right)_{\text{int}} \frac{f_{\text{esc}} \times (F_{1500} \times \mu)_{\text{obs}}}{e^{\tau_{\text{LyC}}^{\text{IGM}}}} \times 10^{0.4 \times A_{1500}}. \quad (2)$$

As discussed in Inoue & Iwata (2008) and V10b, the stochastic nature of the IGM absorption introduces large uncertainties in the estimate of f_{esc} for a single line of sight. Stacking many galaxies among different lines of sight provides strong limits and reduces the variance due to IGM attenuation. Indeed, a deep U -band survey can provide firm constraints for $L \gtrsim L^*$ galaxies down to few per cent of f_{esc} by stacking tens of objects. This has been performed by V10b with deep Very Large Telescope/VIMOS U -band imaging ($\approx 30AB$ at 1σ) of $z \gtrsim 3.4$ LBGs (see also Boutsia et al. 2011).

For $L \ll L^*$ galaxies the situation becomes challenging. One reason is that the knowledge of the redshift is necessary to fix the observed wavelength position of the Lyman limit. Even though a deep U -band survey (e.g. mag- $U \simeq 30$ at 1σ) can still probe f_{esc} down to 20 per cent by stacking about thirty $0.1L^*$ galaxies, the redshift confirmation is not practicable with current facilities, or it is feasible only for a subclass of sources like the Ly α emitters. The lensing magnification greatly facilitates the redshift measurement also for faint galaxies. Additionally, it allows the investigation of f_{esc} down to stronger limits by U -band stacking and/or individual LyC detections. In the following we perform Monte Carlo (MC) simulations to derive predictions for f_{esc} as a function of the U -band depth, lensing magnification and lensing cross-section. In what follows, we adopt r and R to indicate the r -band magnitude ($\sim 1500 \text{ \AA}$ rest frame) for non-magnified and magnified sources, respectively.

3 MAGNIFIED GALAXIES: MC SIMULATIONS

We now want to estimate the feasibility of probing f_{esc} down to <20 per cent for $L \lesssim 0.1L^*$ LBGs. To this aim, from equation (2) it is possible to calculate as a function of f_{esc} and U -band depth the number of sources detectable in the LyC above a certain threshold and the signal-to-noise ratio (S/N) in their LyC stack. We therefore perform MC simulations similar to those described in V10b by assuming reasonable distributions for the quantities involved in equation (2).

3.1 The expected number of magnified LBGs

The probability that a source galaxy is magnified by more than μ can be expressed in terms of the probability density $P(\mu)$: $P(>\mu) = \int_{\mu}^{\infty} P(\mu) d\mu$, where $P(\mu) = -dP(>\mu)/d\mu$. An interesting property of the lensing probability in the source plane is that $P(>\mu) \propto \mu^{-2}$ and therefore $P(\mu) \propto \mu^{-3}$ for $\mu \gg 1$, as can be shown in particular cases and argued to be true in general (see fig. 9 of Lima, Jain & Devlin 2010; Schneider, Ehlers & Falco 1992). A maximum magnification (μ_{max}) is imposed by the size of the source galaxies that we assume to have an intrinsic half-light radius not smaller than 0.2 kpc (assuming circular shape), i.e. $\mu_{\text{max}} = 400$.

Each galaxy cluster produces a region within which the magnification in the source plane is larger than μ_{min} . This area is an effective cross-section for lensing statistics. For our purposes, it is

enough to perform a simple calculation for the cross-section σ_{lens} :

$$\sigma_{\text{lens}} \simeq \frac{1}{\mu_{\text{min}}^2} \sum_{i=1}^{N_{\text{clust}}} \pi \theta_{E,i}^2 \quad (\text{arcmin}^2), \quad (3)$$

where $\theta_{E,i}$ is the Einstein radius of a given cluster. Lenses with the largest θ_E are the best ‘cosmic telescopes’; in particular, we consider in our simulations the case with $\sigma_{\text{lens}} = 10 \text{ arcmin}^2$ and $\mu_{\text{min}} = 1$, corresponding approximately to four to five massive galaxy clusters (see Section 4), and the results can be linearly rescaled to other areas.

Adopting the $z \sim 3$ luminosity function (LF) parameters of Reddy & Steidel (2009) ($\alpha = -1.73$, $M_{AB}^* = -20.97$ and $\phi_* = 1.71 \times 10^{-3} \text{ Mpc}^{-3}$), we can calculate the expected number of star-forming galaxies within the chosen area and redshift interval dz . A further important constraint is that the magnification must produce an apparent magnitude R brighter than the spectroscopic limit for redshift measurements ($R < m_{\text{spec}}$) necessary to fix exactly the wavelength position of the Lyman limit. We adopt $m_{\text{spec}} = 26$ that corresponds to a relatively deep spectroscopic survey (e.g. Vanzella et al. 2009). Together with the limit on μ (< 400) defined above, the chosen m_{spec} implies a limit on the integration of the LF, i.e. $r = 32.5$ (i.e. sources with $r > 32.5$ would require $\mu > 400$ to have $R < 26$). It turns out that within an area $\sigma_{\text{lens}} = 10 \text{ arcmin}^2$ and redshift interval 3.0–3.5, ~ 2300 LBGs are expected in the magnitude range $26.5 < r < 32.5$, 30 of which have $R < m_{\text{spec}} = 26$ (we refer to this sample as $N_{\text{spec}} = 30$). Clearly, the intrinsically faint galaxies are the rarer cases accessible from ground-based spectroscopy, the galaxies being sources that need large magnification μ . We anticipate here that the constraints we discuss below on f_{esc} are mainly dictated by sources belonging to the magnitude bin $26.5 < r < 28$ (i.e. $L = 0.05\text{--}0.16L^*$, on average 24 of the 30 galaxies).

3.2 MC simulations of $L \lesssim 0.1L^*$ LBGs

We run MC simulations on the area $\sigma_{\text{lens}} = 10 \text{ arcmin}^2$ by varying for each galaxy the parameters involved in equation (2) and derive the ionized flux $F_{\text{LyC}}^{\text{Lens}}$ as a function of f_{esc} . The flux is then compared with the depth of the U -band survey under study. The procedure is described in detail in V10b. Briefly, a redshift is randomly extracted from the interval dz (uniformly) and associated with a galaxy with a given r magnitude extracted randomly from the magnitude range 26.5–32.5 (accordingly with the magnitude distribution provided by the LF). The intergalactic transmission $T_{\text{LyC}}^{\text{IGM}} = e^{-\tau_{\lambda}^{\text{IGM}}}$ has been derived from thousands of random realizations at the extracted redshift value and convolved with the adopted U -band filter. The lower limit z_{min} of the redshift range dz is dictated by the adopted filter shape that probes LyC at $z > 3$, 3.2 and 3.3 for the three filters $F336W$, $U\text{-strom}$ and $LBC\text{-}U$, respectively. The ideal upper limit would be $z_{\text{min}} + 0.1$ such that the closest region bluewards of the Lyman limit is probed. However, we relax to $z_{\text{min}} + 0.5$ by allowing more sources to be included (in order to reach $N_{\text{spec}} \simeq 30$ in the three cases). The IGM prescription adopted here includes all the intervening absorption systems and modulates properly the $F_{\text{LyC}}^{\text{Lens}}$ signal for increasing redshift. This is also supported by the recent direct detection of LyC at $\lambda_{\text{rest}} < 830 \text{ \AA}$ (V10b).

The magnification factor (μ) is associated with each galaxy by extracting it randomly from the distribution $P(\mu) \sim \mu^{-3}$, as explained earlier in the text (we assume that galaxies are distributed uniformly over the sky). The f_{esc} is explored in two different regimes of variability: (1) it is fixed to a constant value for all N_{spec} sources spanning the range 0.01–1.00, with steps of 0.01 (i.e. the average $\langle f_{\text{esc}} \rangle = f_{\text{esc}}$ of each galaxy) and (2) it is fixed to values 0 or 1,




Figure 1. MC simulations for magnified LBGs at $3 < z < 3.5$ with intrinsic $L < 0.14L^*$ ($r > 26.5$) and adopting a lensing cross-section of 10 arcmin^2 (see text for details). Top, middle and bottom panels show the results of our MC simulations for U -band depths of 29, 30 and 31 (1σ), respectively. Left-hand panels: the expected S/N in the LyC of the stacked fluxes (median and central 68 per cent interval, solid and dashed lines, respectively). Filled circles are the S/N (median and central 68 per cent interval) in the ‘on/off case’ with 20 per cent of the galaxies with $f_{\text{esc}} = 1$ and 80 per cent with $f_{\text{esc}} = 0$. Horizontal dotted lines mark S/N = 3. Right-hand panels: number of direct detections in the LyC at S/N > 1 (black solid line) and S/N > 2 (red dashed line). The dotted black lines report the number of galaxies in the faint magnitude bins from which they migrate to brighter apparent magnitudes ($R < 26$). The black solid line is the sum of the dotted lines. From top to bottom, the dotted lines refer to the magnitude bins $26.5 < r < 27.0$, $27.0 < r < 27.5$, etc. The red stars mark the expected number of direct detections at S/N > 2 in the ‘on/off case’.

assigned randomly such that 80 per cent of the N_{spec} sample have $f_{\text{esc}} = 0$ and 1 for the remaining sources, i.e. $\langle f_{\text{esc}} \rangle = 0.2$ with large variance (the ‘on/off case’).

In summary, the sample of galaxies derived from the LF in the considered volume (area of 10 arcmin^2 and $dz = 0.5$) and magnitude interval $26.5 < r < 32.5$ has been extracted 10 000 times by varying f_{esc} , $e^{-\tau_{\lambda}^{\text{IGM}}}$, $(L_{1500}/L_{\text{LyC}})_{\text{int}}$, $(F_{1500})_{\text{obs}}$ and μ (no dust has been considered). Fig. 1 shows the results for the $F336W$ case. The resulting median S/N of the stacked LyC fluxes of the galaxies with $R < 26$ is shown as a function of f_{esc} and U -band depth (left-hand panels). The signal-to-noise ratio $S/N_{\text{LyC}}^{\text{stack}}$ is calculated as $[\sum_{i=1}^{N_{\text{spec}}} F_{\text{LyC}}^{i,\text{Lens}} / N_{\text{spec}}] / [1\sigma_U / \sqrt{N_{\text{spec}}}]$, where $1\sigma_U$ is the depth of the U -band survey and $N_{\text{spec}} \simeq 30$, i.e. the number of surviving LBGs with $R < 26$. The number of direct LyC detections above the 2σ limit ($F_{\text{LyC}}^{\text{Lens}} > 2\sigma_U$) is also reported in the same Fig. 1 (right-hand panels). While the median $S/N_{\text{LyC}}^{\text{stack}}$ of the two f_{esc} distributions, constant and ‘on/off’, are compatible if $\langle f_{\text{esc}} \rangle = 0.2$ (as expected from the $S/N_{\text{LyC}}^{\text{stack}}$ calculation), the variance is larger in the ‘on/off’ case. Also, the number of direct detections are different between the two, because by definition we fix a maximum fraction of detectable sources. It turns out that with a U -band depth $\gtrsim 30(1\sigma)$, the S/N of the stacking is $> 1(3)$ for $f_{\text{esc}} > 0.05(0.18)$. The average number of individual detections ($> 2\sigma$) is > 3 for $f_{\text{esc}} > 0.10(0.30)$ if the U -band depth is 31(30). These detections are originated from the magnification of sources with $r > 26.5$ that migrate from their




Figure 2. S/N in the LyC of the stacking (top panels) and the fraction (per cent) of LyC detections with S/N higher than 1 and 2 (bottom panels) are shown as a function of the U -band depth. Three different U -band filters have been used. In the three cases, $\sigma_{\text{lens}} = 10 \text{ arcmin}^2$ and $N_{\text{spec}} \simeq 30$. In all panels the three curves have been calculated for $f_{\text{esc}} = 0.2, 0.15$ and 0.10 from top to bottom, respectively. Horizontal dotted lines mark the one and 3σ levels. In the bottom panels solid and dashed lines refer to $S/N > 1$ and $S/N > 2$, respectively. The insets in the top panels show the adopted filter, with the dotted vertical line marking the position of the Lyman limit at the z_{min} .

original magnitude bins to the brighter lensed regime ($R < 26$). The contribution of each bin to the N_{spec} sample is shown as dotted lines in Fig. 1. As a consistency check, and as expected, their number per magnitude bin flattens if the faint end slope of the LF is $\alpha = -3$ (i.e. the growth of the number counts and the dimming of the lensing cross-section compensate each other). The flattening of the number of direct detections as f_{esc} increases (in the fixed f_{esc} case, Fig. 1) is due to IGM attenuation, i.e. the interception of Lyman limit systems (LLSs) or damped Ly α systems (DLAs) along the line of sight suppresses the ionizing flux even for high f_{esc} values.

Fig. 2 shows the same quantities as a function of the U -band depth by fixing f_{esc} to 10, 15 and 20 per cent for all sources. It has been performed for three filters: *F336W*, *U-strom* and *LBC-U*. Clearly, the deeper U -band surveys give the best constraints to the faint galaxies; in particular, those with $\text{mag-}U > 30$ are the more effective and increase rapidly as the $\text{mag-}U = 31$ limit is approached. The intermediate band filter like the *U-strom* is an optimal solution. It is worth noting that observing a massive cluster with $\theta_E \simeq 1 \text{ arcmin}$ (such as MACS 0717.5+3745, see below), we expect ~ 10 galaxies with $R < 26$ and $r > 26.5$. If all of them have $f_{\text{esc}} = 15$ per cent, then with a U -band (*U-strom*-like) depth of 30.5 we expect to detect three of them at $S/N > 1$ and $S/N_{\text{LyC}} \simeq 3.0$ for the stacking. Therefore, we expect to observe signal already with a single deep pointing (of $\sim 60 \text{ h}$) with a 8–10 m class telescope.

4 THE CURRENT OBSERVATIONS

As mentioned above, the interception of LLSs or DLAs makes IGM attenuation quite stochastic and severe for ionizing radiation at $z \gtrsim 3$; therefore, the observations of magnified galaxies along

different lines of sight are necessary to attenuate its effect. Assuming these absorbers are associated with individual structures with physical size of $\lesssim 50 \text{ kpc}$ (e.g. Yajima, Choi & Nagamine 2012), different lines of sight separated by more than $\sim 10 \text{ arcsec}$ will have a relatively small transverse correlation among high column density absorbers, even along a single cluster.

Currently, there are in the literature several massive clusters of galaxies characterized by relatively large lensing cross-sections, e.g. $\theta_E \sim 0.7\text{--}0.9 \text{ arcmin}$, like MACS 0717.5+3745 (Zitrin et al. 2009), 1E 0657–56 (‘bullet cluster’; Bradac et al. 2009) and A1689 (Broadhurst et al. 2005). Zitrin et al. (2012) recently measured the Einstein radius of 10 000 galaxy clusters from the Sloan Digital Sky Survey. In particular, they identified $\simeq 40$ candidates with $\theta_E > 0.7 \text{ arcmin}$. The largest Einstein radius was $\theta_E \simeq 1.1 \text{ arcmin}$ for the most massive cluster (for a source at $z_s = 2$). The MASSive Cluster Survey (MACS) also provides a statistically complete sample of very X-ray luminous distant clusters of galaxies ($0.3 < z < 0.7$; Ebeling et al. 2010). Therefore, given the existing observations, the area of $\sigma_{\text{lens}} = 10 \text{ arcmin}^2$ considered in this Letter can be easily achieved with five clusters.

It is worth mentioning the ongoing Cluster Lensing And Supernova survey with Hubble (CLASH), a multitreasury programme that, in addition to other available ground-based observations, is observing (524 *HST* orbits) 25 X-ray selected massive galaxy clusters with new *HST* panchromatic imaging capabilities (16 filters): the ACS and both the ultraviolet and infrared channels of the WFC3. While 20 clusters were chosen to be X-ray selected, relatively relaxed clusters with no lensing-selection bias, the five additional clusters were chosen to be high-magnification clusters with $\theta_E > 0.6 \text{ arcmin}$ (Postman et al. 2012). Even though the survey was not designed to provide deep limits on LyC for faint galaxies, the homogeneous panchromatic photometry on a statistically significant sample of massive clusters, together with magnification maps and high S/N spectroscopy on a number of highly magnified galaxies at $z \gtrsim 3$, will yield a very valuable test bed for probing LyC emission at the faint end of the galaxy LF.

5 CONCLUSIONS

We have explored the possibilities offered by gravitational lensing in constraining the ionizing emission from faint galaxies at $z > 3$, which has a great impact on studies of the reionization even at significantly higher redshifts ($z > 6$). We have shown that with current ground-based facilities and the magnification provided by known clusters of galaxies, it is possible to constrain the f_{esc} quantity down to 10–15 per cent for faint luminosities ($L \lesssim 0.1L^*$). In particular, strong lensing provides the opportunity to make a significant step forward in this field in the pre-ELT era. (1) The magnification allows us to measure redshifts for very faint galaxies, i.e. to fix the wavelength position of the Lyman limit and put strong constraints on the LyC with U -band imaging for galaxies down to $L \sim 0.05\text{--}0.1L^*$. (2) The magnification allows us to perform accurate spatial analysis of the LyC emission (if detected), down to hundreds of parsecs. If, on the one hand, the magnified area increases the probability of intercepting a foreground lower- z source that might mimic LyC emission (Vanzella et al. 2010a), on the other hand the possible presence of multiple images of the same background source will help solve the problem, since a real LyC detection will be present in all the counterparts (if μ are similar). The transverse separation of the light paths decreases rapidly as the source redshift is approached. Therefore, the IGM attenuation is practically the same for all the multiple images. (3) It has been suggested that the f_{esc}

increases with decreasing luminosity (e.g. Yajima et al. 2011) and that the ionizing background is mainly produced by a large number of sub- L^* galaxies. A nice feature of the present approach is to put a limit to the process of attributing the ionizing contribution to lower and lower luminosities with the increasing observational depth of null results. This in fact would imply a steepening of the faint end of the LF that in principle can be excluded by the increasing number of highly magnified sources expected to be detected through lensing.

We have shown that deep U -band observations (as in V10b) of five massive galaxy clusters provide a sample of intrinsically faint ($L \lesssim 0.1L^*$) and magnified $z \sim 3$ galaxies (about 30) useful to investigate f_{esc} down to 10–20 per cent. An increasing number of well-studied clusters for this study, with multiband HST photometry, spectroscopy and lens modelling, are becoming available as part of the CLASH project.

ACKNOWLEDGMENTS

We thank the anonymous referee for the fruitful comments. We thank G. Cupani and F. Fontanot for useful discussions. AZ is supported by contract research ‘Internationale Spitzenforschung II-1’ of the Baden-Württemberg Stiftung. We acknowledge financial contribution from the agreement ASI-INAF I/009/10/0 and from the PRIN MIUR 2009 ‘The Intergalactic Medium as a probe of the growth of cosmic structures’ and the Dark Cosmology Centre which is funded by the Danish National Research Foundation.

REFERENCES

- Boutsia K. et al., 2011, *ApJ*, 736, 41
 Bouwens R. J. et al., 2010, *ApJ*, 708, 69
 Bradac M. et al., 2009, *ApJ*, 706, 1201
 Broadhurst T. et al., 2005, *ApJ*, 621, 53
 Ciardi B., Bolton J. S., Maselli A., Graziani L., 2012, *MNRAS*, 423, 558
 Ebeling H., Edge A. C., Mantz A., Barrett E., Henry J. Patrick M. J., van Speybroeck L., 2010, *MNRAS*, 407, 83
 Fernandez E. R., Shull J. M., 2011, *ApJ*, 731, 20
 Gnedin N. Y., Kravtsov A. V., Chen H.-W., 2008, *ApJ*, 672, 765
 Haardt F., Madau P., 2011, *ApJ*, 746, 125
 Inoue A. K., Iwata I., 2008, *MNRAS*, 387, 1681
 Inoue A. K. et al., 2011, *MNRAS*, 411, 2336
 Iwata I. et al., 2009, *ApJ*, 692, 1287
 Kuhlen M., Faucher-Giguère C. A., 2012, *MNRAS*, 423, 862
 Lima M., Jain B., Devlin M., 2010, *MNRAS*, 406, 2352
 Nestor D. B., Shapley A. E., Steidel C. C., Siana B., 2011, *ApJ*, 736, 18
 Postman M. et al., 2012, *ApJ*, 199, 25
 Razoumov A. O., Sommer-Larsen J., 2010, *ApJ*, 710, 1239
 Reddy N. A., Steidel C. C., 2009, *ApJ*, 692, 778
 Robertson B. E., Ellis R. S., Dunlop J. S., McLure R. J., Stark D. P., 2010, *Nat*, 468, 49
 Schneider P., Ehlers J., Falco E. E., 1992, *Gravitational Lenses*. Springer-Verlag, Berlin
 Shapley A. E., Steidel C. C., Pettini M., Adelberger K. L., Erb D. K., 2006, *ApJ*, 651, 688
 Siana B. et al., 2007, *ApJ*, 668, 62
 Steidel C. C., Pettini M., Adelberger K. L., 2001, *ApJ*, 546, 665
 Vanzella E. et al., 2009, *ApJ*, 695, 1163
 Vanzella E., Siana B., Cristiani S., Nonino M., 2010a, *MNRAS*, 404, 1672
 Vanzella E. et al., 2010b, *ApJ*, 725, 1011 (V10b)
 Vanzella E. et al., 2012, *ApJ*, 751, 70 (V12)
 Wise J. H., Cen R., 2009, *ApJ*, 693, 984
 Yajima H., Choi J.-H., Nagamine K., 2011, *MNRAS*, 412, 411
 Yajima H., Choi J.-H., Nagamine K., 2012, *MNRAS*, submitted (arXiv:1112.5691)
 Zitrin A., Broadhurst T., Rephaeli Y., Sadeh S., 2009, *ApJ*, 707, 102
 Zitrin A., Broadhurst T., Bartelmann M., Rephaeli Y., Oguri M., Benítez N., Hao J., Umetsu K., 2012, *MNRAS*, 423, 2308

This paper has been typeset from a \TeX/L\AA\TeX file prepared by the author.