

Journal Pre-proof

Predictors of Bubble Formation and Type Obtained with Pneumatic Dissection During Deep Anterior Lamellar Keratoplasty in Keratoconus

Vincenzo Scordia, Giuseppe Giannaccare, Andrea Lucisano, Mauro Soda, Giovanna Carnovale Scalzo, James Myerscough, Marco Pellegrini, Francesco Verdoliva, Gabriele Piccoli, Cristina Bovone, Massimo Busin

PII: S0002-9394(19)30613-0

DOI: <https://doi.org/10.1016/j.ajo.2019.12.012>

Reference: AJOPHT 11173

To appear in: *American Journal of Ophthalmology*

Received Date: 29 July 2019

Revised Date: 26 November 2019

Accepted Date: 18 December 2019

Please cite this article as: Scordia V, Giannaccare G, Lucisano A, Soda M, Scalzo GC, Myerscough J, Pellegrini M, Verdoliva F, Piccoli G, Bovone C, Busin M, Predictors of Bubble Formation and Type Obtained with Pneumatic Dissection During Deep Anterior Lamellar Keratoplasty in Keratoconus, *American Journal of Ophthalmology* (2020), doi: <https://doi.org/10.1016/j.ajo.2019.12.012>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2019 Published by Elsevier Inc.



PURPOSE: To identify predictors of bubble formation and type during big-bubble deep anterior lamellar keratoplasty (BB-DALK) performed in keratoconus at different stages of severity.

DESIGN: Retrospective Cohort Study.

METHODS: Setting: University Magna Græcia (Catanzaro, Italy); Study Population: Consecutive keratoconus patients undergoing BB-DALK from September 2014 to February 2019; Observation Procedure: Keratometric astigmatism, mean keratometry value (K-mean), highest keratometry value (K-max), thinnest point, anterior segment Optical Coherence Tomography (AS-OCT)-based stage of ectasia. Main Outcome Measures: Rate of bubble formation and type; number and fate of micro/macro-perforation; conversion to mushroom keratoplasty (MK); comparison of parameters in patients with bubble formation vs failure and in type 1 vs type 2 bubble; areas under the curves (AUC) of preoperative parameters for distinguishing between bubble types.

RESULTS: Pneumatic dissection succeeded in 113/155 eyes (72.9%), with 100 type 1 bubbles (88.4%), 11 type 2 (9.8%) and 2 mixed-type (1.8%). Micro-perforations were managed conservatively in type-1 bubbles; macro-perforations occurring in both types of bubbles required conversion to MK. Preoperative K-mean and K-max values were significantly higher in eyes in which bubble formation succeeded (respectively, $P=0.006$ and $P<0.013$). Type 1 bubbles formed in eyes with significantly lower age, K-mean, AS-OCT stages and higher pachymetric values (always $P<0.029$). Age had the highest diagnostic power for discrimination between bubble types, followed by AS-OCT stage, pachymetry, K-mean and astigmatism (respectively, $AUC=0.861$; 0.779 ; 0.748 ; 0.700 ; 0.670).

CONCLUSIONS: Older age and advanced stages of keratoconus were predictors of type 2 bubble formation during BB-DALK that was associated with an increased risk of complications.

Predictors of Bubble Formation and Type Obtained with Pneumatic Dissection During Deep Anterior Lamellar Keratoplasty in Keratoconus

American Journal of Ophthalmology - Original Article

Vincenzo Scordia,^{1,2} Giuseppe Giannaccare,¹ Andrea Lucisano,¹ Mauro Soda,¹ Giovanna Carnovale Scalzo,¹ James Myerscough,³ Marco Pellegrini,⁴ Francesco Verdoliva,¹ Gabriele Piccoli,¹ Cristina Bovone,^{2,3,5} Massimo Busin.^{2,3,5}

¹ *Department of Ophthalmology, University Magna Græcia, Catanzaro, Italy*

² *Istituto Internazionale per la Ricerca in Oftalmologia (IRFO), Forlì, Italy*

³ *Ospedale Privato Villa Igea, Forlì, Italy*

⁴ *Ophthalmology Unit, S. Orsola-Malpighi Hospital, University of Bologna, Bologna, Italy*

⁵ *Department of Morphology, Surgery and Experimental Medicine, University of Ferrara, Ferrara, Italy*

Short Title: Predictors of Bubble Characteristics During DALK

Corresponding Author:

Vincenzo Scordia, MD

Professor of Ophthalmology

University of Magna Græcia, Catanzaro, Italy

Address: Via dei Crociati 40, 88100, Catanzaro, Italy

Mail Address: vscordia@libero.it

Phone/Fax: 0039 09613647133

INTRODUCTION

Characterized by progressive corneal thinning, protrusion, and irregular astigmatism, keratoconus (KC) is the most common corneal ectasia, and, in its early stages, can be managed adequately by spectacle or contact lens correction.¹⁻⁶ However, in cases of reduced visual acuity secondary to corneal scarring, contact lens intolerance or uncorrectable high degree irregular astigmatism, a surgical intervention is mandated.⁷ Deep anterior lamellar keratoplasty (DALK) is increasingly reported as a technique utilized to treat corneal stromal pathologies with healthy endothelium and, being essentially an extraocular intervention, DALK not only eliminates endothelial rejection but also reduces the risks associated with “open-sky” surgery.⁸⁻¹² The most widely used technique to perform DALK is the big-bubble (BB-DALK), as described by Anwar, during which, injection of air in the deep stroma results in cleavage of the deep stroma from either the pre-Descemet’s layer (PDL) or Descemet’s membrane (DM).¹³ This natural plane of separation avoids the interface irregularity that can be associated with manual dissection.¹⁴

Recent studies have demonstrated that the cleavage plane obtained during pneumatic dissection differs, with three possible types of air bubble created.¹⁵ A type 1 bubble, (the most common) originates from the center of the cornea, extending radially, and represents a plane of separation between the deep stroma and the PDL, whereas a type 2 bubble originates from the periphery and extends centripetally, forming between PDL and DM. Finally, both bubbles may coexist to form a mixed-bubble. The corresponding outcomes of BB-DALK depend not only on the formation or failure of the bubble, but also on the type of bubble itself. For example, since the floor of a type 2 bubble consists only of DM, it carries a high risk of perforation which can mandate conversion to full thickness penetrating keratoplasty. Furthermore, even in the absence of perforation, the occurrence of a type-2 bubble has recently been associated with an increased risk of post-operative double anterior chamber formation requiring further intervention.¹⁶ Formation of a type-2 bubble increases, therefore, the intraoperative challenge of DALK, and the identification of predictive factors correlating with its occurrence may help improving the outcomes of DALK.

The aim of this study was to investigate whether statistical analysis of preoperative demographic and clinical parameters may succeed in predicting the success of pneumatic dissection (bubble formation) in addition to the type of bubble obtained with it in keratoconic corneas undergoing DALK.

METHODS

Study Design

In this institutional retrospective cohort study, the records of all consecutive DALK procedures performed in patients affected by KC at a tertiary referral Center (Department of Ophthalmology, University of Magna Graecia, Catanzaro, Italy) from September 2014 to February 2019 were reviewed. The study followed the tenets of the 2013 Declaration of Helsinki and was approved by the local Ethics Committee (Comitato Etico Università Magna Graecia of Catanzaro, Italy). Informed consent was obtained from all patients undergoing surgery. Patients who did not gain useful visual acuity with spectacles and were contact lens were considered candidates for surgery. Eyes with previous hydrops

or evident lesions of DM were excluded. Preoperatively, all patients underwent a complete ophthalmologic evaluation including slit-lamp examination, corrected distance visual acuity (CDVA) expressed both in logarithmic units of the minimum angle of resolution (logMAR) and in Snellen fraction, refraction, tonometry, fundoscopy, endothelial specular microscopy (EM-3000; Tomey, Erlangen, Germany), and anterior segment optical coherence tomography (AS-OCT) (Casia; Tomey, Tokyo, Japan).

Preoperative Data Collection

The following preoperative data were collected for the statistical analysis: age; sex; keratometric astigmatism; K-mean; K-max; thinnest pachymetric measurement; severity of disease graded according to a previously described AS-OCT classification:¹⁷ Stage 1, thinning of apparently normal epithelial and stromal layers at the conus; Stage 2, hyperreflective anomalies occurring at the Bowman's layer level with epithelial thickening at the conus, without or with stromal opacities (2 a and 2 b, respectively); Stage 3, posterior displacement of the hyperreflective structures occurring at the Bowman's layer level with increased epithelial thickening and stromal thinning, without or with stromal opacities (3 a and 3 b, respectively); Stage 4, pan-stromal scar; Stage 5, hydrops, acute onset or healing phase (5 a and 5 b, respectively).

Surgical Procedure

Surgery was performed in all eyes by a single surgeon (V.S.); anesthesia and akinesia were obtained by means of peribulbar injection of 10 mL of a 0.75% ropivacaine solution in all cases but 5, who received general anesthesia. As described previously in detail,¹⁸ pharmacologic mydriasis was induced in all eyes before starting surgery; after, the geometric center of the cornea was marked and a disposable Hessburg-Barron suction trephine (Katena Products Inc, Denville, NJ) was used to create circular incision in the recipient cornea between 8.75 and 9 mm in diameter. The peripheral pachymetric map obtained before surgery by means of the CASIA corneal topographer serve as a reference for the trephination of approximately 80% of the total corneal thickness in depth. Pneumatic dissection was attempted by first advancing a dedicated probe and then a cannula up to 2-3 mm centripetally from the bottom of the deep trephination. Regardless of the success of pneumatic dissection, in all cases the recipient cornea was de-bulked by performing an anterior keratectomy, which removed about 80% of the anterior stroma; even when the residual bed was very thin, this thickness was sufficient to prevent the viscoelastic device (OVD) from breaking through to the surface. When pneumatic dissection succeeded, adhesive viscoelastic substance (IAL-F, Fidia, Padova, Italy) was laid centrally onto the bubble roof and a 15° blade was used to enter the bubble. The inferior arm of a blunt Vannas scissor was inserted into the collapsed bubble through the slit obtained with the 15° blade, which was enlarged to allow completion of the removal of the bubble roof by means of corneal scissors. In case of bubble smaller than 8.75 to 9.00 mm in diameter, opening of the bubble roof was extended only up to the white peripheral ring formed by the air injection; then, a blunt dissector (Galan stromal dissector spatula, Janach, Como, Italy) was used to gently enlarge the dissection up to the trephination edge thus allowing a safer removal of the remaining tissue crown. When a mixed bubble formed, only the type-1 part was opened, leaving the type-2 part untouched and simply waiting for air to reabsorb post-operatively. When pneumatic dissection failed, the procedure was continued, attempting injecting ophthalmic viscoelastic device (VI-VACY sodium hyaluronate 1.55%, I-SPACE Laboratoires, La Ravoire, France) through

a cannula inserted into the same stromal path used for pneumatic dissection, according to the surgical steps previously described.¹⁹ If also OVD-assisted dissection failed, layer-by-layer manual dissection was performed aiming at reaching a level in the deep stroma free from all micro-bubbles. The donor cornea was punched from the endothelial side with a Barron donor punch (Katena Products, Inc.) of the same diameter of the trephine used on the recipient cornea. Then, after staining with 0.06% trypan blue dye (VisionBlue; D.O.R.C., Zuiland, the Netherlands), DM and endothelium was gently stripped off using a dry Weck-Cel sponge. Four interrupted 10-0 nylon sutures secured the graft into the recipient bed and, surgery was completed with 2 running, 10-0 nylon sutures of 16 bites each. Starting the following day, betamethasone 0.2% and chloramphenicol 0.5% eye drops were administered every 2 hours, then tapered off to a single daily steroidal administration one month after partial suture removal and finally discontinued at month 6 after surgery.

Intraoperative Parameters

The following intraoperative data were recorded for each surgery: bubble formation (obtained by means of air or OVD injection); bubble type; micro- or macro-perforations; conversion to MK.

Statistical Analysis

The SPSS statistical software (SPSS Inc., Chicago, Illinois, USA) was used for data analysis. Values were expressed as mean \pm standard deviation (SD). The Mann–Whitney U test was used to compare continuous and ordinal variables between cases with bubble formation vs bubble failure and cases with type-1 vs cases with type-2 bubble formation. Receiver operating characteristic (ROC) curves with calculations of the area under the curve (AUC) were used to describe the accuracy of each parameter for discriminating type-1 and type-2 bubbles; for the statistical analysis, the cases of mixed type bubble were included in the group of type-2 bubble. Sensitivity and specificity of each parameter were determined for the cutoff value whose corresponding point on the ROC curve was nearest to the coordinate (0,100). A multivariate binomial logistic regression was performed to predict the likelihood of a type-1 bubble; a *P* value < 0.05 was considered statistically significant.

RESULTS

A total of 155 consecutive keratoconic eyes of 143 patients undergoing DALK were included in the study with the corresponding demographical and clinical characteristics reported in Table 1. Although they were not considered an exclusion criteria, none of the eyes included in this study had undergone corneal cross-linking or ICR implantation. Pneumatic dissection succeeded in 113 of 155 eyes (72.9%). Type-1 bubble was obtained in 100/113 eyes (88.4%), type-2 in 11/113 eyes (9.8%) and mixed type in 2/113 eyes (1.8%). OVD-assisted dissection succeeded in 25 of the remaining 42 cases (59.0%) in which pneumatic dissection had failed. In the type-1 group, a microperforation occurred in 5 eyes (5%) which was addressed conservatively by air fill of the anterior chamber, with a single case of macroperforation (1%) requiring conversion to a mushroom shaped penetrating keratoplasty. In the type-2 group, we recorded only macroperforations (2 eyes, 18.1%), both of them requiring conversion to MK. Manual

dissection was completed in the remaining 17 eyes. Among these, microperforations occurred in 2 eyes (11.7%), none requiring conversion to MK.

The preoperative K-mean and K-max values were significantly higher in cases with bubble formation (respectively $P = 0.006$ and $P = 0.013$), while no significant differences were found preoperatively for age, gender, keratometric astigmatism, AS-OCT classification and thinnest point between cases with bubble formation and failure (always $P > 0.05$, Table 1).

Cases in which a type-1 bubble was formed showed a significantly lower age ($P < 0.001$), lower K-mean ($P = 0.029$), higher thinnest point ($P = 0.007$) and lower stage of AS-OCT classification ($P < 0.002$) compared to those in which a type-2 bubble was achieved (Table 1). Figure 1 shows a representative case of a type-2 bubble occurring in a patient with severe KC (AS-OCT classification stage 4).

Eyes in which pneumatic dissection succeeded showed significantly higher values of K-mean (63.5 ± 8.5 vs 57.7 ± 5.2 diopters), K-max (66.8 ± 9.2 vs 61.1 ± 5.9 diopters), and lower values of thinnest point (351.3 ± 69.9 vs 387.8 ± 59.5 microns) compared to those in which OVD-dissection was achieved. Conversely, no significant differences in age, gender, keratometric astigmatism and AS-OCT stage were found between the two groups ($P > 0.05$).

The ROC curves of age, keratometric astigmatism, K-mean, thinnest point and AS-OCT stage for discriminating type-1 and type-2 bubbles are shown in Figure 2, part A. Table 2 reports the AUC values of ROC curves with 95% confidence intervals, cutoff values, sensitivity and specificity. Age had the highest diagnostic power for the discrimination between type-1 and type-2 bubble (AUC = 0.861), followed by AS-OCT stage (AUC = 0.779), thinnest point (AUC = 0.748), K-mean (AUC = 0.700), and keratometric astigmatism (AUC = 0.670).

A binomial logistic regression was performed to ascertain the effects of age, keratometric astigmatism and AS-OCT stage on the likelihood of bubble type formation. To avoid multicollinearity, K-mean and thinnest point were not included in the regression model, since they were strongly correlated with the AS-OCT stage (respectively, $R = 0.675$ and $R = -0.690$, both $P < 0.001$). The logistic regression model was statistically significant, $\chi^2(3) = 28.098$, $P < 0.001$. The model explained 46.2% (Nagelkerke R^2) of the variance in the type of bubble and correctly classified 91.4% of cases. The ROC curve of the regression model is shown in Figure 2, part B. The AUC was 0.916 (95% CI: 0.825 to 1), with a sensitivity of 90.9% and a specificity of 89.5%.

Increasing age and more advanced AS-OCT stages were associated with an increased likelihood of type-2 bubble (Table 3).

DISCUSSION

Our study was designed to investigate whether preoperative demographic and clinical parameters were able to predict the formation and the type of bubble during BB-DALK for keratoconus, in order to identify potential eyes at high risk of type-2 bubble formation. Other than conventional keratometry data, clinical parameters included a novel AS-OCT-based grading system of keratoconus.¹⁷ This technology has been reported to be instrumental in the analysis of the progressive structural changes of the deep stroma during the evolution of the disease. These modifications often involve the precise layer into which the air is injected, and therefore, may have a critical role in the success rate

or type of bubble obtained by means of air injection.

In our series, preoperative maximum and mean keratometric readings values differed significantly between the two groups in which the bubble formed (regardless of the type) or failed, with higher values correlated with successful DALK completion. These results confirm the outcomes of previous studies reporting a higher success rate in eyes with advanced keratoconus, where the more ectatic tissue allows an easier access to a deeper plane, usually revealed by a sudden decrease of tissue resistance to the advancement of the cannula.²⁰⁻²³ Conversely, eyes with milder stages of keratoconus, where the failure of pneumatic dissection was higher, might benefit from OVD injection through the same track that was used unsuccessfully for air injection bubble.^{19,24}

Several parameters including age, K-mean, thinnest point and AS-OCT staging differed significantly between cases in which type-1 bubble occurred compared to type-2 bubbles. In particular, milder stages of KC (higher pachymetric values, lower K-mean, and early AS-OCT stages from 1 to 3 a) in addition to younger age were associated with type-1 bubble formation. This correlation between age and bubble type occurrence, here reported for the first time, may suggest that the PDL may undergo changes with aging, similarly to DM which is modified by the formation of advanced glycation end products, with alterations in its biophysical properties.²³ However, to date, all in vitro studies examining the PDL were carried out in adult eyes, and, therefore, histological data pertaining to younger eyes are not yet available. The reason for which previous studies have not identified an association between age and bubble type is likely to be related to the heterogeneous etiological inclusion criteria applied in such studies, with a heavier skew towards older patients when including non-KC related disease.²¹ In our analysis, the cut off value of 55 years had the highest values of sensitivity and specificity for the discrimination of bubble type occurrence. However, further histopathology studies are required to detect the possible variations occurring at a cellular level in the PDL in relation to both patient's age and severity of KC.

In our series we found the involvement of posterior corneal surface (AS-OCT stage 3 b onwards) to be a significant risk factor for type-2 bubble formation. We hypothesize that the deep scarring may fuse the PDL to the posterior stroma or degrade the PDL itself, preventing type-1 bubble formation due to the direct air dispersion at the level of DM. Recently, posterior stromal scars, resulting from different corneal diseases, have been associated with the failure of pneumatic dissection,^{21,22} and this poor prognostic value was found to be stronger in patients with KC compared to other corneal pathologies.²¹ Accordingly, in order to homogenize our outcomes and eliminate multiple variables, we enrolled in this study only keratoconic patients.

We are aware that intraoperative variables such as the depth of placement of the cannula during pneumodissection may influence the success rate of the surgery. Therefore, we decided to analyze data obtained from a single-surgeon experience that used a reproducible approach to reach the pre-Descemet's stroma based on a deep trephination.¹⁸ However, a further prospective study is needed to validate in a second subset the predictive factors detected in this work.

When performing pneumatic dissection, type-1 bubble formation is desirable because the floor of the bubble includes PDL, providing strength to the thin layer and reducing the risk of perforation, which are usually small and manageable conservatively. Instead,

the type-2 floor is fragile, consisting only of DM, and, despite recent techniques proposed to optimally manage this bubble,^{25,26} the completion of DALK surgery remains a challenge, with conversion to PK often needed. It is therefore unsurprising that in this series the conversion rate to MK is also much higher in the group of patients with a type-2 bubble than those with a type-1 bubble (18.1% vs 1%). In order to reduce the rate of perforation, different surgical techniques could be suggested as a safer alternative to conventional big-big-bubble DALK, such as diamond knife assisted or pachimetry guided manual techniques.^{27,28} Another choice could be also not to postpone too much the surgery thus performing surgery in less advanced stages of keratoconus in which bubble type 1 is more likely to occur.

In conclusion, this study highlighted that during DALK for keratoconus, the combination of age and stage of disease predicts the type of bubble to be obtained. Specifically, advanced stages of the disease detected by AS-OCT and increasing age were found to be strong predictors of the occurrence of type-2 bubble. Therefore, a detailed preoperative assessment of KC may provide useful information for the surgical planning of DALK in terms of both timing and technique employed. Early surgical intervention should be considered in such cases before the development of deep stromal scarring associated in BB-DALK with subsequent type-2 bubble formation and increased risk of conversion to PK. On the other hand, in case of severe KC cases, the use of manual techniques can be taken into account.

Funding/Support: None.

Financial Disclosure: *The authors have no proprietary or commercial interest in any materials discussed in the article. M.B.: Reimbursement of travel expenses and royalties - Moria, Antony, France.*

Acknowledgments: None.

Journal Pre-proof

REFERENCES

1. Nottingham J. Practical Observations on Conical Cornea: and on the Short Sight, and Other Defects of Vision Connected with it. London: John Churchill; 1854:1e19.
2. Amsler M. Le kératocone fruste au Javal. *Ophthalmologica* 1938;96:77-83.
3. Amsler M. Keratocone Classique et keratocone fruste; arguments unitaires. *Ophthalmologica* 1946;111:96-101.
4. Krachmer JH, Feder RS, Belin MW. Keratoconus and related non inflammatory corneal thinning disorders. *Surv Ophthalmol* 1984;28(4):293-322.
5. Kennedy RH, Bourne WM, Dyer JA. A 48-year clinical and epidemiologic study of keratoconus. *Am J Ophthalmol* 1986;101(3):267-273.
6. Rabinowitz YS. Keratoconus. *Surv Ophthalmol* 1998;42(4):297-319.
7. Kubaloglu A, Sari ES, Unal M. Long-term results of deep anterior lamellar keratoplasty for the treatment of keratoconus. *Am J Ophthalmol* 2011;151(5):760-767.
8. Reinhart WJ, Musch DC, Jacobs DS et al. Deep anterior lamellar keratoplasty as an alternative to penetrating keratoplasty a report by the American Academy of Ophthalmology. *Ophthalmology* 2011;118(1):209-218.
9. Coster DJ, Lowe MT, Keane MC, Williams KA, Australian Corneal Graft Registry Contributors. A comparison of lamellar and penetrating keratoplasty outcomes: a registry study. *Ophthalmology* 2014;121(5):979-987.
10. Borderie VM, Sandali O, Bullet J, Gaujoux T, Touzeau O, Laroche L. Long-term results of deep anterior lamellar versus penetrating keratoplasty. *Ophthalmology* 2012;119(2):249-255.
11. Keane M, Coster D, Ziaei M, Williams K. Deep anterior lamellar keratoplasty versus penetrating keratoplasty for treating keratoconus. *Cochrane Database Syst Rev* 2014;(7):CD009700.
12. Giannaccare G, Weiss JS, Sapigni L et al. Immunologic stroma rejection after deep anterior lamellar keratoplasty with grafts of a larger size (9 mm) for various stromal diseases. *Cornea* 2018;37(8):967-972.
13. Anwar M, Teichmann KD. Big-bubble technique to bare Descemet's membrane in anterior lamellar keratoplasty. *J Cataract Refract Surg* 2002;28(3):398-403.
14. Knutsson KA, Rama P, Paganoni G. Modified big-bubble technique compared to manual dissection deep anterior lamellar keratoplasty in the treatment of keratoconus. *Acta Ophthalmol* 2015;93(5):431-438.
15. Dua HS, Mastropasqua L, Faraj L et al. Big bubble deep anterior lamellar keratoplasty: the collagen layer in the wall of the big bubble is unique. *Acta Ophthalmol* 2015;93(5):427-430.
16. Myerscough J, Friehmann A, Bovone C, Mimouni M, Busin M. Management of type 2 bubble formed during big-bubble deep anterior lamellar keratoplasty. *Cornea* 2019;38(6):e20.

17. Sandali O, El Sanharawi M, Temstet C et al. Fourier-domain optical coherence tomography imaging in keratoconus: a corneal structural classification. *Ophthalmology* 2013;120(12):2403-2412.
18. Scorcia V, Lucisano A, Pietropaolo R, Savoca Corona V, Scorcia G, Busin M. Red reflex-guided big-bubble deep anterior lamellar keratoplasty: a simple technique to judge dissection depth. *Cornea* 2015;34(9):1035-1038.
19. Scorcia V, De Luca V, Lucisano A et al. Results of viscobubble deep anterior lamellar keratoplasty after failure of pneumatic dissection. *Br J Ophthalmol* 2018;102(9):1288-1292.
20. Bhatt UK, Fares U, Rahman I, Said DG, Maharajan SV, Dua HS. Outcomes of deep anterior lamellar keratoplasty following successful and failed 'big bubble'. *Br J Ophthalmol* 2012;96(4):564-569.
21. Borderie VM, Touhami S, Georgeon C, Sandali O. Predictive factors for successful type 1 big bubble during deep anterior lamellar keratoplasty. *J Ophthalmol* 2018; 13;2018:4685406.
22. Ozmen MC, Yesilirmak N, Aydin B, Ceylanoglu KS, Atalay HT, Akata F. Prediction of Descemet membrane perforation during deep anterior lamellar keratoplasty in patients with keratoconus with stromal scar. *Eye Contact Lens* 2018;44 Suppl 2:S176-S179.
23. Schlötzer-Schrehardt U, Bachmann BO, Tourtas T et al. Ultrastructure of the posterior corneal stroma. *Ophthalmology* 2015;122(4):693-639.
24. Muftuoglu O, Toro P, Hogan RN et al. Sarnicola air-visco bubble technique in deep anterior lamellar keratoplasty. *Cornea* 2013;32(4):527-532.
25. Goweida MB, Ragab AM, Liu C. Management of Type 2 Bubble Formed During Big Bubble Deep Anterior Lamellar Keratoplasty. *Cornea* 2019;38(2):189-193.
26. Goweida MB, Ragab AM, Liu C. Reply. *Cornea* 2019; 38(6):e20-e21.
27. Rama P, Knutsson KA, Razzoli G, Matuska S, Viganò M, Paganoni G. Deep anterior lamellar keratoplasty using an original manual technique. *Br J Ophthalmol* 2013;97(1):23-7;
28. Vajpayee RB, Maharana PM, Sharma N, Agarwal T, Jhanji V. Diamond knife-assisted deep anterior lamellar keratoplasty to manage keratoconus. *J Cataract Refract Surg* 2014;40(2):276-82.

FIGURE CAPTIONS:

Figure 1: Representative images from a patient with severe keratoconus - Anterior segment Optical Coherence Tomography scan of a patient with severe keratoconus (Stage 4) showing a pan-stomal scar (Part A). Intraoperative image of bubble 2 formation during pneumatic dissection in the same patient (Part B).

Figure 2: Receiver operating characteristic (ROC) curves for discriminating type 1 and type 2 bubbles - ROC curves of age, keratometric astigmatism, K-mean, thinnest point and AS-OCT stage (Part A). ROC curve of the multivariate logistic regression model including age, corneal cylinder and AS-OCT stage as independent variables (Part B).

Table 1: Parameters of patients undergoing DALK^a - Comparison of demographical and clinical parameters of patients according to the success or failure of bubble formation and the type of bubble formed during DALK.^a

Parameter	Overall (n = 155)	Bubble Success			Bubble Type		
		Formation (n = 113)	Failure (n = 42)	<i>P</i>	Type 1 (n = 100)	Type 2 (n = 11)	<i>P</i>
Age (years)	41.4 ± 15.7	41.7 ± 15.9	40.3 ± 15.2	0.704	39.5 ± 15.1	58.7 ± 8.7	<0.001
Gender (Male/Female)	87/68	69/49	18/19	0.293	60/45	7/4	0.678
Keratometric astigmatism (D) ^b	4.9 ± 3.0	5.0 ± 3.0	4.5 ± 2.8	0.258	5.1 ± 3.0	3.6 ± 3.0	0.065
K mean ^c (D) ^b	62.9 ± 8.6	64.0 ± 8.4	59.4 ± 8.3	0.006	63.4 ± 8.2	69.4 ± 8.8	0.029
K max ^d (D) ^b	66.2 ± 9.4	67.2 ± 9.2	62.7 ± 9.3	0.013	66.8 ± 9.1	71.9 ± 9.9	0.091
Thinnest point (µm)	350.1 ± 75.2	346.1 ± 72.5	362.8 ± 83.1	0.181	353.8 ± 67.5	279.2 ± 88.2	0.007
AS-OCT ^e classification				0.441			0.002
Stage 1	22 (14.2%)	14 (11.9%)	8 (21.6%)		14 (13.3%)	0 (0.0%)	
Stage 2 a	53 (34.2%)	42 (35.6%)	11 (29.7%)		40 (38.1%)	1 (9.1%)	
Stage 2 b	18 (11.6%)	12 (10.2%)	6 (16.2%)		10 (9.5%)	2 (18.2%)	
Stage 3 a	16 (10.3%)	15 (12.7%)	1 (2.7%)		15 (14.3%)	0 (0.0%)	

Stage 3 b	27 (17.4%)	21 (17.8%)	6 (16.2%)	16 (15.2%)	4 (36.4%)
Stage 4	19 (12.3%)	14 (11.9%)	5 (13.5%)	10 (9.5%)	4 (36.4%)

^aDALK = deep anterior lamellar keratoplasty; ^bD = diopters; ^cK mean = mean keratometric value; ^dK max = highest keratometry value; ^eAS-OCT = anterior segment-Optical Coherence Tomography.

Journal Pre-proof

Table 2: Areas under the curves (AUC) of parameters for the discrimination between bubble types - AUC with 95% confidence intervals (CIs), sensitivity, and specificity of parameters for the discrimination between type 1 and type 2 bubbles. Sensitivity and specificity were determined at the cutoff value whose corresponding point on the receiver operating characteristic (ROC) curve was nearest to the coordinate (0,100).


Parameter	AUC	95% CI	Cutoff value	Sensitivity (%)	Specificity (%)
Age	0.861	0.773 to 0.949	55.5 years	83.9	90.9
Keratometric astigmatism	0.670	0.476 to 0.864	3.6 D	67.6	63.6
K mean ^a	0.700	0.567 to 0.834	63.9 D	60.0	72.7
Thinnest point	0.748	0.586 to 0.910	307.5 μ m	75.2	63.6
AS-OCT ^b classification	0.779	0.649 to 0.910	Stage 3a	75.2	72.7

^aK mean = mean keratometric value; ^bAS-OCT = anterior segment-Optical Coherence Tomography.

Table 3: Multivariate logistic regression predicting the likelihood of a type 1 bubble - Multivariate logistic regression predicting the likelihood of the occurrence of a type 1 bubble based on age, keratometric astigmatism and AS-OCT^a classification.

Parameter	B	Odds ratio	95% CI	<i>P</i>
Age	-0.078	0.925	0.874 to 0.979	0.007
Keratometric astigmatism	0.309	1.362	0.973 to 1.906	0.072
AS-OCT ^a classification	-0.753	0.471	0.280 to 0.790	0.004

^aAS-OCT = *anterior segment-Optical Coherence Tomography*.



Journal Pre-proof

