


This document is confidential and is proprietary to the American Chemical Society and its authors. Do not copy or disclose without written permission. If you have received this item in error, notify the sender and delete all copies.

Indomethacin Co-Crystals and their Parent Mixtures: does the Intestinal Barrier Recognize them Differently?

Journal:	<i>Molecular Pharmaceutics</i>
Manuscript ID:	Draft
Manuscript Type:	Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Ferretti, Valeria; University of Ferrara, Department of Chemical and Pharmaceutical Sciences Dalpiaz, Alessandro; University of Ferrara, Department of Chemical and Pharmaceutical Sciences Bertolasi, Valerio; University of Ferrara, Department of Chemical and Pharmaceutical Sciences Ferraro, Luca; University of Ferrara, Department of Life Sciences and Biotechnology Beggiato, Sarah; University of Ferrara, Department of Life Sciences and Biotechnology Spisni, Enzo; University of Bologna, Department of Biological, Geological and Environmental Sciences Pavan, Barbara; University of Ferrara, Department of Chemical and Pharmaceutical Sciences

SCHOLARONE™
Manuscripts



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Indomethacin Co-Crystals and their Parent Mixtures: does the Intestinal Barrier Recognize them Differently?

Valeria Ferretti[†], Alessandro Dalpiaz^{†,*}, Valerio Bertolasi[†], Luca Ferraro[‡], Sarah Beggiato[‡], Enzo Spisni[§] and Barbara Pavan[‡]

[†]Department of Chemical and Pharmaceutical Sciences, University of Ferrara, Ferrara, Italy

[‡]Department of Life Sciences and Biotechnology, University of Ferrara, Ferrara, Italy

[§]Department of Biological, Geological and Environmental Sciences, University of Bologna, Bologna, Italy

1
2
3 **ABSTRACT:** Co-crystals are crystalline complexes of two or more molecules bound together in
4
5
6 crystal lattices through non-covalent interactions. The solubility and dissolution properties of co-
7
8
9 crystals can allow to increase the bioavailability of poorly water soluble active pharmaceutical
10
11 ingredients (APIs). It is currently believed that the co-crystallization strategy should not induce changes
12
13 on the pharmacological profile of the APIs, even if it is not yet clear whether a co-crystal would be
14
15 defined as a physical mixture or as a new chemical entity. In order to clarify these aspects, we chose
16
17 indomethacin as guest poorly aqueous soluble molecule and compared its properties with those of its
18
19 co-crystals obtained with 2-hydroxy-4-methyl-pyridine (co-crystal *1*), 2-methoxy-5-nitroaniline (co-
20
21 crystal *2*) and saccharine (co-crystal *3*). In particular, we performed a systematic comparison among
22
23 indomethacin, its co-crystals and their parent physical mixtures by evaluating via HPLC analysis the
24
25 API dissolution profile, its ability to permeate across intestinal cell monolayers (NCM460) and its oral
26
27 bioavailability in rat. The indomethacin dissolution profile was not altered by the presence of co-
28
29 crystallizing agents as physical mixtures, whereas significant changes were observed by the dissolution
30
31 of the co-crystals. Furthermore, there was a qualitative concordance between the API dissolution
32
33 patterns and the relative oral bioavailabilities in rats. Co-crystal *1* induced a drastic decrease of the
34
35 transepithelial electrical resistance (TEER) value of NCM460 cell monolayers, whereas its parent
36
37 mixture did not evidence any effect. The saccharin-indomethacin mixture induced a drastic decrease of
38
39 the TEER value of monolayers, whereas its parent co-crystal *3* did not induce any effects on their
40
41 integrity, being anyway able to increase the permeation of indomethacin. Taken together, these results
42
43 demonstrate for the first time different effects induced by co-crystals and their parent physical mixtures
44
45 on a biologic system, findings that could raise serious concerns about the use of co-crystal strategy to
46
47 improve API bioavailability without performing appropriate investigations.

48
49
50
51
52
53
54
55 **KEYWORDS:** *co-crystals, indomethacin, saccharin, drug permeation, NCM 460 cells,*
56
57
58 *bioavailability*

INTRODUCTION

The therapeutic efficacy of a pharmaceutical formulation depends on its bioavailability, i.e. the absorption extent and rate of the active pharmaceutical ingredient (API) into the bloodstream following its administration. The bioavailability of a solid pharmaceutical formulation may depend in turn on the dissolution profile of its components, in particular of the API. In the case of highly lipophilic API, and therefore of a compound poorly soluble in water but capable of effectively permeate through biological membranes (Biopharmaceutical Classification System - BCS - class II), the dissolution process is the limiting factor of its absorption; in this case the bioavailability is highly dependent on both dissolution rate and maximum amount dissolved of the APIs themselves.¹

The solid state form of an API is determinant in influencing its solubility and dissolution rate. In general, the amorphous phases are easier to solubilize than crystalline solids and, among them, the metastable polymorphs can offer solubility or dissolution advantages with respect to stable ones. On the other hand, the risks of phase transformation of amorphous solids or metastable polymorphs do not allow their employment in manufacturing pharmaceutical solid dosage forms.^{2,3} The crystal engineering of pharmaceutical solids may be therefore very useful to optimize the API stability and bioavailability, and the co-crystals seem to be promising in this context.⁴ A co-crystal can be considered as a crystalline complex of two or more molecules bound together in the crystal lattice through non-covalent interactions, often including hydrogen bonding. Pharmaceutical co-crystals are obtained by an API and a co-crystal former.⁵ It is known that the solubility and dissolution properties of co-crystals can be similar to those of amorphous compounds, *i.e.* higher than the parent crystalline pure phases. As a consequence, pharmaceutical co-crystals give the opportunities to increase bioavailability of APIs showing, at the same time, the stability of their stable crystalline forms.^{6,7}

1
2
3 Currently, several APIs are known to improve their solubility profile and bioavailability when
4 co-crystallized.⁷⁻⁹ These APIs include the anticonvulsant carbamazepine,¹⁰ nonsteroidal anti-
5 inflammatory drugs such as indomethacin and meloxicam,^{11,12} the flavonoid quercetin,¹³ or other
6 molecules employed as model drugs.¹⁴⁻¹⁷

7
8
9
10
11
12 It is currently believed that the co-crystallization strategy should not induce changes on the
13 pharmacological profile of the APIs. Indeed, co-crystal design requires changes on crystal structures
14 that essentially alter hydrogen bonding motifs rather than covalent bonds of the API, thus retaining its
15 safety and therapeutic properties.^{11,18} On the other hand, the regulatory status regarding the use of co-
16 crystals in pharmaceutical products appears still unsettled. In particular, it is not yet clear whether a co-
17 crystal would be defined as a physical mixture (enabling its classification within current compendial
18 guidelines) or as a new chemical entity requiring full safety and toxicology testing.^{7,8} In this context,
19 FDA has taken the position that a co-crystal may be treated as a drug product intermediate.⁸

20
21
22
23
24
25
26
27
28
29
30
31
32 In this study we evaluated the properties of (i) indomethacin, chosen as guest molecule poorly
33 soluble in aqueous environment,¹¹ (ii) its two new co-crystals with 2-hydroxy-4-methyl-pyridine (co-
34 crystal **1**), 2-methoxy-5-nitroaniline (co-crystal **2**) and (iii) a previously described indomethacin-
35 saccharine co-crystal^{11,19} (co-crystal **3**). The schematic representation of indomethacin and the co-
36 formers are reported in Figure 1. In particular, the dissolution, the permeation across NCM460 cell
37 monolayers employed as an *in vitro* model of human intestinal epithelial barrier, and the bioavailability
38 after oral administration to rats of indomethacin, its co-crystals and their parent mixtures (**1**, **2** and **3**,
39 respectively) have been investigated. Overall, the results indicate, for the first time, that strongly
40 different effects on the integrity of intestinal cell monolayers can derive by the dissolution of co-
41 crystals or their parent mixtures.

MATERIALS AND METHODS

Materials and Reagents. γ -indomethacin, 2-hydroxy-4-methyl-pyridine, 2-methoxy-5-nitro-aniline, saccharine, methanol, acetonitrile, ethyl acetate, isoamyl acetate and water were high performance liquid chromatography (HPLC) grade from Sigma Aldrich (Milan, Italy). All other reagents and solvents were of analytical grade (Sigma-Aldrich). NCM-460 cells were kindly provided by Dr. Antonio Strillacci, University of Bologna, Italy. The male Sprague-Dawley rats were provided by Charles-River (Milan, Italy).

Synthesis of Adducts. Two new co-crystals containing the indomethacin API have been synthesized and characterized by X-ray crystallography: **Co-crystal 1**: γ -indomethacin and 2-hydroxy-4-methyl-pyridine 1:1; **Co-crystal 2**: γ -indomethacin and 2-methoxy-5-nitro-aniline 1:1. Two other co-crystals have been synthesized and characterized but not used in the present work because of their poor reproducibility: **Co-crystal a**: γ -indomethacin and 4-nitropyridine-N-oxide monohydrate 1:1:1; **Co-crystal b**: γ -indomethacin and pyridine-N-oxide 1:1. Details of the X-ray crystallographic analysis for all the four crystals are reported in Supplementary Table 1. An equimolar quantity of indomethacin and co-crystal partner was dissolved in the minimum quantity of isoamyl acetate and left for slow evaporation at room temperature. Crystals were observed after a few days. **Co-crystal 3**, containing saccharine as the coformer, has been obtained by solvent slow evaporation of an equimolar saccharine/ γ -indomethacin solution prepared according to ref. 19. The phase and composition of the co-crystals **1**, **2** and **3** have been checked by X-ray powder crystallography, comparing the experimental spectra with those calculated from the single-crystal crystallography structures (supplementary Figures. S1-S3)

Experimental – X-Ray. The crystallographic data for the four co-crystals **1**, **2**, **a**, **b** were

1
2
3 collected on a Nonius Kappa CCD diffractometer at room temperature using graphite-monochromated
4
5 MoK α radiation ($\lambda = 0.71073 \text{ \AA}$). Data sets were integrated with the Denzo-SMN package²⁰ and
6
7 corrected for Lorentz-polarization effects. The structures were solved by direct methods with the
8
9 SIR97 suite of programs²¹ and refinement was performed on F^2 by full-matrix least-squares methods
10
11 with all non-hydrogen atoms anisotropic. The N/O-H atoms were found in the difference Fourier map
12
13 and refined isotropically; all other hydrogen atoms were included on calculated positions, riding on
14
15 their carrier atoms. All calculations were performed using SHELXL-97²² implemented in the WINGX
16
17 system of programs.²³ The ORTEPIII²⁴ diagrams of co-crystals **1** and **2** are shown in Figure 2. Powder
18
19 diffraction spectra for co-crystals **1**, **2** and **3** were recorded, at room temperature, on a Bruker D-8
20
21 Advance diffractometer with graphite monochromatized Cu K α radiation ($\lambda = 1.5406 \text{ \AA}$). The data
22
23 were recorded at 2θ steps of 0.02° with 1 s/step. Crystallographic data for the structural analysis of the
24
25 four new compounds have been deposited at the Cambridge Crystallographic Data Center, 12 Union
26
27 Road, Cambridge, CB2 1EZ, UK, and are available free of charge from the Director on request quoting
28
29 the deposition number CCDC 1005832-1005835 for **1**, **2**, **a** and **b**, respectively.
30
31
32
33
34
35

36 **HPLC Analysis.** The quantification of the indomethacin was performed by HPLC. The
37
38 chromatographic apparatus consisted of a modular system (model LC-10 AD VD pump and model
39
40 SPD- 10A VP variable wavelength UV–vis detector; Shimadzu, Kyoto, Japan) and an injection valve
41
42 with 20 μL sample loop (model 7725; Rheodyne, IDEX, Torrance, CA, USA). Separation was
43
44 performed at room temperature on a reverse phase column Hypersil BDS C-18, 5U, equipped with a
45
46 guard column packed with the same Hypersil material (Alltech Italia Srl BV, Milan, Italy). Data
47
48 acquisition and processing were accomplished with a personal computer using CLASS-VP Software,
49
50 version 7.2.1 (Shimadzu Italia, Milan, Italy). The detector was set at 319 nm. The mobile phase
51
52 consisted of a mixture of methanol and 0.2 phosphoric acid (75:25 v/v). The flow rate was 1 mL/min.
53
54
55
56
57 The compound 9-phenyl- carbazole was employed as internal standard in extraction procedures of
58
59
60

1
2
3 indomethacin from rat blood (see below). The retention times for indomethacin and 9-phenyl-carbazole
4
5 were 4.0 and 13.5 minutes, respectively.
6
7

8 The chromatographic precision for each compound was evaluated by repeated analysis ($n = 6$)
9
10 of the same samples (100 μM). For indomethacin and 9-phenyl-carbazole (employed as internal
11
12 standard in the extraction procedures) dissolved in aqueous phase the values were obtained for 100 μM
13
14 (0.036 mg/ml) solutions and were represented by the relative standard deviation (RSD) values ranging
15
16 between 0.63% and 0.74%, respectively.
17
18

19 The calibration curves of indomethacin dissolved in phosphate buffer 200 mM and in PBS 10
20
21 mM were linear over the ranges of 50 μM (0.018 mg/ml) - 1500 μM (0.54 mg/ml) and 2 μM (0.00072
22
23 mg/ml) – 500 μM (0.18 mg/ml), respectively ($n = 8$, $r > 0.997$, $P < 0,0001$). The limit of quantification
24
25 for indomethacin was 625 nM (224 ng/ml, 2.24 ng injected) with a signal-to-noise ratio of 10, whereas
26
27 the limit of detection was 188 nM (67 ng/ml, 0.67 ng injected) with a signal-to-noise ratio of 3.
28
29

30 A preliminary analysis performed with 100 μM solutions showed that hydroxy-4-methyl-
31
32 piridine, 2-methoxy-5-nitro-aniline and saccharin did not interfere with the indomethacin and 9-
33
34 phenylcarbazone retention times.
35
36
37

38 **Dissolution Studies.** For the dissolution studies, the samples were micronized and sieved using
39
40 stainless steel standard-mesh sieves (mesh size 106 μm). In each experiment, the solid powders were
41
42 added to 12 ml of phosphate buffer 200 mM and incubated at 37°C under gentle shaking (100 rpm) in
43
44 a water bath. The amounts of sieved samples added to the buffer solution were 57.6 mg of
45
46 indomethacin; 75.2 mg of co-crystal *1*; 84.7 of co-crystal *2*; 86.9 of co-crystal *3*; 57.6 mg of
47
48 indomethacin mixed with 17.6 of 2-hydroxy-4-methyl-piridine, or 27.1 mg of 2-methoxy-5-nitro-
49
50 aniline or 29.3 mg of saccharin for mixtures *1*, *2* or *3*, respectively. Aliquots (200 μL) were withdrawn
51
52 from the resulting slurry at fixed time intervals and filtered through regenerated cellulose filters (0.45
53
54
55
56
57
58
59
60

1
2
3 μm). The resulting filtered samples were diluted 1:10 in water, then 10 μl was injected into the HPLC
4
5
6 system in order to quantify the indomethacin concentrations.
7

8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
Dissolution experiments were conducted also in PBS 10 mM at 37°C with the same procedure
above described, with the only difference that the filtered samples obtained from the slurry of mixture
3 were not diluted 1:10, but directly injected into the HPLC system. All the values obtained were the
mean of three independent experiments.

Cell Culture. The NCM460 cell line was grown in DMEM + Glutamax supplemented with 10 %
fetal bovine serum (FBS), 100 U/mL penicillin and 100 g/mL streptomycin at 37 °C in a humidified
atmosphere of 95%, with 5% of CO₂. For maximum viability, NCM460 cells were subcultured in fresh
and spent growth medium in 1:1 ratio. All cell culture reagents were provided by Invitrogen (Life
Technologies, Milan, Italy).

Differentiation of NCM460 Cells to Polarized Monolayers. Differentiation to NCM460 cell
monolayers was performed modifying the method reported by Dalpiaz and co-workers.²⁵ Briefly, after
two passages, confluent NCM460 cells were seeded at a density of 10⁵ cells/mL in 1:1 ratio fresh and
spent culture medium in 12-well Millicell inserts (Millipore, Milan, Italy) consisting of 1.0 μm pore
size polyethylene terephthalate (PET) filter membranes, whose surface was 1.12 cm². Filters were
presoaked for 24 h with fresh culture medium, and then the upper compartment (apical, A) received
400 μL of the diluted cells, whereas the lower (basolateral, B) received 2 mL of the medium in the
absence of cells. Half volume of the culture medium was replaced every two days with fresh medium to
each of the apical and basolateral compartments. The integrity of the cell monolayers was monitored by
measuring the transepithelial electrical resistance (TEER) by means of a voltmeter (Millicell-ERS;
Millipore, Milan, Italy). The measured resistance value was multiplied by the area of the filter to obtain
an absolute value of TEER, expressed as $\Omega\cdot\text{cm}^2$. The background resistance of blank inserts not plated
with cells was around 35 $\Omega\cdot\text{cm}^2$ and was deducted from each value. The homogeneity and integrity of

1
2
3 the cell monolayer were also monitored by phase contrast microscopy. Based on these parameters, cell
4
5 monolayers reached confluence and epithelial polarization after 6 days and monolayers with TEER
6
7 stable value around $180 \Omega \cdot \text{cm}^2$ were used for permeation studies. At this time, the medium was
8
9 replaced with low serum fresh medium (1 % FBS) in both the apical and basal compartments.
10
11

12 **Permeation Studies Across Cell Monolayers.** For permeation studies, inserts were washed twice
13
14 with pre-warmed PBS buffer in the apical compartment (A, 400 μL) and basolateral (B, 2 ml), then
15
16 PBS buffer containing 5 mM glucose at 37°C was added to the apical compartment. The sieved
17
18 powders (mesh size 106 μm) were added to the apical compartments in the following amounts: 1.92 mg
19
20 of indomethacin; 2.5 mg of co-crystal *1*; 2.8 mg of co-crystal *2*; 2.9 of mg co-crystal *3*; 1.92 mg of
21
22 indomethacin mixed to 0.59 mg of 2-hydroxy- 4 -methyl- pyridine, or 0.90 mg of 2 -methoxy-5-nitro-
23
24 aniline, or 0.98 mg of saccharin for mixtures *1*, *2* or *3*, respectively. During permeation experiments,
25
26 Millicell inserts loaded with the powders were continuously swirled on an orbital shaker (100 rpm;
27
28 model 711/CT, ASAL, Cernusco, Milan, Italy) at 37°C . At programmed time points the inserts were
29
30 removed and transferred into the subsequent wells containing fresh PBS, then basolateral PBS was
31
32 harvested, filtered through regenerated cellulose filters (0.45 μm) and injected (10 μL) into the HPLC
33
34 system for the determination of the concentration of indomethacin.
35
36
37
38
39
40

41 At the end of incubation the apical slurries were withdrawn, filtered and injected into HPLC
42
43 system (10 μl) after 1:10 dilution, with the exception of the apical sample of the mixture *3* that was
44
45 directly injected after filtration, without dilution. After the withdrawn of apical samples, 400 μl of PBS
46
47 were inserted in the apical compartments and TEER measurements were performed.
48
49 Permeation experiments were also conducted using cell-free inserts in the same conditions described
50
51 above. All the values obtained were the mean of three independent experiments.
52
53
54
55
56
57
58
59
60

1
2
3 Apparent permeability coefficients (P_{app}) of indomethacin were calculated according to the
4
5 following equation:²⁶⁻²⁸
6
7

$$P_{app} = \frac{\frac{dc}{dt} V_r}{S_A C} \quad (1)$$

8
9
10
11

12 where P_{app} is the apparent permeability coefficient in cm/min; dc/dt is the flux of drug across the filters,
13 calculated as the linearly regressed slope through linear data; V_r is the volume in the receiving
14 compartment (basolateral = 2 mL); S_A is the diffusion area (1.13 cm²); and C is the compound
15 concentration in the donor chamber (apical) detected at 60 min and chosen as approximate apical
16 concentration.
17
18
19
20
21
22
23

24 **Statistical Analysis about Permeation Studies.** Statistical comparisons between apparent
25 permeability coefficients or between apical concentrations of indomethacin were performed by one
26 way ANOVA followed by Dunnett's post-test; statistical comparisons between transepithelial electrical
27 resistance before and after incubation with the sieved samples was performed by one way ANOVA
28 followed by Bonferroni post-test. $P < 0.001$ was considered statistically significant. All the
29 calculations were performed by using the computer program Graph Pad Prism (GraphPad Software
30 Incorporated, La Jolla, CA, USA) that was employed also for the linear regression of the cumulative
31 amounts of the compounds in the basolateral compartments of the Millicell systems. The quality of fit
32 was determined by evaluating the correlation coefficients (r) and P values.
33
34
35
36
37
38
39
40
41
42
43
44

45 ***In Vivo* Administration of Indomethacin: Intravenous Infusion.** Male Sprague Dawley rats
46 (200–250 g) kept fasting since 24 hours received a femoral intravenous infusion of 0.90 mg/mL
47 indomethacin dissolved in a medium constituted by 20% (v/v) DMSO and 80% (v/v) physiologic
48 solution, with a rate of 0.2 mL/min for 5 min. Four rats were employed for femoral intravenous
49 infusions. At the end of infusion and at fixed time points within 24 hours blood samples (300 μ L) were
50 collected and inserted in heparinized test tubes that were centrifuged at 4°C for 15 min at 1,500 x g;
51
52
53
54
55
56
57
58
59
60

1
2
3 100 μL of plasma were then withdrawn and immediately quenched in 300 μL of ethanol (4 $^{\circ}\text{C}$); 100 μL
4
5 of internal standard (100 μM 9-phenyl-carbazole dissolved in ethanol) was then added. After
6
7 centrifugation at 13,000 $\times g$ for 10 min, 400 μL aliquots were reduced to dryness under a nitrogen
8
9 stream and stored at -20°C until analysis. The samples were dissolved in 150 μL of mobile phase
10
11 (methanol and 0.2 phosphoric acid 75:25 v/v), and, after centrifugation, 10 μL was injected into the
12
13 HPLC system for indomethacin assay. All the values obtained were the mean of four independent
14
15 experiments.
16
17
18

19
20 The efficacy of indomethacin extraction from blood samples was determined by recovery
21
22 experiments, comparing the peak areas extracted from 10 μM (3.58 $\mu\text{g}/\text{ml}$) blood test samples at 4 $^{\circ}\text{C}$
23
24 with those obtained by injection of an equivalent concentration of the drug dissolved in their mobile
25
26 phase. The average recovery \pm SD of indomethacin from rat blood resulted $87,4 \pm 3.9\%$. The
27
28 concentrations of this compound were therefore referred to as peak area ratio with respect to the
29
30 internal standard 9-phenyl-carbazole. The precision of the method based on peak area ratio, calculated
31
32 for 10 μM (3.6 $\mu\text{g}/\text{ml}$) solutions, was represented by RSD values of 0.93% . The calibration of
33
34 indomethacin was performed by employing eight different concentrations in whole blood at 4 $^{\circ}\text{C}$
35
36 ranging from 2 μM (0.72 $\mu\text{g}/\text{ml}$) to 50 μM (18.0 $\mu\text{g}/\text{ml}$) and expressed as peak area ratios of the
37
38 compounds to the internal standard versus concentration. The calibration curve resulted linear ($n = 8$, r
39
40 $= 0.990$, $P < 0.0001$). The accuracy of extraction method was determined with respect to the calibration
41
42 curve and was described by relative errors comprised between -2.63% and 0.24% .
43
44
45
46
47

48
49 The *in vivo* half-life of indomethacin in the blood was calculated by nonlinear regression
50
51 (exponential decay) of concentration values in the time range within 24 hours after infusion and
52
53 confirmed by linear regression of the log concentration values versus time. The area under the
54
55 concentration-time curve (AUC) value was calculated by the trapezoidal method within 24 hours, the
56
57 remaining area was determined as the ratio between the indomethacin concentration detected at 24 hour
58
59
60

1
2
3 and the elimination constant (k_{el}), that was obtained from the slope of the semilogarithmic (-slope ·
4
5
6 2.3). All the calculations were performed by using the computer program Graph Pad Prism.
7

8 ***In Vivo Administration: Oral Administration of Indomethacin, its Co-Crystals and Parent***
9
10 **Mixtures.** The sieved powders were mixed with palatable food in order to induce their oral assumption
11
12 by male Sprague Dawley rats (200–250 g) kept fasting since 24 hours. The following doses were
13
14 administered: 0.90 mg of indomethacin; 1.18 mg of co-crystal *1*; 1.32 mg of co-crystal *2*; 1.36 mg of
15
16 co-crystal *3*; 0.90 mg of indomethacin mixed to 0.28 mg of 2-hydroxy-4 -methyl-pyridine, or 0.42 mg
17
18 of 2 -methoxy-5-nitro-aniline, or 0.46 mg of saccharin for mixtures *1*, *2* or *3*, respectively. Four
19
20 rats/group were employed for the oral administration experiments. At the end of administration and at
21
22 fixed time points within 24 hours blood samples (300 μ L) were collected, then extracted and analyzed
23
24 as above described. All the concentration values obtained for indomethacin were the mean of four
25
26 independent experiments. The AUC values referred to each orally administered treatment were
27
28 calculated as above described. The absolute bioavailability values of indomethacin, referred to the oral
29
30 administered samples, were obtained as the ratio between their oral AUC values and AUC of the
31
32 intravenous administration of the drug. All the calculations were performed by using the computer
33
34 program Graph Pad Prism.
35
36
37
38
39

40 **Statistical Analysis about *in Vivo Administration of Indomethacin.*** Statistical comparisons
41
42 between absolute bioavailability values were performed by one way ANOVA followed by Dunnet
43
44 post-test. $P < 0.001$ was considered statistically significant. All the calculations were performed by
45
46 using the computer program Graph Pad Prism.
47
48
49

50 51 52 53 **RESULTS** 54 55 56 57 58 59 60

1
2
3 **Indomethacin Co-Crystals.** Indomethacin was co-crystallized with two model molecules, 2-
4 hydroxy-4-methyl-pyridine (co-crystal *1*), 2-methoxy-5-nitroaniline (co-crystal *2*). Their chemical
5 structures along with that of the previously reported^{11,19} indomethacin-saccharin co-crystal (co-crystal
6 *3*) are reported in Figure 1. The X-ray three-dimensional structures for the three co-crystals are shown
7 in Figure 2. The hydrogen bonds between indomethacin and co-crystallizing agents are evidenced by
8 dashed lines. Figure 2 evidences that the covalent bonds of each single molecule are not altered in the
9 co-crystallized structures.
10
11
12
13
14
15
16
17
18

19 **Dissolution Studies: Co-Crystals Can Significantly Change the Dissolution Profiles of**
20 **Indomethacin.** Figure 3A reports a comparison between the dissolution profiles in 200 mM phosphate
21 buffer at 37°C of γ -indomethacin, as free drug, co-crystallized or mixed in the parent mixtures. The
22 saturation concentration of free γ -indomethacin, reached after about two hours of its incubation into the
23 buffer, was about 1.8 mg/ml. The dissolution profile of γ -indomethacin was not altered by the presence
24 of the co-crystallizing agents when mixed with the drug. Differently, the co-crystallized powders
25 induced significant changes of indomethacin dissolution profiles. In particular, the co-crystal *1* and the
26 co-crystal *2* induced an increase up to three times and a 50% decrease of γ -indomethacin saturation
27 concentration, respectively, without affecting the dissolution rate. Finally, the co-crystal *3* induced not
28 only a significant enhancement of the saturation concentration of γ -indomethacin (up to four times) but,
29 differently from the other co-crystals, also an increase of the drug dissolution rate, being 30 min the
30 time necessary to reach the saturation conditions.
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

48 The indomethacin solubility and dissolution profiles represented in Figure 3A drastically
49 changed when the powders were incubated in PBS 10 mM, the medium employed for the drug
50 permeation studies across NCM460 cell monolayers (see below). In particular, under these
51 experimental conditions, the saturation concentration of free γ -indomethacin was reduced of about 50%
52
53
54
55
56
57
58
59
60

1
2
3 when compared to that measured when it was dissolved in 200 mM phosphate buffer (Figure 3B); the
4
5 mixtures **1** and **2** showed dissolution profiles similar to that of free γ -indomethacin, whereas the
6
7
8 mixture **3** induced a drastic decrease of the saturation concentration of the drug, showing a mean \pm S.D
9
10 value of 0.0012 ± 0.0004 mg/ml (about 0.2% of the saturation concentration of the free drug). The co-
11
12 crystal **1** was instead associated to an increase of indomethacin saturation concentration. Differently
13
14 from the results obtained under 200 mM phosphate buffer conditions (see above), the co-crystal **2** led
15
16 to an indomethacin dissolution profile similar to those displayed by the free drug. Finally, the co-
17
18 crystal **3** induced a very fast dissolution of indomethacin that was, however, followed by a sudden
19
20 precipitation of the drug; this event was completed within 60 minutes, showing indomethacin
21
22 concentration values about 0.003 mg/ml.
23
24
25
26

27 **Permeation Studies: Co-Crystals Can Induce Different Effects on Cell Monolayers with** 28 29 **Respect to their Parent Mixtures**

30
31
32 The PBS was used as incubation medium for the permeation studies of indomethacin across an
33
34 *in vitro* model of human intestinal wall, i.e. NCM460 cell monolayers.²⁹ In order to simulate an oral
35
36 administration, the powders of γ -indomethacin, its co-crystals or the parent mixtures were introduced in
37
38 the apical compartment of the “millicell” systems with the same ratio between solid powders and
39
40 incubation medium adopted for dissolution studies. The indomethacin permeation profiles, expressed
41
42 by the cumulative concentrations in the receiving basolateral compartments are reported in Figure 4.
43
44 The linear profiles indicate constant permeation conditions during the analysis time period (60 min) for
45
46 all samples. The straight line related to mixture **3** ($r = 0.970$, $P = 0.001$) was characterized by
47
48 indomethacin concentration values strongly lower than those of the straight lines of the other powders
49
50 ($r \geq 0.998$, $P < 0.0001$). The apparent permeability coefficients (P_{app}) of indomethacin (Table 1) have
51
52 been calculated on the basis of the resulting slopes of the linear fits and the indomethacin
53
54 concentrations detected in the apical compartments after one hour of incubation of the powders (Table
55
56
57
58
59
60

1
2
3 1), chosen as approximate apical concentrations. These latter values appeared essentially in line with
4
5 those obtained from dissolution studies of indomethacin powders in 10 mM PBS (Figure 3B). Indeed,
6
7 the drug concentrations obtained from the powders constituted by γ -indomethacin, mixtures **1** and **2**
8
9 and co-crystal **2** were not statistically dissimilar among them (about 1000 μ M, $P > 0.05$). On the other
10
11 hand, mixture **3** induced a drastic reduction of indomethacin concentration ($P < 0.001$), showing values
12
13 close to the drug limit of quantification. Furthermore, the co-crystal **1** induced an increase of
14
15 indomethacin concentration of about three times with respect to the free drug powder ($P < 0.001$),
16
17 whereas the co-crystal **3** significantly reduced the γ -indomethacin concentration ($P < 0.001$), even if in
18
19 a less drastic manner than detected from dissolution experiments in 10 mM PBS in the absence of cells
20
21 (Figure 3B). The slope of the permeation profile (Figure 4) and the apical concentration of
22
23 indomethacin related to mixture **3** (Table 1) appeared too low to obtain a reliable P_{app} value of
24
25 indomethacin dissolved from this sample.
26
27
28
29
30
31

32 A comparison of the P_{app} values of γ -indomethacin (Table 1) obtained in the presence ($150 \cdot 10^{-5}$
33
34 $\pm 6 \cdot 10^{-5}$ cm/min) and in the absence ($429 \cdot 10^{-5} \pm 18 \cdot 10^{-5}$ cm/min) of NCM460 cell monolayers
35
36 indicated a significant lower permeation of the drug in the presence of cells ($P < 0.001$), confirming the
37
38 validity of the monolayer as an *in vitro* model of a physiologic barrier. This behavior appeared in
39
40 agreement with the transepithelial electrical resistance (TEER) values (about $180 \Omega \cdot \text{cm}^2$) attributed to
41
42 the monolayers before their incubation with the powders (Table 1). The apparent coefficient values
43
44 across the monolayer of indomethacin dissolved from mixtures **1** and **2** and co-crystal **2** did not
45
46 significantly differ from the P_{app} obtained for the free γ -indomethacin (Table 1, $P > 0.05$). On the other
47
48 hand, the co-crystal **1** induced a consistent increase of indomethacin permeation ($P_{app} = 301 \cdot 10^{-5} \pm 11$
49
50 $\cdot 10^{-5}$ cm/min, $P < 0.001$). This phenomenon was associated to the ability of this co-crystal to impair the
51
52 tight junctions among the cells of the monolayer. This ability was evidenced by the drastic reduction of
53
54
55
56
57
58
59
60

1
2
3 TEER values ($P < 0.001$) measured after 60 min of incubation with this sample ($21 \pm 1 \Omega \cdot \text{cm}^2$) in
4
5 comparison to the value obtained before incubation (185 ± 11). Moreover, by monitoring the
6
7 monolayer after its incubation with co-crystal **1** by phase contrast microscopy evidenced a complete
8
9 separation of the cells (*data not shown*). It is remarkable that mixture **1** incubation did not alter neither
10
11 the monolayer integrity, as monitored by phase contrast microscopy (*data not shown*) nor the TEER
12
13 value (before incubation = $181 \pm 9 \Omega \cdot \text{cm}^2$; after incubation = $158 \pm 7 \Omega \cdot \text{cm}^2$; $P > 0.05$). The same
14
15 profile has been also registered for the powder of free γ -indomethacin, the mixture **2** and the co-crystals
16
17 **2** and **3**. It is interesting to observe that the mixture **3** induced a cell monolayer fragmentation, as
18
19 monitored by phase contrast microscopy (*data not shown*) and indicated by the TEER value (before
20
21 incubation = $180 \pm 10 \Omega \cdot \text{cm}^2$; after incubation = $36 \pm 2 \Omega \cdot \text{cm}^2$; $P < 0.001$). The co-crystal **3** has been
22
23 characterized by a P_{app} value of $374 \cdot 10^{-5} \pm 16 \cdot 10^{-5} \text{ cm/min}$, significantly higher with respect to that
24
25 obtained with the powder of free γ -indomethacin ($P < 0.001$). Interestingly, the permeation
26
27 enhancement induced by co-crystal **3** was not related with the monolayer fragmentation (*data not*
28
29 *shown*).

30
31
32
33
34
35
36
37 ***In Vivo* Administration of Indomethacin: its Oral Bioavailability is Modulated by Co-**
38
39 **crystallization.** After intravenous infusion of 0.90 mg indomethacin, the drug concentration in the rat
40
41 bloodstream was $13.2 \pm 1.4 \mu\text{g/ml}$. This value decreased during time with an apparent first order
42
43 kinetic (Figure 5A) confirmed by the linearity of the semilogarithmic plot reported in the inset of
44
45 Figure 5A ($n = 8$, $r = 0.983$, $P < 0.0001$), showing an half-life value of 8.94 ± 0.38 hours.
46
47

48
49 The rat blood indomethacin concentrations within 24 hours after its intravenous infusion (IV) as
50
51 free-drug or the oral administration of 0.90 mg indomethacin as sieved powders of free γ -drug, its co-
52
53 crystals and the parent mixtures are reported in Figure 5B. In order to better compare among them the
54
55 results, Figure 5C reports a section of Figure 5B, focused on the profiles obtained within eight hours
56
57
58
59
60

1
2
3 after of the oral administration of the powders. It can be observed that the free γ -indomethacin powder
4
5 induced a concentration peak in the rat bloodstream of about 5 $\mu\text{g/ml}$ two hours after the
6
7 administration. A similar profile was obtained with mixtures *1* and *2*, whereas mixture *3* was
8
9 characterized by a profile showing a peak concentration of about 3.5 $\mu\text{g/ml}$ two hours after its
10
11 administration.
12
13

14
15 The co-crystal *1* induced a concentration peak in the rat bloodstream of about 5 $\mu\text{g/ml}$ 30
16
17 minutes after the administration, the same time required for co-crystal *3* to induce a concentration peak
18
19 of about 3 $\mu\text{g/ml}$. The co-crystal *2* profile was instead characterized by a peak lower than 2 $\mu\text{g/ml}$
20
21 obtained between 1 and 2 hours after the administration. In general, the profiles of the co-crystals
22
23 appeared characterized by a decrease of indomethacin blood concentration within 4 hours after their
24
25 administration with a rate lower than those observed following free γ -indomethacin and the parent
26
27 mixtures administrations.
28
29
30
31

32
33 The AUC values of the profiles reported in Figure 5B were employed for the calculation of
34
35 absolute bioavailabilities (F) of the solid formulations, reported in Table 1. In particular the free γ -
36
37 indomethacin powder was characterized by an F value of $23.5 \pm 0.8\%$, not statistically dissimilar ($P >$
38
39 0.05) from those of mixtures *1*, *2* and *3*. The co-crystals *1* and *3* were characterized by a significant
40
41 higher ($P < 0.001$) F values than free γ -indomethacin ($37.7 \pm 1.5\%$ and $33.58 \pm 0.59\%$, respectively),
42
43 while, co-crystal *2* induced a relative small, but significant ($P < 0.001$) decrease of bioavailability with
44
45 respect to the powder of the free drug (F value = $20.1 \pm 0.9\%$).
46
47
48
49
50

51 DISCUSSION

52
53
54
55
56
57
58
59
60

1
2
3 Co-crystallization, by enhancing BCS class II API solubility, has been proposed as a new
4 strategy to increase drug bioavailability.⁹⁻¹⁷ However, experimental evidence about the effects of the
5 co-crystals on the permeation of APIs across intestinal barriers and on intestinal epithelial barrier
6 integrity is not reported in literature. These studies should be relevant as the disruption of intestinal
7 epithelial tight junctions has two undesirable consequences: unwanted substances such as endotoxins
8 are allowed into the body, and depolarization of cells may be promoted.^{30,31} Moreover, the regulatory
9 status regarding the use of co-crystals in pharmaceutical products appears still unsettled, being
10 necessary to clarify whether the co-crystal would be defined as a physical mixture or as a new chemical
11 entity requiring full safety and toxicology testing.^{7,8} In the attempt to contribute to clarify these aspects,
12 we have chosen indomethacin as a model BCS class II API and compared its properties with those of
13 three of its co-crystals along with their parent physical mixtures. To our knowledge, this type of study
14 is absolutely novel, being not only focused to a systematic comparison among the behavior of powders
15 constituted by the pure API, its co-crystals and their parent mixtures, but also involving the analysis of
16 the API permeation across an *in vitro* intestinal barrier model.
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35

36 The dissolution studies were firstly performed in a 200 mM phosphate buffer, pH 7.4. The
37 relative high ionic strength of this buffer was necessary to maintain the stability of pH value during the
38 dissolution processes.¹⁹ As already described, we have observed that the dissolution profile of
39 indomethacin was not altered by the presence of the co-crystallizing agents in the form of physical
40 mixtures with the API, whereas significant changes were observed by the dissolution of the co-crystals.
41 A qualitative concordance between the API dissolution patterns in the 200 mM phosphate buffer and its
42 absorption in the rat bloodstream after the oral administration of the powders, has been observed. In
43 particular, dissolution (Figure 3A) and bioavailability (Figure 5C) profiles of pure γ -indomethacin were
44 similar to those of its physical mixtures with the co-crystallizing molecules; on the other hand either
45 indomethacin solubility or its bioavailability were significantly increased by co-crystals **1** and **3**, and
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 weakly decreased by co-crystal 2. It is worth noting that, to allow a direct comparison among the
4 powder constituted by the free drug, its co-crystals and the parent mixtures we intentionally decided to
5 do not follow the suitable formulation strategies suggested to improve the plasma levels of APIs orally
6 administered as co-crystals.⁹ This can explain the relatively weak, although significant, changes of
7 bioavailability induced by the co-crystallization observed in the present study. However, the
8 accordance between dissolution and bioavailability profiles above described is in line with literature
9 data concerning pharmaceutical co-crystals.¹⁰⁻¹⁷ Since this correlation does not provide any information
10 on the effective role of the co-crystals in influencing the API absorption mechanisms across the
11 intestinal barrier, we decided to perform permeation experiments across NCM460 cell monolayers.
12 These studies are absolutely innovative for systems involving pharmaceutical co-crystals and their
13 parent mixtures.
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28

29 Various cell monolayer models that mimic the human intestinal epithelial barrier have been
30 developed, providing ideal systems for the rapid *in vitro* assessment of the intestinal absorption of drug
31 candidates. We have chosen the human normal colonic epithelial NCM460 cells being them an
32 immortalized, non-transformed cell line, derived from primary cells of the normal human transverse
33 colonic mucosa.²⁹ As these cells are not of tumor origin neither transfected ones, they retain more
34 closely the physiological characteristics of the normal human colon compared to the pathologically or
35 experimentally transformed cell lines. In this context, it is worth noting that TEER developed by the
36 NCM460 cells are within the range reported for intact sheets of human colonic mucosa.^{32,33}
37
38 Furthermore, the lipophilic nature of indomethacin enables the molecule to diffuse quickly and to get
39 absorbed completely through the intestinal membrane after oral ingestion, resulting almost equally
40 permeable in the colon and small intestine.^{34,35}
41
42
43
44
45
46
47
48
49
50
51
52
53
54

55 The permeation studies were performed by glucose-enriched PBS as dissolution medium of the
56 indomethacin powder, being the concentration of 200 mM phosphate buffer too high to allow the cell
57
58
59
60

1
2
3 survival. Moreover, PBS represented the simplest medium where to dissolve indomethacin from its
4
5 powders, in order to study the permeation properties across NCM460 cells in the absence of other
6
7 interfering substances. It is indeed known that simulated intestinal buffers can induce TEER changes of
8
9 the monolayers and have inhibitory activity towards efflux transporters expressed on the cell
10
11 membranes.³⁶ As described above, the dissolution profiles in PBS of indomethacin showed some
12
13 marked differences with respect to the patterns obtained in 200 mM phosphate buffer, attributable to
14
15 the PBS relatively weak buffering power.
16
17
18

19
20 The suspensions obtained by the introduction of the solid powders containing indomethacin in
21
22 the apical compartments of the “millicell” systems allowed us to simulate an oral administration. The γ -
23
24 indomethacin crystals appeared able to maintain the integrity of the monolayer characterized by TEER
25
26 values around $180 \Omega \cdot \text{cm}^2$. Moreover, a comparison of the permeability values obtained in the absence
27
28 and in the presence of cell monolayer validated the ability of this preparation to behave as a
29
30 physiologic barrier. We have instead observed that the physical mixture of indomethacin and saccharin
31
32 induced a drastic decrease of the TEER value of the monolayer, whose cells appeared to loose
33
34 completely their mutual contacts after one hour of incubation. Surprisingly, the incubation with the
35
36 parent co-crystal **3** allowed to maintain the integrity of the monolayer as well as of its TEER value, and
37
38 induced an increase of indomethacin permeation across the NCM460 cells, with respect to γ -
39
40 indomethacin crystals.
41
42
43
44

45
46 We have also observed that co-crystal **1** induced a drastic decrease of the TEER value of the
47
48 monolayer, whose cells appeared completely separated after one hour of incubation. The permeation
49
50 profile of indomethacin dissolved from co-crystal **1** showed indeed the highest values with respect to
51
52 all other cases, probably due to both the loss of the barrier effect of the monolayer and the increased
53
54 dissolution of the API. Surprisingly, the incubation with the parent physical mixture **1** did not induce
55
56 any changes on the monolayer integrity, as evidenced by unaffected TEER value after one hour of
57
58
59
60

1
2
3 incubation. Moreover, the permeation profile of indomethacin dissolved from this mixture was the
4
5 same to that obtained with solid γ -indomethacin. No significant differences between γ -indomethacin
6
7 and co-crystal **2** or mixture **2** were observed, as far as the integrity of the monolayer and API
8
9 permeation profile is concerned.
10
11

12 13 14 15 **CONCLUSIONS**

16
17
18
19
20 To the best of our knowledge this is the first study demonstrating different effects induced by
21
22 co-crystals and their parent physical mixtures on a biologic system, findings that could raise serious
23
24 concerns about the use of co-crystal strategy to improve API bioavailability without performing
25
26 appropriate investigations. In this case co-crystal **1** was found to induce a drastic decrease of the TEER
27
28 value NCM460 cell monolayers, whereas its parent mixture did not evidence any effect. On the other
29
30 hand, the physical mixture of saccharin and indomethacin was able to induce a drastic decrease of the
31
32 TEER value of NCM460 monolayers, whereas its parent co-crystal **3** did not evidence any effect on
33
34 the integrity of the monolayers, being anyway able to increase the permeation of indomethacin across
35
36 the monolayers. On the basis of the present experimental data we cannot explain the reason(s) of these
37
38 phenomena, but it is clearly evidenced that the biological effects of a co-crystal and its parent mixture
39
40 can be drastically different, even if this is not to be taken as a general rule. Indeed, any difference was
41
42 registered by our permeation measurements between co-crystal **2** and its parent physical mixture on
43
44 NCM460 cell monolayer.
45
46
47
48
49

50
51 Our results seem to open new perspectives about the application of pharmaceutical products
52
53 containing co-crystals. New and appropriate investigations appear therefore necessary in order to
54
55 evaluate the potential new applications and the potential damaging effects of pharmaceutical co-
56
57 crystals.
58
59
60

AUTHOR INFORMATION

Corresponding Author

*Department of Chemical and Pharmaceutical Sciences, University of Ferrara, via Fossato di Mortara 19, I-44121, Ferrara, Italy. Phone: +39-0532-455274. Fax: +39-0532- 455953. E-mail: dla@unife.it.

REFERENCES

(1) Dressman, J.; Reppas, C. Drug Solubility: How to Measure it, how to Improve it. *Adv. Drug Del. Rev.* **2007**, 59, 531-532.

(2) Morissette, S. L.; Soukasene, S.; Levinson, D.; Cima, M. J.; Almarsson O. Elucidation of Crystal form Diversity of the HIV Protease Inhibitor Ritonavir by High-Throughput crystallization. *Proc. Natl. Acad. Sci. U.S.A* **2003**, 100, 2180–2184.

(3) Heinz, A.; Strachan, C. J.; Gordon, K. C.; Rades, T. Analysis of Solid-State Transformations of Pharmaceutical Compounds Using Vibrational Spectroscopy. *J. Pharm. Pharmacol.* **2009**, 61, 971–988.

(4) Schultheiss, N.; Newman, A. Pharmaceutical Cocrystals and their Physicochemical Properties. *Cryst. Growth Des.* **2009**, 9, 2950–2967.

(5) Jones, W.; Motherwell, W. D. S.; Trask, A. V. Pharmaceutical Cocrystals: an Emerging Approach to Physical Property Enhancement. *MRS Bulletin* **2006**, 31, 875-879.

(6) Remenar, J. F.; Morissette, S. L.; Peterson, M. L.; Moulton, B.; MacPhee, J. M.; Guzman, H. R.; Almarsson, O. Crystal Engineering of Novel Cocrystals of a Triazole Drug with 1,4-Dicarboxylic Acids. *J. Am. Chem. Soc.* **2003**, 125, 8456-8457.

- 1
2
3 (7) Brittain, H. G. Cocrystal Systems of Pharmaceutical Interest: 2010. *Cryst. Growth Des.*
4
5 **2012**, 12, 1046–1054.
6
7
8 (8) Brittain, H. G. Cocrystal Systems of Pharmaceutical Interest: 2011. *Cryst. Growth Des.*
9
10 **2012**, 12, 5823–5832.
11
12 (9) Childs, S. L.; Kandi, P.; Lingireddy, S. R. Formulation of a Danazol Cocrystal with
13 Controlled Supersaturation Plays an Essential role in Improving Bioavailability. *Mol. Pharm.* **2013**, 10,
14 3112–3127.
15
16
17 (10) Hickey, M. B.; Peterson, M. L.; Scoppettuolo, L. A.; Morrisette, S. L.; Vetter, A.; Guzman,
18 H.; Remenar, J. F.; Zhang, Z.; Tawa, M. D.; Haley, S.; Zaworotko, M. J.; Almarsson, O. Performance
19 Comparison of a Cocrystal of Carbamazepine with Marketed Product. *Eur. J. Pharm. Biopharm.* **2007**,
20 67, 112–119.
21
22 (11) Jung, M. S.; Kim, J. S.; Kim, M. S.; Alhalaweh, A.; Cho, W.; Hwang, S. J.; Velaga, S. P.
23 Bioavailability of Indomethacin-Saccharin Cocrystals. *J. Pharm. Pharmacol.* **2010**, 62, 1560–1568.
24
25 (12) Weyna, D. R.; Cheney, M. L.; Shan, N.; Hanna, M.; Zaworotko, M. J.; Sava, V.; Song, S.;
26 Sanchez-Ramos, J. R. Improving Solubility and Pharmacokinetics of Meloxicam via Multiple-
27 Component Crystal Formation. *Mol. Pharm.* **2012**, 9, 2094–2102.
28
29 (13) Smith, A. J.; Kavuru, P.; Wojtas, L.; Zaworotko, M. J.; Shytle, R. D. Cocrystals of
30 Quercetin with Improved Solubility and Oral Bioavailability. *Mol. Pharm.* **2011**, 8, 1867–1876.
31
32 (14) McNamara, D. P.; Childs, S. L.; Giordano, J.; Iarriccio, A.; Cassidy, J.; Shet, M. S.;
33 Mannion, R.; O'Donnell, E.; Park, A. Use of a Glutaric Acid Cocrystal to Improve Oral Bioavailability
34 of a low Solubility API. *Pharm. Res.* **2006**, 23, 1888–1897.
35
36 (15) Bak, A.; Gore, A.; Yanez, E.; Stanton, M.; Tufekcic, S.; Syed, R.; Akrami, A.; Rose, M.;
37 Surapaneni, S.; Bostick, T.; King, A.; Neervannan, S.; Ostovic, D.; Koparkar, A. The Co-Crystal
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 Approach to Improve the Exposure of a Water-Insoluble Compound: AMG 517 Sorbic Acid Co-
4
5 Crystal Characterization and Pharmacokinetics. *J. Pharm. Sci.* **2008**, 97, 3942–3956.

6
7
8 (16) Stanton, M. K.; Kelly, R. C.; Colletti, A.; Langley, M.; Munson, E. J.; Peterson, M. L.;
9
10 Roberts, J.; Wells, M. Improved Pharmacokinetics of AMG 517 through Co-Crystallization part 2:
11
12 Analysis of 12 Carboxylic Acid Co-Crystals. *J. Pharm. Sci.* **2011**, 100, 2734–2743.

13
14
15 (17) Zheng, W.; Jain, A.; Papoutsakis, D.; Dannenfelser, R. M.; Panicucci, R.; Garad, S.
16
17 Selection of Oral Bioavailability Enhancing Formulations during Drug Discovery. *Drug Dev. Ind.*
18
19 *Pharm.* **2012**, 38, 235–247.

20
21
22 (18) Bethune, S. J.; Schultheiss N.; Henck, J. O. Improving the Poor Aqueous Solubility of
23
24 Nutraceutical Compound Pterostilbene through Cocrystal Formation. *Cryst. Growth Des.* **2011**, 11,
25
26 2817–2823.

27
28
29 (19) Basavoju, S.; Bostrom, D.; Velaga, S. P. Indomethacin-Saccharin Cocrystals: Design,
30
31 Synthesis and Preliminary Pharmaceutical Characterization. *Pharm. Res.* **2007**, 25, 530-541.

32
33
34 (20) Otwinowski, Z.; Minor, W. Processing of X-Ray Diffraction Data Collected in Oscillation
35
36 Mode. In *Methods in Enzymology*, 276, *Macromolecular Crystallography*, Part A, Carter, C.W. Jr &
37
38 Sweet, R.M. Eds., Academic Press: New York, 1997; pp 307-326

39
40
41 (21) Altomare, A.; Burla, M. C.; Camalli, M.; Cascarano, G.; Giacovazzo, C.; Guagliardi, A.;
42
43 Moliterni, A. G.; Polidori, G.; Spagna, R. SIR97: a New Tool for Crystal Structure Determination and
44
45 Refinement. *J. Appl. Crystallogr.* **1999**, 32, 115–119.

46
47
48 (22) Sheldrick, G. M. *SHELX97, Program for Crystal Structure Refinement*; University of
49
50 Göttingen, Göttingen, Germany, 1997

51
52
53 (23) Farrugia, L. J. WinGX Suite for Small-Molecule Single-Crystal Crystallography. *J. Appl.*
54
55 *Crystallogr.* **1999**, 32, 837–838.

1
2
3 (24) Burnett, M. N.; Johnson, C. K. *ORTEPIII. Report ORNL-6895*; Oak Ridge National
4 Laboratory, Oak Ridge, Tennessee, USA, 1996.
5
6

7
8 (25) Dalpiaz, A.; Paganetto, G.; Pavan, B.; Fogagnolo, M.; Medici, A.; Beggiato, S.; Perrone,
9 D. Zidovudine and Ursodeoxycholic Acid Conjugation: Design of a New Prodrug Potentially Able to
10 Bypass the Active Efflux Transport Systems of the Central Nervous System. *Mol. Pharm.* **2012**, 9(4),
11 957-968.
12
13
14
15

16
17 (26) Artursson, P.; Karlson, J. Correlation Between Oral Absorption in Humans and Apparent
18 Drug Permeability Coefficients in Human Intestinal Epithelial (Caco-2) Cells. *Biochem. Biophys. Res.*
19 *Commun.* **1991**, 175, 880–885.
20
21
22
23

24 (27) Pal, D.; Udata, C.; Mitra, A. K. Transport of Cosalane, a Highly Lipophilic Novel Anti-
25 HIV Agent, across Caco-2 Cell Monolayers. *J. Pharm. Sci.* **2000**, 89, 826–833.
26
27
28

29 (28) Raje, S.; Cao, J.; Newman, A. H.; Gao, H.; Eddington, N. D. Evaluation of the Blood-
30 Brain Barrier Transport, Population Pharmacokinetics, and Brain Distribution of Benzotropine Analogs
31 and Cocaine Using in vitro and in vivo Techniques. *J. Pharmacol. Exp. Ther.* **2003**, 307, 801–808.
32
33
34
35

36 (29) Moyer, M. P.; Manzano, L.; Merriman, R.; Stauffer, J.; Tanzer, L. R. NCM460, a Normal
37 Human Colon Mucosal Epithelial Cell Line. *In Vitro Cell Dev. Biol. Anim.* **1996**, 32, 315–317.
38
39
40

41 (30) Laurent-Puig, P.; Blons, H.; Cugnenc, P. H. Sequence of Molecular Genetic Events in
42 Colorectal Tumorigenesis. *Eur. J. Cancer Prev.* **1999**, 8, Suppl 1, S39–S47.
43
44

45 (31) Rao, R. K.; Seth, A.; Sheth P. Recent Advances in Alcoholic Liver Disease I. Role of
46 Intestinal Permeability and Endotoxemia in Alcoholic Liver Disease. *Am. J Physiol. Gastrointest. Liver*
47 *Physiol.* **2004**, 286, G881–G884.
48
49
50

51 (32) Sahi, J.; Nataraja, S. G.; Layden, T. J.; Goldstein, J. L.; Moyer, M. P.; Rao M. C. Cl⁻
52 Transport in an Immortalized Human Epithelial Cell Line (NCM460) Derived from the Normal
53 Transverse Colon. *Am. J. Physiol.* **1998**, 275 (4Pt1), C1048-1057.
54
55
56
57
58
59
60

1
2
3 (33) Liu, Z. H.; Shen, T. Y.; Zhang P.; Ma, Y. L.; Moyer, M. P.; Qin, H. L. Protective Effects of
4 Lactobacillus Plantarum against Epithelial Barrier Dysfunction of Human Colon Cell Line NCM460.
5
6
7
8 *World J. Gastroenterol.* **2010**, 16, 5759-5765.

9
10 (34) ElShaer, A.; Hanson, P.; Mohammed, A. R. A Novel Concentration Dependent Amino
11 Acid Ion Pair Strategy to Mediate Drug Permeation using Indomethacin as a Model Insoluble Drug.
12
13
14
15 *Eur. J. Pharm. Sci.* **2014**, 62, 124-131.

16
17 (35) Ehrhardt, C.; Kim K. J. *Drug Absorption Studies: In Situ, In Vitro and In Silico Models.*
18
19 Springer: New York, 2008.

20
21 (36) Ingels, F.; Defermec, S.; Destexhe, E.; Oth, M., Van den Mooter, G.; Augustijns, P.
22 Simulated Intestinal Fluid as Transport Medium in the Caco-2 Cell Culture Model. *Int. J. Pharm.* **2002**,
23
24
25
26
27 232, 183–192.

Table 1. Data related to indomethacin *in vitro* permeation studies and *in vivo* oral bioavailability.

Permeation studies were performed by using “millicell” filters alone (filter) or coated by NCM460 cell monolayers (cells). Indomethacin was introduced in the donor compartment as sieved powder of γ -indomethacin, or its co-crystals, or the parent mixtures. The apical concentrations detected at the end of incubation were employed for the calculation of the apparent permeation coefficients (P_{app}). Permeation studies were performed after cell cultures reached the confluence using parallel sets of “millicell” well plates with similar TEER values (TEER 0 min). The TEER values were measured again at the end of incubation (TEER 60 min). All data related to permeation studies are reported as the mean \pm S.D. of three independent experiments. Bioavailability data were obtained after oral administration to rats of the powders and are reported as the mean \pm S.D. of four independent experiments.

Powder	Permeation condition	Apical concentrations at 60 min (μM)	P_{app} ($\cdot 10^{-5}$ cm/min)	TEER ($\Omega\cdot\text{cm}^2$) 0 min	TEER ($\Omega\cdot\text{cm}^2$) 60 min	Absolute Bioavailability (%)
γ -Indomethacin	Cells	1010 \pm 40	150 \pm 6	181 \pm 10	163 \pm 8	23.5 \pm 0.8
Mixture 1	Cells	989 \pm 42	165 \pm 7	181 \pm 9	158 \pm 7	25.7 \pm 1.6
Co-crystal 1	Cells	2708 \pm 84 (*)	301 \pm 14 (*)	185 \pm 11	21 \pm 1 (**)	37.7 \pm 1.5 (***)
Mixture 2	Cells	1047 \pm 50	147 \pm 10	183 \pm 10	152 \pm 7	24.4 \pm 1.2
Co-crystal 2	Cells	1013 \pm 38	145 \pm 6	177 \pm 9	159 \pm 8	20.1 \pm 0.9 (***)
Mixture 3	Cells	3.2 \pm 0.1 (*)	--	180 \pm 10	36 \pm 2 (**)	24.8 \pm 0.9
Cocrystal 3	Cells	442 \pm 18 (*)	374 \pm 16 (*)	183 \pm 10	164 \pm 8	33.6 \pm 0.6 (***)
γ -Indomethacin	Filter	1015 \pm 45	429 \pm 18 (*)	--	--	--

(*) $P < 0.001$ versus γ -indomethacin permeating across NCM460 cell monolayers

(**) $P < 0.001$ versus TEER at “time 0” (0 min)

(***) $P < 0.001$ versus absolute bioavailability of γ - indomethacin

Caption of Figures

Figure 1. Schematic representation of indomethacin and the co-formers 2-hydroxy-4-methyl-pyridine, 2-methoxy-5-nitroaniline and saccharine in co-crystals **1**, **2** and **3**, respectively.

Figure 2. (a) ORTEPIII view and atom numbering scheme for co-crystals **1**; (b) ORTEPIII view and atom numbering scheme for co-crystals **2**; (c) indomethacin-saccharine complex **3** (from Ref. 19).

Thermal ellipsoids are drawn at the 40% probability level. Hydrogen bonds are drawn as dashed lines.





Figure 3. Solubility and dissolution profiles in phosphate buffer 200 mM [A] and PBS 10 mM [B] at 37°C for γ -indomethacin as free drug, or co-crystallized, or mixed in the parent mixtures. Data are reported as the mean \pm S.D. of three independent experiments.


Figure 4. Permeation kinetics of indomethacin after introduction in the “Millicell” apical compartments of powders constituted by free γ -indomethacin, its co-crystals or the parent mixtures of γ -indomethacin with co-crystallizing agents. The permeations were analyzed across monolayers obtained by NCM460 cells. The permeation of free γ -indomethacin was analyzed across the Millicell filters alone (filter) or coated by monolayers. The cumulative amounts in the basolateral receiving compartments were linear within 60 min ($r \geq 0.97$, $P \leq 0.001$). The resulting slopes of the linear fits were used for the calculation of permeability coefficients (P_{app}). All data are reported as mean \pm SD of three independent experiments.

1
2
3 **Figure 5.** [A] Elimination profile of indomethacin after 0.90 mg infusion to rats. The elimination
4 followed an apparent first order kinetic, confirmed by the semilogarithmic plot reported in the inset (n
5 = 8, $r = 0.983$, $P < 0.0001$). The half-life of indomethacin was calculated to be 8.94 ± 0.38 hours.
6
7


8
9
10 [B] Blood indomethacin concentrations ($\mu\text{g/ml}$) after intravenous infusion (IV) or oral administration
11 of 0.90 mg dose to rats within 24 hours. The oral formulations were constituted by the sieved powders
12 of free γ -indomethacin, its co-crystals and the parent mixtures.
13
14

15
16 [C] Detailed blood indomethacin concentrations ($\mu\text{g/ml}$) after oral administration of 0.90 mg dose to
17 rats within 8 hours. All data reported in Figure are expressed as the mean \pm SD of four independent
18 experiments.
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60


**2-hydroxy-4-methyl-pyridine****Indomethacin****2-methoxy-5-nitroaniline****Saccharine**



a



b



c

