

Members of the DEPRESSD HADS Group:

Kira E. Riehm, Lady Davis Institute for Medical Research, Jewish General Hospital, Montréal, Québec, Canada;
Danielle B. Rice, Lady Davis Institute for Medical Research, Jewish General Hospital, Montréal, Québec, Canada;
Marleine Azar, Lady Davis Institute for Medical Research, Jewish General Hospital, Montréal, Québec, Canada;
Xin Wei Yan, Lady Davis Institute for Medical Research, Jewish General Hospital, Montréal, Québec, Canada;
Mahrukh Imran, Lady Davis Institute for Medical Research, Jewish General Hospital, Montréal, Québec, Canada;
Matthew J. Chiovitti, Lady Davis Institute for Medical Research, Jewish General Hospital, Montréal, Québec, Canada;
Nazanin Saadat, Lady Davis Institute for Medical Research, Jewish General Hospital, Montréal, Québec, Canada;
Jill T. Boruff, Schulich Library of Physical Sciences, Life Sciences, and Engineering, McGill University, Montréal, Québec, Canada ; Pim Cuijpers, Department of Clinical, Neuro and Developmental Psychology, Amsterdam Public Health Research Institute, Vrije Universiteit, Amsterdam, the Netherlands; Simon Gilbody, Hull York Medical School and the Department of Health Sciences, University of York, Heslington, York, United Kingdom; John P. A. Ioannidis, Department of Medicine, Department of Health Research and Policy, Department of Biomedical Data Science, Department of Statistics, Stanford University, Stanford, California, USA; Lorie A. Kloda, Library, Concordia University, Montréal, Québec, Canada; Scott B. Patten, Department of Community Health Sciences, University of Calgary, Calgary, Alberta, Canada; Roy C. Ziegelstein, Department of Medicine, Johns Hopkins University School of Medicine, Baltimore, Maryland, USA; Sarah Markham, Department of Biostatistics and Health Informatics, King's College London, London, UK; Melissa Henry, Lady Davis Institute for Medical Research, Jewish General Hospital, Montréal, Québec, Canada; Zahinoor Ismail, Hotchkiss Brain Institute and O'Brien Institute for Public Health, University of Calgary, Calgary, Alberta, Canada; Carmen G. Loiselle, Ingram School of Nursing, McGill University, Montréal, Québec, Canada; Nicholas D. Mitchell, Department of Psychiatry, University of Alberta, Edmonton, Alberta, Canada; Marcello Tonelli, Department of Medicine, University of Calgary, Calgary, Alberta, Canada; Samir Al-Adawi, Department of Behavioural Medicine, College of Medicine & Health Sciences, Sultan Qaboos University, Muscat, Oman; Kevin R. Beck, Department of Psychiatry, Singapore General Hospital, Singapore; Anna Beraldi, kbo Lech-Mangfall-Klinik für Psychiatrie, Psychotherapie und Psychosomatik, Garmisch-Partenkirchen, Bayern, German; Charles N. Bernstein, Department of Medicine, University of Manitoba, Winnipeg, Manitoba, Canada; Birgitte Boye, Psychosomatic and C-L psychiatry adult, Division of Psychiatry and Addiction, Oslo University Hospital, Oslo, Norway; Natalie Büel-Drabe, Department of

Psychiatry and Psychotherapy, University Hospital Zürich, Zürich Switzerland; Adomas Bunevicius, Neuroscience Institute, Lithuanian University of Health Sciences, Kaunas, Lithuania; Ceyhun Can, Adana City Training and Research Hospital, Adana, Turkey; Gregory Carter, School of Medicine and Public Health, University of Newcastle, Callaghan NSW, Australia; Chih-Ken Chen, Community Medicine Research Center, Keelung Chang Gung Memorial Hospital and Chang Gung University College of Medicine, Keelung, Taiwan; Gary Cheung, Department of Psychiatry, University of Auckland, Auckland, New Zealand; Kerrie Clover, Centre for Brain and Mental Health Research, University of Newcastle, Callaghan NSW, Australia; Gema Costa-Requena, Clinical Psychology, Department of Psychiatry, Hospital Universitari Vall d'Hebron, Universitat Autònoma de Barcelona, CIBERSAM, Barcelona, Spain; Daniel Cukor, Rogosin Institute, New York, New York, USA; Eli Dabscheck, The Alfred Hospital, Prahran, VIC, Australia; Federico M. Daray, Institute of Pharmacology, School of Medicine, University of Buenos Aires, Buenos Aires, Argentina; Jennifer De Souza, Birmingham and Solihull Mental Health Foundation Trust, Birmingham, UK; Marie Dorow, Institute of Social Medicine, Occupational Health and Public Health, University of Leipzig, Leipzig, Germany; Marina G. Downing, Monash Epworth Rehabilitation Research Centre, Epworth HealthCare, Melbourne VIC, Australia; Anthony Feinstein, Department of Psychiatry, University of Toronto, Toronto, Ontario, Canada; Panagiotis P. Ferentinos, 2nd Department of Psychiatry, Attikon General Hospital, National and Kapodistrian University of Athens, Athens, Greece; Felix H. Fischer, Department of Psychosomatic Medicine, Center for Internal Medicine and Dermatology, Charité - Universitätsmedizin Berlin, corporate member of Freie Universität Berlin, Humboldt-Universität zu Berlin, and Berlin Institute of Health, Berlin, Germany; Alastair J. Flint, University Health Network, Toronto, Ontario, Canada; Maiko Fujimori, Section of Psychological Science, Division of Health Care Research, Center for Public Health Sciences, National Cancer Center, Tokyo, Japan; Pamela Gallagher, School of Psychology, Dublin City University, Dublin, Ireland; Milena Gandy, The Department of Psychology, Macquarie University, Sydney, Australia; Luigi Grassi, Institute of Psychiatry, Department of Biomedical and Specialty Surgical Sciences, University of Ferrara, Ferrara, Italy; Martin Härter, Department of Medical Psychology, University of Hamburg, Hamburg, Germany; Asuncion Hernando, Department of Medicine, Universidad Europea de Madrid, Villaviciosa de Odon, Spain; Josef Jenewein, Clinic Zugersee, Center for Psychiatry and Psychotherapie, Oberwil-Zug, Switzerland; Nathalie Jetté, Department of Neurology, Icahn School of Medicine at Mount Sinai, New York, New York, USA; Miguel Julião, Equipa Comunitária de Suporte em Cuidados Paliativos de Sintra, Portugal; Monika Keller, Division of Psychooncology,

Department of General Internal Medicine and Psychosomatics, University Hospital Heidelberg, Germany; Sung-Wan Kim, Department of Psychiatry, Chonnam National University Medical School, Republic of Korea; Marie Kjærgaard, Endocrinology Research Group, Medical Clinic, University Hospital of North Norway, Norway; Sebastian Köhler, Department of Psychiatry and Neuropsychology, School for Mental Health and Neuroscience, Maastricht University, Maastricht, The Netherlands; Hans-Helmut König, Department of Health Economics and Health Services Research, University Medical Center Hamburg-Eppendorf, Hamburg, Germany; Lalit K.R. Krishna, Department of Palliative Medicine, National Cancer Centre, Singapore; Yu Lee, Department of Psychiatry, Kaohsiung Chang Gung Memorial Hospital, Chang Gung University College of Medicine, Kaohsiung, Taiwan; Margrit Löbner, Institute of Social Medicine, Occupational Health and Public Health, University of Leipzig, Leipzig, Germany; Wim L. Loosman, Onze Lieve vrouw Gasthuis, Amsterdam, The Netherlands; Anthony W. Love, Department of Psychology, Victoria University, Victoria, Australia; Bernd Löwe, Department of Psychosomatic Medicine and Psychotherapy, University Medical Center Hamburg-Eppendorf, Hamburg, Germany; Ulrik F. Malt, Department of Research and Education Division of Surgery and Clinical Neuroscience, University of Oslo, Oslo, Norway; Ruth Ann Marrie, Department of Medicine, Max Rady College of Medicine, Rady Faculty of Health Sciences, Winnipeg, Manitoba, Canada; Rocio Martin-Santos, Department of Psychiatry and Psychology, Hospital Clinic, IDIBAPS, CIBERSAM, Barcelona, Spain; Loreto Massardo, Centro de Biología Celular y Biomedicina, Facultad de Medicina y Ciencia, Universidad San Sebastián, Santiago, Chile; Yutaka Matsuoka, Division of Health Care Research, Center for Public Health Sciences, National Cancer Center, Tokyo, Japan; Anja Mehnert, Department of Medical Psychology and Medical Sociology, University of Leipzig, Germany; Ioannis Michopoulos, 2nd Department of Psychiatry, Attikon General Hospital, National and Kapodistrian University of Athens, Athens, Greece; Laurent Misery, Department of Dermatology, University Hospital of Brest, Brest, France; Ricard Navines, Department of Psychiatry and Psychology, Hospital Clinic, IDIBAPS, CIBERSAM, Barcelona, Spain; Christian J. Nelson, Department of Psychiatry and Behavioral Sciences, Memorial Sloan Kettering Cancer Center, New York, New York, USA; Chong Guan Ng, Department of Psychological Medicine, Faculty of Medicine, University of Malaya, Kuala Lumpur, Malaysia; Meaghan L. O'Donnell, Phoenix Australia, Carlton VIC, Australia; Suzanne J. O'Rourke, School of Health in Social Sciences, University of Edinburgh, Edinburgh, Scotland; Ahmet Öztürk, Sabahattin Zaim University, Istanbul, Turkey; Alexander Pabst, Institute of Social Medicine, Occupational Health and Public Health (ISAP), Medical Faculty, University of Leipzig, Leipzig, Germany; Julie A. Pasco, Deakin

University, IMPACT – the Institute for Mental and Physical Health and Clinical Translation, School of Medicine, Geelong, Victoria, Australia; Jurate Peceliuniene, Vilnius University, Faculty of Medicine, Clinic of Internal Diseases, Family Medicine and Oncology, Vilnius, Lithuania; Luis Pintor, Instituto de Investigaciones Biomédicas Augusto Pi i Sunyer (IDIBAPS), Barcelona, Spain; Jennie L. Ponsford, School of Psychological Sciences, Monash University, Melbourne VIC, Australia; Federico Pulido, HIV Unit, Hospital 12 de Octubre, imas12, UCM, Madrid, Spain; Terence J. Quinn, Institute of Cardiovascular and Medical Sciences, University of Glasgow, Glasgow, UK; Silje E. Reme, Department of Psychology, Faculty of Social Sciences, University of Oslo, Oslo, Norway; Katrin Reuter, Private Practice for Psychotherapy and Psycho-oncology, Freiburg, Germany; Nina Rieckmann, Institute of Public Health, Charité - Universitätsmedizin Berlin, corporate member of the Freie Universität Berlin, Humboldt-Universität zu Berlin, and the Berlin Institute of Health, Berlin, Germany; Steffi G. Riedel-Heller, Institute of Social Medicine, Occupational Health and Public Health (ISAP), Medical Faculty, University of Leipzig, Leipzig, Germany; Alasdair G. Rooney, Division of Psychiatry, University of Edinburgh, Edinburgh, UK; Roberto Sánchez-González, Department of Psychiatry, Institut de Neuropsiquiatria i Addiccions, Centre Emili Mira, Parc de Salut Mar, Barcelona, Spain; Rebecca M. Saracino, Department of Psychiatry and Behavioral Sciences, Memorial Sloan Kettering Cancer Center, New York, USA; Melanie P. J. Schellekens, Scientific Research Department, Helen Dowling Institute, Bilthoven, The Netherlands; Martin Scherer, Institute of Primary Medical Care, University Medical Center Hamburg-Eppendorf, Hamburg, Germany; Marcelo L. Schwarzbald, Department of Internal Medicine, Federal University of Santa Catarina, Florianópolis, Santa Catarina, Brazil; Vesile Senturk Cankorur, Department of Psychiatry, Faculty of Medicine, Ankara University, Ankara, Turkey; Juwita Shaaban, Department of Family Medicine, School of Medical Sciences, Universiti Sains Malaysia, Kelantan, Malaysia; Louise Sharpe, School of Psychology, The University of Sydney, Sydney NSW, Australia; Michael Sharpe, Department of Psychological Medicine, University of Oxford, Oxford, UK; Sébastien Simard, Département des sciences de la santé, Université du Québec à Chicoutimi (UQAC), Québec, Canada; Susanne Singer, University Medical Centre Mainz, Institute of Medical Biostatistics, Epidemiology and Informatics, Mainz, Germany; Lesley Stafford, Centre for Women's Mental Health, Royal Women's Hospital, Parkville, Australia; Jon Stone, Centre for Clinical Brain Sciences, University of Edinburgh, Edinburgh, UK; Natalie A. Strobel, Kurongkurl Katitjin, Edith Cowan University, Perth, Western Australia, Australia; Serge Sultan, Département de Psychologie, Faculté des arts et des sciences, Université de Montréal, Québec, Canada; Antonio L. Teixeira, University of Texas Health Science Center

at Houston, Houston, Texas, USA; Istvan Tiringier, Pécs University, Medical School, Institute of Behavioral Sciences, Pécs, Hungary; Mira Tschorn, Department of Psychiatry and Psychotherapy, CCM, Charité – Universitätsmedizin Berlin, corporate member of Freie Universität Berlin, Humboldt-Universität zu Berlin, and Berlin Institute of Health, Berlin, Germany; Ka-Yee Tung, Kwai Chung Hospital, Hong Kong SAR, China; Alyna Turner, Faculty of Health and Medicine, School of Medicine and Public Health, The University of Newcastle, Callaghan, NSW, Australia; Michael Wagner, Department of Neurodegenerative Diseases and Geriatric Psychiatry, University Hospital Bonn, Bonn, Germany; Jane Walker, Department of Psychiatry, University of Oxford, Oxford, UK; Mark Walterfang, Neuropsychiatry Unit, Royal Melbourne Hospital, Melbourne, Australia; Liang-Jen Wang, Department of Child and Adolescent Psychiatry, Kaohsiung Chang Gung Memorial Hospital and Chang Gung University College of Medicine, Kaohsiung, Taiwan; Siegfried B. Weyerer, Central Institute of Mental Health, Medical Faculty Mannheim/Heidelberg University, Mannheim, Germany; Jennifer White, Department of Physiotherapy, School of Primary and Allied Health Care, Monash University, Melbourne, Australia; Lana J. Williams, Deakin University, IMPACT – the Institute for Mental and Physical Health and Clinical Translation, School of Medicine, Geelong, Victoria, Australia; Lai-Yi Wong, Kwai Chung Hospital, Hong Kong SAR, China.