

An existence result for a constrained two-phase transition model with metastable phase for vehicular traffic

Mohamed Benyahia, Carlotta Donadello, Nikodem Dymski, Massimiliano D. Rosini

July 24, 2018

Abstract

In this paper we study a phase transition model for vehicular traffic flows. Two phases are taken into account, according to whether the traffic is light or heavy. We assume that the two phases have a non-empty intersection, the so called *metastable* phase. The model is given by the Lighthill-Whitham-Richards model in the *free-flow* phase and by the Aw-Rascle-Zhang model in the *congested* phase. In particular, we study the existence of solutions to Cauchy problems satisfying a local point constraint on the density flux. We prove that if the constraint F is higher than the minimal flux f_c^- of the metastable phase, then constrained Cauchy problems with initial data of bounded total variation admit globally defined solutions. We also provide sufficient conditions on the initial data that guarantee the global existence of solutions also in the case $F < f_c^-$. These results are obtained by applying the wave-front tracking technique.

2010 Mathematical Subject Classification. Primary: 35L65, 90B20, 35L45

Key words. Conservation laws, phase transitions, Lighthill-Whitham-Richards model, Aw-Rascle-Zhang model, point constraint on the density flux, Cauchy problem, wave-front tracking.

1 Introduction

In this paper we study one of the constrained phase transition models of hyperbolic conservation laws introduced in [12]. The application of such model is, for instance, the modelling of vehicular traffic along a road with pointlike inhomogeneities characterized by limited capacity, such as speed bumps, construction sites, tollbooths, etc.

The model considers two different phases corresponding to the *congested phase* Ω_c and the *free-flow phase* Ω_f . The model is given by a 2×2 system of conservation laws in the congested phase, coupled with a scalar conservation law in the free-flow phase. The coupling is achieved via *phase transitions*, namely discontinuities between two states belonging to different phases and satisfying the Rankine-Hugoniot conditions.

The first two-phase model has been proposed by Colombo in [9]. The motivation stems from experimental data, according to which the density flux represented in the fundamental diagram is one-dimensional for high velocities, while it covers a two-dimensional domain for low velocities, see [9, Figure 1.1]. For this reason, it is reasonable to describe the dynamics in the congested regime with a 2×2 system of conservation laws and those in the free regime with a scalar conservation law.

Later, Goatin proposed in [16] a two-phase model obtained by coupling the ARZ model by Aw, Rascle and Zhang [4, 21] for the congested phase Ω_c , with the LWR model by Lighthill, Whitham and Richards [18, 19] for the free-flow phase Ω_f . We recall that this model has been recently generalized in [5].

Both the models introduced in [9] and [16] assume that $\Omega_c \cap \Omega_f = \emptyset$. The first two-phase model that considers a metastable phase $\Omega_c \cap \Omega_f \neq \emptyset$ has been introduced in [7]. We also recall that, differently from [5, 16], for the models in [7, 9] the density flux function vanishes at a maximal density, whose inverse corresponds to the average length of the vehicles. Here we consider the case $\Omega_c \cap \Omega_f \neq \emptyset$. For this reason, in order to ensure the well-posedness of the Cauchy problems, see [9, Remark 2], we also assume that Ω_f is characterized by a unique value of the velocity, V . At last, we consider an heterogeneous traffic with vehicles having different lengths and allow the density flux function to vanish at different densities.

These two-phase models have been recently generalized in [6, 12] by considering Riemann problems, namely Cauchy problems for piecewise constant initial data with a single jump, coupled with a constraint on the density flux, so that at the interface $x = 0$ the density flux of the solution must be lower than a given constant quantity F . This condition is referred to as *unilateral point constraint* and can be thought of as a pointwise bottleneck at $x = 0$ that hinders the density flow, see [20] and the references therein. In vehicular traffic, a point constraint accounts for inhomogeneities of the road and models, for instance, the presence of a toll gate across which the flow of the vehicles cannot exceed its capacity F .

In the case in which no constraint conditions are enforced, existence results for the Cauchy problems for the above mentioned two-phase transition models have already been established, see [5, 7, 11, 16]. In the present paper, we focus on the constrained version proposed in [12] for the model introduced in [5] and prove an existence result for constrained Cauchy problems. More precisely, we use the Riemann solvers established in [5] and [12] in a wave-front tracking scheme and prove that the obtained approximate solution u_n converges (up to a subsequence) to a globally defined solution of the constrained Cauchy problem with general **BV**-initial data, at least in the case $F \geq f_c^-$, where the threshold value f_c^- is the minimal density flux of the metastable phase, see Figure 1. At last, in the case $F < f_c^-$ we give sufficient conditions on the initial data that ensure the convergence of u_n to a globally defined solution of the constrained Cauchy problem.

The paper is organized as follows. In the next section we introduce the notations used throughout the paper, the model, the definitions of solutions to the unconstrained and constrained Cauchy problems, the main result in Theorem 2.2 and at last the Riemann solvers for the unconstrained and constrained Riemann problems. In Section 3 we apply the model to reproduce the traffic across a toll gate. Finally, in the last section we defer the technical proofs.

2 Notations, definitions and main result

In this section we state the main assumptions on the parameters, collect useful notations, see Figure 1, give the definition of solutions, state the main result in Theorem 2.2 and at last introduce the Riemann solvers.




Figure 1: Notations.

2.1 Notations

Denote by $\rho \geq 0$ and $v \geq 0$ the density and the velocity of the vehicles, respectively. Let $u \doteq (\rho, v)$ and $f(u) \doteq v\rho$ be the density flux. If $V > 0$ is the unique velocity in the free-flow phase Ω_f and ρ^+ is the maximal density in Ω_f , then

$$\Omega_f \doteq \{u \in \mathbb{R}_+^2 : \rho \leq \rho^+, v = V\},$$

where $\mathbb{R}_+ \doteq [0, \infty)$. If the velocity V is reached in the congested phase Ω_c for densities ranging in $[\rho^-, \rho^+] \subset (0, \infty)$, then

$$\Omega_c \doteq \{u \in \mathbb{R}_+^2 : v \leq V, w^- \leq v + p(\rho) \leq w^+\},$$

where $w^\pm \doteq p(\rho^\pm) + V$. Above $p \in \mathbf{C}^2((0, \infty); \mathbb{R})$ is an anticipation factor, which takes into account drivers' reactions to the state of traffic in front of them. We assume that

$$p(0) = 0, \quad p'(\rho) > 0, \quad 2p'(\rho) + p''(\rho)\rho > 0 \quad \text{for every } \rho > 0. \quad (1)$$

Typical choices for p are $p(\rho) \doteq \rho^\gamma$ with $\gamma > 0$, see [4], and $p(\rho) \doteq V_{\text{ref}} \ln(\rho/\rho_{\text{max}})$ with $V_{\text{ref}} > 0$ and $\rho_{\text{max}} > 0$, see [16].

Let $f_c^\pm \doteq V\rho^\pm$ and $R \doteq p^{-1}(w^+) > 0$ be the maximal density (in the congested phase). Let

$$\Omega \doteq \Omega_f \cup \Omega_c, \quad \Omega_f^- \doteq \{u \in \Omega_f : \rho \in [0, \rho^-]\}, \quad \Omega_f^+ \doteq \{u \in \Omega_f : \rho \in [\rho^-, \rho^+]\}, \quad \Omega_c^- \doteq \Omega_c \setminus \Omega_f^+.$$

Notice that $\Omega_f \cap \Omega_c = \Omega_f^+$. We assume that

$$v < p'(\rho)\rho \text{ for every } (\rho, v) \in \Omega_c. \quad (2)$$

The (extended) Lagrangian marker $\mathbf{w}: \Omega \rightarrow [w^- - 1, w^+]$ is defined by

$$\mathbf{w}(u) \doteq \begin{cases} v + p(\rho) & \text{if } u \in \Omega_c, \\ w^- - 1 + \frac{\rho}{\rho^-} & \text{if } u \in \Omega_f^-. \end{cases}$$

Let $\mathbf{W}: \Omega \rightarrow [w^-, w^+]$ be defined by

$$\mathbf{W}(u) \doteq \max\{w^-, \mathbf{w}(u)\}.$$

The 2×2 system of conservation laws describing the traffic in the congested phase has two characteristic families of Lax curves. In the (ρ, f) -plane the Lax curves in Ω_c of the first and second characteristic families passing through $\bar{u} = (\bar{\rho}, \bar{v}) \in \Omega_c$ are respectively described by the graphs of the maps

$$\begin{aligned} [p^{-1}(\mathbf{w}(\bar{u}) - V), p^{-1}(\mathbf{w}(\bar{u}))] \ni \rho \mapsto \mathfrak{L}_{\bar{u}}(\rho) &\doteq f(\rho, \mathbf{w}(\bar{u}) - p(\rho)), \\ [p^{-1}(w^- - \bar{v}), p^{-1}(w^+ - \bar{v})] \ni \rho \mapsto \bar{v} \rho. \end{aligned}$$

Remark 2.1. Conditions (1) and (2) ensure that for any $w \in [w^-, w^+]$ the map $\rho \mapsto \mathfrak{L}_w(\rho) = (w - p(\rho)) \rho$ is strictly concave and strictly decreasing in $[p^{-1}(w - V), p^{-1}(w)]$. Indeed, for any $w \in [w^-, w^+]$ and $\rho \in [p^{-1}(w - V), p^{-1}(w)]$, we have that $(\rho, w - p(\rho)) \in \Omega_c$ and therefore

$$\mathfrak{L}'_w(\rho) = w - p(\rho) - p'(\rho) \rho < 0, \quad \mathfrak{L}''_w(\rho) = -2p(\rho) - p''(\rho) \rho < 0.$$

If for instance $p(\rho) \doteq V_{\text{ref}} \ln(\rho/\rho_{\text{max}})$ with $V_{\text{ref}} > 0$ and $\rho_{\text{max}} > 0$, then $p'(\rho) \rho = V_{\text{ref}}$ and (2) is equivalent to require $V < V_{\text{ref}}$, while (1) is trivial. If for instance $p(\rho) \doteq \rho^\gamma$ with $\gamma > 0$, then $\Omega_c = \{u \in \mathbb{R}_+^2 : v \leq V, w^- - v \leq \rho^\gamma \leq w^+ - v\}$ and therefore

$$\min_{u \in \Omega_c} (p'(\rho) \rho - v) = \min_{u \in \Omega_c} (\gamma \rho^\gamma - v) = \gamma w^- - (\gamma + 1)V,$$

hence (2) is equivalent to require $(\gamma + 1)V < \gamma w^-$, while (1) is trivial.

We introduce the following functions, see Figure 1:

$$\begin{aligned} \omega: \Omega_c &\rightarrow \Omega_f^+, & u = \omega(\bar{u}) &\iff \begin{cases} \mathbf{w}(u) = \mathbf{w}(\bar{u}), \\ v = V, \end{cases} \\ \mathbf{v}^\pm: \Omega &\rightarrow \Omega_c, & u^\pm = \mathbf{v}^\pm(\bar{u}) &\iff \begin{cases} \mathbf{w}(u^\pm) = w^\pm, \\ v^\pm = \bar{v}, \end{cases} \\ \mathbf{u}_*: \Omega^2 &\rightarrow \Omega_c, & u_* = \mathbf{u}_*(u_\ell, u_r) &\iff \begin{cases} \mathbf{w}(u_*) = \mathbf{W}(u_\ell), \\ v_* = v_r, \end{cases} \\ \Lambda: \{(u_\ell, u_r) \in \Omega^2 : \rho_\ell \neq \rho_r\} &\rightarrow \mathbb{R}, & \Lambda(u_\ell, u_r) &\doteq \frac{f(u_r) - f(u_\ell)}{\rho_r - \rho_\ell}. \end{aligned}$$

Notice that:

- the point $\omega(\bar{u})$ is the intersection of the Lax curve of the first characteristic family passing through \bar{u} and Ω_f^+ , namely the Lax curve of the second characteristic family passing through $(0, V)$;
- for any $w \in [w^-, w^+]$ the point $(p^{-1}(w), 0)$ is the intersection of the Lax curve of the first characteristic family corresponding to w and the segment $\{(\rho, v) \in \Omega_c : v = 0\}$, namely the Lax curve of the second characteristic family passing through $(p^{-1}(w^\pm), 0)$;
- the point $\mathbf{v}^\pm(\bar{u})$ is the intersection of the Lax curve of the second characteristic family passing through \bar{u} and $\{u \in \Omega_c : \mathbf{w}(u) = w^\pm\}$, namely the Lax curve of the first characteristic family passing through $(p^{-1}(w^\pm), 0)$;
- for any $u_\ell, u_r \in \Omega_c$ the point $\mathbf{u}_*(u_\ell, u_r)$ is the intersection between the Lax curve of the first characteristic family passing through u_ℓ and the Lax curve of the second characteristic family passing through u_r ;
- $\Lambda(u_\ell, u_r)$ is the speed of a discontinuity (u_ℓ, u_r) , that in the (ρ, f) -coordinates coincides with the slope of the segment connecting u_ℓ and u_r .

Observe that by definition $\mathbf{v}^\pm(\bar{u}) = \mathbf{u}_*((p^{-1}(w^\pm), 0), \bar{u})$ and $\omega(\bar{u}) = \mathbf{u}_*(\bar{u}, (0, V))$.

We denote by \mathcal{R} and \mathcal{R}_F the Riemann solver and the constrained Riemann solver introduced in [5] and [12], respectively, see Section 2.3 for more details.

2.2 The constrained Cauchy problem

We study the constrained Cauchy problem for the phase transition model

$$\begin{array}{ll}
 \text{Free-flow} & \text{Congested flow} \\
 \left\{ \begin{array}{l} u \in \Omega_f, \\ \rho_t + (\rho V)_x = 0, \\ v = V, \end{array} \right. & \left\{ \begin{array}{l} u \in \Omega_c, \\ \rho_t + (\rho v)_x = 0, \\ (\rho \mathbf{w}(u))_t + (\rho \mathbf{w}(u) v)_x = 0, \end{array} \right. \quad (3)
 \end{array}$$

with initial datum

$$u(0, x) = u^o(x) \quad (4)$$

and local point constraint on the density flux at $x = 0$

$$f(u(t, 0_{\pm})) \leq F, \quad (5)$$

where $F \in [0, f_c^+]$ is a given constant quantity. To this aim we apply the wave-front tracking algorithm, which is based on the definition of the Riemann solvers defined in the next sections.

Introduce, see Figure 2, $v_F^{\pm} \in [0, V]$ and $w_F \in [w^- - 1, w^+]$ defined by the following conditions:

$$\begin{array}{llll}
 \text{if } F = f_c^+ : & v_F^+ \doteq V, & v_F^- \doteq V, & w_F \doteq w^+, \\
 \text{if } F \in [f_c^-, f_c^+) : & v_F^+ \doteq V, & v_F^- + p(F/v_F^-) = w^+, & w_F \doteq p(F/V) + V, \\
 \text{if } F \in (0, f_c^-) : & v_F^+ + p(F/v_F^+) = w^-, & v_F^- + p(F/v_F^-) = w^+, & w_F \doteq w^- - 1 + \frac{F}{f_c^-}, \\
 \text{if } F = 0 : & v_F^+ \doteq 0, & v_F^- \doteq 0, & w_F \doteq w^- - 1.
 \end{array}$$




Figure 2: Geometrical meaning of w_F , v_F^{\pm} and Ξ_F in the case $F \in (0, f_c^-)$. The curve in the figure on the left is the graph of Ξ_F , which corresponds to the horizontal solid segment in the figure on the right.

For any $F \in (0, f_c^-)$, let $\Xi_F : [v_F^-, v_F^+] \rightarrow [w^-, w^+]$ be given by $\Xi_F(v) \doteq v + p(F/v)$, see Figure 2. Notice that Ξ_F is strictly decreasing because by (2)

$$\left(\frac{F}{v}, v\right) \in \Omega_c \quad \Rightarrow \quad \Xi_F'(v) = 1 - p' \left(\frac{F}{v}\right) \frac{F}{v^2} < 0,$$

moreover it is strictly convex because by (1)

$$\Xi_F''(v) = \left[2p' \left(\frac{F}{v}\right) + p'' \left(\frac{F}{v}\right) \frac{F}{v} \right] \frac{F}{v^3} > 0.$$

The notion of solution to Cauchy problem (3), (4) necessarily involves both the notions of solution to the Cauchy problems for LWR and ARZ models, that have to be combined by defining which phase transitions are admissible, see [5]. Since the characteristic field corresponding to the free phase is linearly degenerate, a

discontinuity between states in Ω_f is entropic if and only if it satisfies the corresponding Rankine-Hugoniot condition, namely its speed of propagation is V . For this reason we consider only the entropy-entropy flux pair

$$\mathbf{E}^k(u) \doteq \begin{cases} 0 & \text{if } v \geq k, \\ \frac{\rho}{p^{-1}(\mathbb{W}(u) - k)} - 1 & \text{if } v < k, \end{cases} \quad \mathbf{Q}^k(u) \doteq \begin{cases} 0 & \text{if } v \geq k, \\ \frac{f(u)}{p^{-1}(\mathbb{W}(u) - k)} - k & \text{if } v < k, \end{cases}$$

for $u \in \Omega$ and $k \in [0, V]$, which is obtained by adapting the entropy-entropy flux pair introduced in [1] for the ARZ model.

Definition 2.1. *Let $u^\circ \in \mathbf{BV}(\mathbb{R}; \Omega)$. We say that $u \in \mathbf{L}^\infty((0, \infty); \mathbf{BV}(\mathbb{R}; \Omega)) \cap \mathbf{C}^0(\mathbb{R}_+; \mathbf{L}_{\text{loc}}^1(\mathbb{R}; \Omega))$ is a solution to Cauchy problem (3), (4) if the following holds:*

(S.1) *Condition (4) holds for a.e. $x \in \mathbb{R}$, namely*

$$u(0, x) = u^\circ(x) \quad \text{for a.e. } x \in \mathbb{R}.$$

(S.2) *For any $\phi \in \mathbf{C}_c^\infty((0, \infty) \times \mathbb{R}; \mathbb{R})$ we have*

$$\int_0^\infty \int_{\mathbb{R}} (\rho \phi_t + f(u) \phi_x) \left(\frac{1}{\mathbb{W}(u)} \right) dx dt = \begin{pmatrix} 0 \\ 0 \end{pmatrix}.$$

(S.3) *For any $k \in [0, V]$ and $\phi \in \mathbf{C}_c^\infty((0, \infty) \times \mathbb{R}; \mathbb{R})$ such that $\phi \geq 0$ we have*

$$\int_0^\infty \int_{\mathbb{R}} (\mathbf{E}^k(u) \phi_t + \mathbf{Q}^k(u) \phi_x) dx dt \geq 0.$$

We recall the existence result proved in [5, Theorem 2.8].

Theorem 2.1. *Cauchy problem (3), (4) with initial datum $u^\circ \in \mathbf{L}^1 \cap \mathbf{BV}(\mathbb{R}; \Omega)$ admits a solution u in the sense of Definition 2.1; moreover there exist two constants C° and L° such that for any $t, s \geq 0$*

$$\text{TV}(u(t)) \leq \text{TV}(u^\circ), \quad \|u(t)\|_{\mathbf{L}^\infty(\mathbb{R}; \Omega)} \leq C^\circ, \quad \|u(t) - u(s)\|_{\mathbf{L}^1(\mathbb{R}; \Omega)} \leq L^\circ |t - s|.$$

In the following definition we introduce the notion of solution to constrained Cauchy problem (3), (4), (5), which is obtained by adapting that introduced in Definition 2.1 for Cauchy problem (3), (4).

Definition 2.2. *Let $u^\circ \in \mathbf{BV}(\mathbb{R}; \Omega)$. We say that $u \in \mathbf{L}^\infty((0, \infty); \mathbf{BV}(\mathbb{R}; \Omega)) \cap \mathbf{C}^0(\mathbb{R}_+; \mathbf{L}_{\text{loc}}^1(\mathbb{R}; \Omega))$ is a solution to constrained Cauchy problem (3), (4), (5) if the following holds:*

(CS.1) *Condition (4) holds for a.e. $x \in \mathbb{R}$, namely*

$$u(0, x) = u^\circ(x) \quad \text{for a.e. } x \in \mathbb{R}.$$

(CS.2) *For any $\phi \in \mathbf{C}_c^\infty((0, \infty) \times \mathbb{R}; \mathbb{R})$ we have*

$$\int_0^\infty \int_{\mathbb{R}} (\rho \phi_t + f(u) \phi_x) dx dt = 0 \tag{6}$$

and if $\phi(\cdot, 0) \equiv 0$ then

$$\int_0^\infty \int_{\mathbb{R}} (\rho \phi_t + f(u) \phi_x) \mathbb{W}(u) dx dt = 0. \tag{7}$$

(CS.3) *For any $k \in [0, V]$ and $\phi \in \mathbf{C}_c^\infty((0, \infty) \times \mathbb{R}; \mathbb{R})$ such that $\phi(\cdot, 0) \equiv 0$ and $\phi \geq 0$ we have*

$$\int_0^\infty \int_{\mathbb{R}} (\mathbf{E}^k(u) \phi_t + \mathbf{Q}^k(u) \phi_x) dx dt \geq 0. \tag{8}$$

(CS.4) *Condition (5) holds for a.e. $t > 0$, namely*

$$f(u(t, 0_\pm)) \leq F \quad \text{for a.e. } t > 0.$$

In the following proposition we state which discontinuities are admissible for the solutions to (3), (4), (5).

Proposition 2.1. *Let u be a solution of constrained Cauchy problem (3), (4), (5) in the sense of Definition 2.2. Then u has the following properties:*

- Any discontinuity $\delta(t)$ of $x \mapsto u(t, x)$ satisfies the first Rankine-Hugoniot jump condition

$$\left[\rho(t, \delta(t)_+) - \rho(t, \delta(t)_-) \right] \dot{\delta}(t) = f(u(t, \delta(t)_+)) - f(u(t, \delta(t)_-)), \quad (9)$$

and if $\delta(t) \neq 0$, then it satisfies also the second Rankine-Hugoniot jump condition

$$\begin{aligned} & \left[\rho(t, \delta(t)_+) \mathbb{W}(u(t, \delta(t)_+)) - \rho(t, \delta(t)_-) \mathbb{W}(u(t, \delta(t)_-)) \right] \dot{\delta}(t) \\ &= f(u(t, \delta(t)_+)) \mathbb{W}(u(t, \delta(t)_+)) - f(u(t, \delta(t)_-)) \mathbb{W}(u(t, \delta(t)_-)). \end{aligned} \quad (10)$$

- Any discontinuity of u away from the constraint is classical, i.e. it satisfies the Lax entropy inequalities.
- Non-classical discontinuities of u may occur only at the constraint location $x = 0$, and in this case the (density) flux at $x = 0$ does not exceed the maximal flux F allowed by the constraint.

Proof. These properties follow directly from (CS.2), (CS.3) and (CS.4). Let us just underline that (9), (10) are equivalent to

$$\begin{aligned} & \left[v(t, 0_+) - \dot{\delta}(t) \right] \rho(t, 0_+) = \left[v(t, 0_-) - \dot{\delta}(t) \right] \rho(t, 0_-), \\ & \left[\mathbb{W}(u(t, 0_+)) - \mathbb{W}(u(t, 0_-)) \right] \left[v(t, 0_-) - \dot{\delta}(t) \right] \rho(t, 0_-) = 0. \end{aligned}$$

In particular phase transitions and shocks are admissible because for them $\mathbb{W}(u(t, 0_+)) = \mathbb{W}(u(t, 0_-))$, while the contact discontinuities are admissible because for them $\dot{\delta}(t) = v(t, 0_\pm)$. \square

Remark 2.2. *Differently from any solution to Cauchy problem (3), (4), a solution u to constrained Cauchy problem (3), (4), (5) does not satisfy in general the second Rankine-Hugoniot condition (10) along $x = 0$*

$$\rho(t, 0_-) \mathbb{W}(u(t, 0_-)) v(t, 0_-) = \rho(t, 0_+) \mathbb{W}(u(t, 0_+)) v(t, 0_+) \quad \text{for a.e. } t > 0.$$

Indeed the (extended) linearized momentum $\rho \mathbb{W}(u)$ is conserved across (classical) shocks and phase transitions, but in general it is not conserved across non-classical shocks even if they are between states in Ω_c . As a consequence, a solution to (3), (4), (5) taking values in Ω_c is not necessarily a weak solution to the 2×2 system of conservation laws in (3) for the congested flow. For this reason in (7) (and then also in (8)) we consider test functions ϕ such that $\phi(\cdot, 0) \equiv 0$.

This is in the same spirit of the solutions considered in [6, 12–15] for traffic through locations with reduced capacity. However, with this choice for the test functions in (7) and (8) we loose the possibility to better characterize the (density) flux at $x = 0$ associated to non-classical shocks. In fact, differently from what is proved in [10] for the LWR model and in [3] for the ARZ model, we cannot ensure that the flux of the non-classical shocks of any solution is equal to the maximal flux F allowed by the constraint. Nevertheless, in Section 4.4 we can give sufficient conditions ensuring that the solutions constructed with our wave-front tracking algorithm have this property, see Proposition 4.4.

Let $[w^- - 1, w^+] \ni w \mapsto \hat{u}(w, F) = (\hat{r}(w, F), \hat{v}(w, F)) \in \Omega_c$ and $[0, V] \ni v \mapsto \check{u}(v, F) = (\check{r}(v, F), \check{v}(v, F)) \in \Omega$ be defined in the (v, w) -coordinates by, see Figures 3 and 4,

$$\hat{v}(w, F) \doteq \begin{cases} \Xi_F^{-1}(w) & \text{if } w > \max\{w^-, w_F\}, \\ v_F^+ & \text{if } w_F < w \leq w^-, \\ V & \text{if } w \leq w_F, \end{cases} \quad \hat{w}(w, F) \doteq \begin{cases} w & \text{if } w > \max\{w^-, w_F\}, \\ w^- & \text{if } w_F < w \leq w^-, \\ w_F & \text{if } w \leq w_F, \end{cases} \quad (11a)$$

$$\check{v}(v, F) \doteq \begin{cases} V & \text{if } v > v_F^+, \\ v & \text{if } v \in [v_F^-, v_F^+], \\ v_F^- & \text{if } v < v_F^-, \end{cases} \quad \check{w}(v, F) \doteq \begin{cases} w_F & \text{if } v > v_F^+, \\ \Xi_F(v) & \text{if } v \in [v_F^-, v_F^+], \\ w^+ & \text{if } v < v_F^-, \end{cases} \quad (11b)$$

where $\hat{w} \doteq \mathbf{w} \circ \hat{u}$ and $\check{w} \doteq \mathbf{w} \circ \check{u}$.




Figure 3: Geometrical meaning of \hat{u} and \tilde{u} defined in (11) in the case $F \in (f_c^-, f_c^+)$.

Remark 2.3. Notice that

$$f(\hat{u}(w, F)) = f(\tilde{u}(v, F)) = F.$$

Moreover, $w \mapsto \hat{u}(w, F)$ and $v \mapsto \tilde{u}(v, F)$ are continuous if and only if $F \geq f_c^-$, and in this case they are Lipschitz continuous. On the other hand, if $F < f_c^-$, then $w \mapsto \hat{u}(w, F)$ and $v \mapsto \tilde{u}(v, F)$ are only left-continuous. Moreover $\hat{w}(w, F) \geq w$ and $\check{v}(v, F) \geq v$. At last, $w \mapsto \hat{w}(w, F)$ and $v \mapsto \check{v}(v, F)$ are non-decreasing, while $w \mapsto \hat{v}(w, F)$ and $v \mapsto \check{w}(v, F)$ are non-increasing.

Denote by TV_+ and TV_- the positive and negative total variations, respectively. For any $u: \mathbb{R} \rightarrow \Omega$ let

$$\hat{\Upsilon}(u) \doteq \text{TV}_+(\hat{v}(\mathbf{w}(u), F); (-\infty, 0)) + \text{TV}_-(\hat{w}(\mathbf{w}(u), F); (-\infty, 0)), \quad (12a)$$

$$\check{\Upsilon}(u) \doteq \text{TV}_+(\check{v}(v, F); (0, \infty)) + \text{TV}_-(\check{w}(v, F); (0, \infty)). \quad (12b)$$

For any $u \in \Omega$ and $k \in [0, V]$ let

$$\mathbf{N}_F^k(u) \doteq \begin{cases} f(u) \left[\frac{k}{F} - \frac{1}{p^{-1}(\mathbf{W}(u) - k)} \right]_+ & \text{if } F \neq 0, \\ k & \text{if } F = 0, \end{cases} \quad [w]_+ \doteq \begin{cases} w & \text{if } w > 0, \\ 0 & \text{otherwise.} \end{cases}$$

We are now in the position to state the main result of the paper.

Theorem 2.2. Let $u^\circ \in \mathbf{L}^1 \cap \mathbf{BV}(\mathbb{R}; \Omega)$ and $F \in [0, f_c^+]$ satisfy one of the following conditions:

(H.1) $F \in [f_c^-, f_c^+]$;

(H.2) $F \in [0, f_c^-)$ and $\hat{\Upsilon}(u^\circ) + \check{\Upsilon}(u^\circ)$ is bounded.

Then the approximate solutions u_n constructed in Section 4.1 converge to a solution $u \in \mathbf{C}^0(\mathbb{R}_+; \mathbf{BV}(\mathbb{R}; \Omega))$ of constrained Cauchy problem (3), (4), (5) in the sense of Definition 2.2. Moreover for all $t, s \in \mathbb{R}_+$ the following estimates hold

$$\text{TV}(u(t)) \leq C_F^\circ, \quad \|u(t) - u(s)\|_{\mathbf{L}^1(\mathbb{R}; \Omega)} \leq L_F^\circ |t - s|, \quad \|u(t)\|_{\mathbf{L}^\infty(\mathbb{R}; \Omega)} \leq R + V, \quad (13)$$




Figure 4: Geometrical meaning of \hat{u} and \tilde{u} defined in (11) in the case $F \in (0, f_c^-)$.

where C_F^o and L_F^o are constants that depend on u^o and F . Furthermore, non-classical discontinuities of u can occur only at the constraint location $x = 0$, and if for any $k \in [0, V]$ and $\phi \in \mathbf{C}_c^\infty((0, \infty) \times \mathbb{R}; \mathbb{R})$ such that $\phi \geq 0$ we have

$$\lim_{n \rightarrow \infty} \int_0^T \mathbf{N}_F^k(u_n(t, 0_-)) \phi(t, 0) dt = \int_0^T \mathbf{N}_F^k(u(t, 0_-)) \phi(t, 0) dt, \quad (14)$$

then the (density) flow at $x = 0$ is the maximal flow F allowed by the constraint.

As in [2, 5, 11], the proof of the above theorem is based on the wave-front tracking algorithm, see [8, 17] and the references therein. The details of the proof are deferred to Section 4.

Remark 2.4. If $F \in [f_c^-, f_c^+]$, then $w \mapsto \hat{u}(w, F)$ and $v \mapsto \tilde{u}(v, F)$ are Lipschitz continuous and therefore $\hat{\Upsilon}(u^o) + \tilde{\Upsilon}(u^o)$ is obviously bounded if u^o has bounded total variation.

2.3 The constrained Riemann problem

For completeness, we conclude this section by giving the definitions of the Riemann solvers \mathcal{R} and \mathcal{R}_F introduced in [5] and [12], associated to Riemann problem (3), (15) and to constrained Riemann problem (3), (5), (15), respectively, and used in Section 4 to prove Theorem 2.2.

We recall that Riemann problems for (3) are Cauchy problems with initial condition of the form

$$u(0, x) = \begin{cases} u_\ell & \text{if } x < 0, \\ u_r & \text{if } x > 0. \end{cases} \quad (15)$$

Definition 2.3. The Riemann solver $\mathcal{R}: \Omega^2 \rightarrow \mathbf{L}^\infty(\mathbb{R}; \Omega)$ associated to Riemann problem (3), (15) is defined as follows.

- (R.1) If $u_\ell, u_r \in \Omega_f$, then $\mathcal{R}[u_\ell, u_r]$ consists of a contact discontinuity (u_ℓ, u_r) with speed of propagation V .
- (R.2) If $u_\ell, u_r \in \Omega_c$, then $\mathcal{R}[u_\ell, u_r]$ consists of a 1-wave $(u_\ell, \mathbf{u}_*(u_\ell, u_r))$ and of a 2-contact discontinuity $(\mathbf{u}_*(u_\ell, u_r), u_r)$.

(R.3) If $u_\ell \in \Omega_c^-$ and $u_r \in \Omega_f^-$, then $\mathcal{R}[u_\ell, u_r]$ consists of a 1-rarefaction $(u_\ell, \omega(u_\ell))$ and a contact discontinuity $(\omega(u_\ell), u_r)$.

(R.4) If $u_\ell \in \Omega_f^-$ and $u_r \in \Omega_c^-$, then $\mathcal{R}[u_\ell, u_r]$ consists of a phase transition $(u_\ell, \mathbf{v}^-(u_r))$ and a 2-contact discontinuity $(\mathbf{v}^-(u_r), u_r)$.

Since $(t, x) \mapsto \mathcal{R}[u_\ell, u_r](x/t)$ does not in general satisfy constraint condition (5), we introduce

$$\begin{aligned} \mathcal{D}_1 &\doteq \{(u_\ell, u_r) \in \Omega \times \Omega : f(\mathcal{R}[u_\ell, u_r](t, 0_\pm)) \leq F\} \\ &= \{(u_\ell, u_r) \in \Omega_f \times \Omega_f : f(u_\ell) \leq F\} \\ &\quad \cup \{(u_\ell, u_r) \in \Omega_c \times \Omega : f(\mathbf{u}_*(u_\ell, u_r)) \leq F\} \\ &\quad \cup \{(u_\ell, u_r) \in \Omega_f^- \times \Omega_c^- : \min\{f(u_\ell), f(\mathbf{v}^-(u_r))\} \leq F\}, \end{aligned}$$

$\mathcal{D}_2 \doteq \Omega^2 \setminus \mathcal{D}_1$ and the constrained Riemann solver \mathcal{R}_F in the following

Definition 2.4. The constrained Riemann solver $\mathcal{R}_F: \Omega^2 \rightarrow \mathbf{L}^\infty(\mathbb{R}; \Omega)$ associated to constrained Riemann problem (3), (5), (15) is defined as

$$\mathcal{R}_F[u_\ell, u_r](x) \doteq \begin{cases} \mathcal{R}[u_\ell, u_r](x) & \text{if } (u_\ell, u_r) \in \mathcal{D}_1, \\ \begin{cases} \mathcal{R}[u_\ell, \hat{u}_\ell](x) & \text{if } x < 0, \\ \mathcal{R}[\check{u}_r, u_r](x) & \text{if } x > 0, \end{cases} & \text{if } (u_\ell, u_r) \in \mathcal{D}_2, \end{cases}$$

where $\hat{u}_\ell \doteq \hat{\mathbf{u}}(\mathbf{w}(u_\ell), F) \in \Omega_c$ and $\check{u}_r \doteq \check{\mathbf{u}}(v_r, F) \in \Omega$ are defined by (11).

In Figure 5 we clarify the selection criterion (11) for \hat{u}_ℓ and \check{u}_r . We point out that \hat{u}_ℓ and \check{u}_r satisfy the following




Figure 5: The selection criterion (11) for $\hat{u}_\ell \doteq \hat{\mathbf{u}}(\mathbf{w}(u_\ell), F)$ and $\check{u}_r \doteq \check{\mathbf{u}}(v_r, F)$ exploited in Definition 2.4 in the case $(u_\ell, u_r) \in \mathcal{D}_2$ and $F \in (0, f_c^-)$. In the first picture u_ℓ^1, u_ℓ^2 represent the left state in two different cases and $\hat{u}_\ell^1, \hat{u}_\ell^2$ are the corresponding \hat{u}_ℓ . Analogously in the second and third pictures for u_r^1, u_r^2 and $\check{u}_r^1, \check{u}_r^2$.

general properties.

- If $(u_\ell, u_r) \in \mathcal{D}_2$, then $\mathbf{w}(u_\ell) > \mathbf{w}(\check{u}_r)$ and $v_r > \hat{v}_\ell$.
- If $(u_\ell, u_r) \in \mathcal{D}_2$ and $u_\ell \in \Omega_f^-$, then $\mathbf{w}(\hat{u}_\ell) = w^-$.
- If $(u_\ell, u_r) \in \mathcal{D}_2$ and $u_r \in \Omega_f$, then $\check{v}_r = V$.

It is easy to prove that $(t, x) \mapsto \mathcal{R}[u_\ell, u_r](x/t)$ and $(t, x) \mapsto \mathcal{R}_F[u_\ell, u_r](x/t)$ are solutions to Riemann problems (3), (15) and (3), (5), (15) in the sense of Definitions 2.1 and 2.2, respectively.

We recall that both \mathcal{R} and \mathcal{R}_F are $\mathbf{L}_{\text{loc}}^1$ -continuous, see [12, Propositions 2 and 3].

3 Example

In this section we apply model (3), (4), (5) to simulate the traffic across, for instance, a toll gate located at $x = 0$ and with capacity F . More specifically, let w^- and w^+ be the Lagrangian markers corresponding to vehicles that are initially at rest in $[x_A, x_B]$ and $[x_B, 0)$, respectively. The resulting initial condition is

$$u^\circ(x) \doteq \begin{cases} u_\ell & \text{if } x \in [x_A, x_B), \\ u_r & \text{if } x \in [x_B, 0), \\ u_0 & \text{if } x \in \mathbb{R} \setminus [x_A, 0), \end{cases}$$

where $u_0 \doteq (0, V)$, $u_\ell \doteq (p^{-1}(w^-), 0)$ and $u_r \doteq (p^{-1}(w^+), 0)$, see Figure 6.




Figure 6: Notations used to describe the solution constructed in Section 3.

The resulting solution can be constructed by solving the Riemann problems corresponding to the discontinuities of u° and by considering the interactions of the waves between themselves or with the point constraint $x = 0$. We describe below the solution and its construction in more details. Let

$$\hat{u}_\ell \doteq \hat{u}(w^-, F), \quad \hat{u}_r \doteq \hat{u}(w^+, F), \quad \tilde{u} \doteq \tilde{u}(V, F), \quad u_* \doteq u_*(u_\ell, \hat{u}_r).$$

At $x = 0$ we apply \mathcal{R}_F and obtain a backward rarefaction $R_0(u_r, \hat{u}_r)$, a stationary non-classical shock $NS_0(\hat{u}_r, \tilde{u})$ and a forward contact discontinuity $CD_0(\tilde{u}, u_0)$, which moves with speed V . At $x = x_B$ we apply \mathcal{R} and obtain a stationary contact discontinuity $CD_B(u_\ell, u_r)$. Let C and E be the starting and final interaction points between CD_B and R_0 . During such interaction we have that CD_B accelerates, while R_0 crosses CD_B and eventually changes its values. After time $t = t_E$ we have that CD_B moves with speed $\hat{v}_r > 0$ and interacts with NS_0 at G . At G we apply \mathcal{R}_F and obtain a backward rarefaction $R_G(u_*, \hat{u}_\ell)$ and a stationary non-classical shock $NS_G(\hat{u}_\ell, \tilde{u})$.

At $x = x_A$ we apply \mathcal{R} and obtain a stationary phase transition $PT_A(u_0, u_\ell)$. Let D and F be the starting and final interaction points between PT_A and R_0 . During the time interval (t_D, t_F) we have that PT_A accelerates and R_0 starts to disappear. After time $t = t_F$ we have that PT_A moves with speed $\hat{v}_r > 0$. Let H and I be the starting and final interaction points between PT_A and R_G . Then, during the time interval (t_H, t_I) we have that PT_A accelerates and R_G starts to disappear. After time $t = t_I$ we have that PT_A moves with speed $\hat{v}_\ell > 0$. Finally, PT_A interacts with NS_G at L and then moves with speed V .

In Figure 7 we represent in different coordinates the quantitative evolution of the solution corresponding to $p(\rho) \doteq \rho^2$ and to the data

$$x_A = -8, \quad x_B = -5, \quad w^- = 1, \quad w^+ = 6/5, \quad V = 3/5, \quad F = \sqrt{3}/5.$$

Such solution is obtained by the explicit analysis of the wave-fronts interactions with computer-assisted computation of the interaction times and front slopes.

We finally observe that, once the overall picture of the solution is known, it is possible to express in a closed form the time at which the last vehicle passes through $x = 0$, indeed $t_L = [(x_B - x_A)\rho_\ell - x_B\rho_r]/F \approx 24.4716$.




Figure 7: The solution constructed in Section 3. Above we let $\mathbf{f}_* = f(\mathbf{u}_*)$.

4 Proof of Theorem 2.2

In this section we prove Theorem 2.2. More precisely, in Section 4.1 we construct a grid \mathcal{G}_n , approximate Riemann solvers $\mathcal{R}_n, \mathcal{R}_{F,n}$ and an approximate solution $u_n = (\rho_n, v_n)$ to constrained Cauchy problem (3), (4), (5). In Section 4.2 we prove that the approximate solution u_n is well defined globally in time by introducing a non-increasing Temple functional \mathcal{T}_n , which strictly decreases any time the number of the discontinuities of u_n increases. In Section 4.3 we prove that u_n converges to u , which is a solution to (3), (4), (5) and satisfies the estimates listed in (13). At last in Section 4.4 we consider the flux density of the non-classical shocks.

We choose to study the total variation in the (v, w) -coordinates rather than in the (ρ, v) -coordinates. This choice is in fact convenient to describe the grid, the approximate Riemann solvers and ease the forthcoming analysis, because the total variation of u_n in these coordinates does not increase after any interaction away from $x = 0$. Furthermore, the entropy pairs in the (v, w) -coordinates are well defined, but in the (ρ, v) -coordinates are multi-valued at the vacuum.

For simplicity we assume below that $n \in \mathbb{N}$ is sufficiently large. Moreover we simplify the notation by letting

$$\mathbf{w}_\ell \doteq \mathbf{w}(u_\ell), \quad \hat{\mathbf{u}}_\ell \doteq \hat{\mathbf{u}}(\mathbf{w}_\ell, F), \quad \check{\mathbf{u}}_\ell \doteq \check{\mathbf{u}}(v_\ell, F)$$

and so on, where $\hat{\mathbf{u}}$ and $\check{\mathbf{u}}$ are defined in (11).

4.1 The approximate solution

In this section we apply the wave-front tracking algorithm to construct an approximate solution u_n in the space \mathbf{PC} of piecewise constant functions taking finitely many values. To do so we introduce a grid \mathcal{G}_n in Ω and approximate Riemann solvers $\mathcal{R}_n, \mathcal{R}_{F,n} : \mathcal{G}_n \times \mathcal{G}_n \rightarrow \mathbf{PC}(\mathbb{R}; \mathcal{G}_n)$.

The grid

We introduce in Ω a grid $\mathcal{G}_n \doteq \Omega \cap \mathcal{P}$, see Figure 8, with \mathcal{P} given in the (v, w) -coordinates by

$$\left(\bigcup_{i=0}^{M \cdot 2^n} \{v^i\} \right) \times \left(\bigcup_{i=0}^{N \cdot 2^n} \{w^i\} \right),$$

where M, N, v^i and w^i , are defined as follows:




Figure 8: The grid \mathcal{G}_n corresponding to $F \in (0, f_c^-)$ and $n = 2$. The curve in the figure on the left is the support of Ξ_F , which corresponds to (a portion of) the horizontal line in the figure on the right.

- If $F = 0$, then we let $M = 1, N = 2$,

$$w^i \doteq \begin{cases} w^- - 1 + i 2^{-n} & \text{if } i \in \{0, \dots, 2^n\}, \\ w^- + (i - 2^n) 2^{-n} (w^+ - w^-) & \text{if } i \in \{2^n + 1, \dots, 2 \cdot 2^n\}, \end{cases}$$

and

$$v^i \doteq i 2^{-n} V \quad \text{if } i \in \{0, \dots, 2^n\}.$$

- If $F \in (0, f_c^-)$, then we let $M = 3, N = 3$,

$$w^i \doteq \begin{cases} w^- - 1 + i 2^{-n} (w_F - w^- + 1) & \text{if } i \in \{0, \dots, 2^n\}, \\ w_F + (i - 2^n) 2^{-n} (w^- - w_F) & \text{if } i \in \{2^n + 1, \dots, 2 \cdot 2^n\}, \\ w^- + (i - 2 \cdot 2^n) 2^{-n} (w^+ - w^-) & \text{if } i \in \{2 \cdot 2^n + 1, \dots, 3 \cdot 2^n\}, \end{cases}$$

and

$$v^i \doteq \begin{cases} i 2^{-n} v_F^- & \text{if } i \in \{0, \dots, 2^n\}, \\ \Xi_F^{-1}(w^{4 \cdot 2^n - i}) & \text{if } i \in \{2^n + 1, \dots, 2 \cdot 2^n\}, \\ v_F^+ + (i - 2 \cdot 2^n) 2^{-n} (V - v_F^+) & \text{if } i \in \{2 \cdot 2^n + 1, \dots, 3 \cdot 2^n\}. \end{cases}$$

- If $F \in [f_c^-, f_c^+]$, then we let $M = 2, N = 3$,

$$w^i \doteq \begin{cases} w^- - 1 + i 2^{-n} & \text{if } i \in \{0, \dots, 2^n\}, \\ w^- + (i - 2^n) 2^{-n} (w_F - w^-) & \text{if } i \in \{2^n + 1, \dots, 2 \cdot 2^n\}, \\ w_F + (i - 2 \cdot 2^n) 2^{-n} (w^+ - w_F) & \text{if } i \in \{2 \cdot 2^n + 1, \dots, 3 \cdot 2^n\}, \end{cases}$$

and

$$v^i \doteq \begin{cases} i 2^{-n} v_F^- & \text{if } i \in \{0, \dots, 2^n\}, \\ \Xi_F^{-1}(w^{4 \cdot 2^n - i}) & \text{if } i \in \{2^n + 1, \dots, 2 \cdot 2^n\}. \end{cases}$$

Notice that if $F \in \{f_c^-, f_c^+\}$, then not necessarily $w^i \neq w^{i+1}$.

The approximate Riemann solvers

An approximate solution $u_n \in \mathbf{PC}(\mathbb{R}; \mathcal{G}_n)$ to (3), (4), (5) is constructed by applying the approximate Riemann solvers $\mathcal{R}_n, \mathcal{R}_{F,n} : \mathcal{G}_n \times \mathcal{G}_n \rightarrow \mathbf{PC}(\mathbb{R}; \mathcal{G}_n)$, which are obtained by approximating the rarefactions. More precisely, for any $(u_\ell, u_r) \in \mathcal{G}_n \times \mathcal{G}_n$ such that $w_\ell = w_r$ and $v_\ell = v^h < v_r = v^{h+k}$, we let

$$\mathcal{R}_n[u_\ell, u_r](\xi) \doteq \begin{cases} u_\ell & \text{if } \xi \leq \Lambda(u_\ell, u_1), \\ u_j & \text{if } \Lambda(u_{j-1}, u_j) < \xi \leq \Lambda(u_j, u_{j+1}), \quad 1 \leq j \leq k-1, \\ u_r & \text{if } \xi > \Lambda(u_{k-1}, u_r), \end{cases}$$

where $u_0 \doteq u_\ell$, $u_k \doteq u_r$ and $u_j \in \mathcal{G}_n$ is such that $v_j \doteq v^{h+j}$ and $w_j = w_\ell$. The Riemann solver $\mathcal{R}_{F,n}$ is defined as follows:

1. If $f(\mathcal{R}_n[u_\ell, u_r](0_\pm)) \leq F$, then $\mathcal{R}_{F,n}[u_\ell, u_r] \doteq \mathcal{R}_n[u_\ell, u_r]$.
2. If $f(\mathcal{R}_n[u_\ell, u_r](0_\pm)) > F$, then

$$\mathcal{R}_{F,n}[u_\ell, u_r](\xi) \doteq \begin{cases} \mathcal{R}_n[u_\ell, \hat{u}_\ell](\xi) & \text{if } \xi < 0, \\ \mathcal{R}_n[\check{u}_r, u_r](\xi) & \text{if } \xi \geq 0. \end{cases}$$

The approximate solution

An approximate solution $u_n \in \mathbf{PC}(\mathbb{R}_+ \times \mathbb{R}; \mathcal{G}_n)$ to (3), (4), (5) can be constructed as follows. As a first step we approximate the initial datum u^o with $u_n^o \in \mathbf{PC}(\mathbb{R}; \mathcal{G}_n)$ such that

$$\begin{aligned} \|v_n^o\|_{\mathbf{L}^\infty} &\leq \|v^o\|_{\mathbf{L}^\infty}, \quad \text{TV}(v_n^o) \leq \text{TV}(v^o), \quad \lim_{n \rightarrow \infty} \|v_n^o - v^o\|_{\mathbf{L}_{\text{loc}}^1} = 0, \quad \hat{\Upsilon}(u_n^o) \leq C \hat{\Upsilon}(u^o), \\ \|\bar{w}_n^o\|_{\mathbf{L}^\infty} &\leq \|\bar{w}^o\|_{\mathbf{L}^\infty}, \quad \text{TV}(\bar{w}_n^o) \leq \text{TV}(\bar{w}^o), \quad \lim_{n \rightarrow \infty} \|\bar{w}_n^o - \bar{w}^o\|_{\mathbf{L}_{\text{loc}}^1} = 0, \quad \check{\Upsilon}(u_n^o) \leq C \check{\Upsilon}(u^o), \end{aligned} \quad (16)$$

for a constant C . The approximate solution u_n is then obtained by gluing together the approximate solutions computed by applying $\mathcal{R}_{F,n}$ at $x = 0$ at time $t = 0$ and at any time a wave-front reaches $x = 0$, and by applying \mathcal{R}_n at any discontinuity of u_n^o away from $x = 0$ or at any interaction between wave-fronts away from $x = 0$. As usual, in order to extend the construction globally in time we have to ensure that only finitely many interactions may occur in finite time. In Section 4.2 we prove that $u_n(t, \cdot)$ is well defined for all $t > 0$ and belongs to $\mathbf{PC}(\mathbb{R}_+ \times \mathbb{R}; \mathcal{G}_n)$. Finally, in Section 4.3 we prove that u_n converges (up to a subsequence) in $\mathbf{L}_{\text{loc}}^1$ to a limit u , which results to be a constrained solution to (3), (4), (5) in the sense of Definition 2.2.

4.2 A priori estimates

In this section we prove the main a priori estimates on the sequence of approximate solutions $(u_n)_n$. We prove in Proposition 4.1 that u_n takes values in \mathcal{G}_n and we estimate $\text{TV}(u_n(t, \cdot))$ uniformly in n and t . This together with Proposition 4.2 guarantee that the number of interactions and the number of the discontinuities of u_n are both bounded globally in time.

Observe that any Contact Discontinuity (CD) has non-negative speed (of propagation), any Shock (S) or Rarefaction Shock (RS) has negative speed, all the Non-classical Shocks (NSs) are stationary and the speed of all the possible Phase Transitions (PTs) ranges in the interval $(-f_c^- / (p^{-1}(w^-) - \rho^-), V)$. Below we say that (u_ℓ, u_r) is a null wave if $u_\ell = u_r$. Notice that if (u_ℓ, u_r) is a PT then $u_\ell \in \Omega_f^-$ and $u_r \in \Omega_c^-$, moreover if (u_ℓ, u_r) is a PT with $w_r > w^-$ then $\rho_\ell = 0$.

Let u_n be an approximate solution. Let $\sharp(t)$ be the number of waves/discontinuities of $u_n(t, \cdot)$ and introduce $\mathcal{T}_n : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ defined as

$$\mathcal{T}_n(t) \doteq \text{TV}(v_n(t, \cdot)) + \text{TV}(\bar{w}_n(t, \cdot)) + 2\hat{\Upsilon}_n(t) + 2\check{\Upsilon}_n(t),$$

where $\hat{\Upsilon}_n(t) \doteq \hat{\Upsilon}(u_n(t, \cdot))$ and $\check{\Upsilon}_n(t) \doteq \check{\Upsilon}(u_n(t, \cdot))$. Conventionally, we assume that u_n is left continuous in time, i.e. $u_n(t, \cdot) \equiv u_n(t_-, \cdot)$. Then also \mathcal{T}_n is left continuous in time. By the monotonicity of $w \mapsto \hat{v}(w)$, $w \mapsto \hat{w}(w)$, $v \mapsto \check{v}(v)$, $v \mapsto \check{w}(v)$, see Remark 2.3, and the definitions of $\hat{\Upsilon}$ and $\check{\Upsilon}$ given in (12), we have that

$$\begin{aligned} \hat{\Upsilon}_n(t) &= \text{TV}_+(\hat{v}(\bar{w}_n(t, \cdot)); (-\infty, 0)) + \text{TV}_-(\hat{w}(\bar{w}_n(t, \cdot)); (-\infty, 0)) \\ &= \sum_{x \in \text{CD}_n} \left\{ [\hat{v}(\bar{w}_n(t, x_+)) - \hat{v}(\bar{w}_n(t, x_-))]_+ + [\hat{w}(\bar{w}_n(t, x_-)) - \hat{w}(\bar{w}_n(t, x_+))]_+ \right\}, \end{aligned}$$

$$\begin{aligned}\check{Y}_n(t) &= \text{TV}_+ \left(\check{v}(v_n(t, \cdot), F); (0, \infty) \right) + \text{TV}_- \left(\check{w}(v_n(t, \cdot), F); (0, \infty) \right) \\ &= \sum_{x \in \text{RS}_n} \left\{ [\check{v}(v_n(t, x_+), F) - \check{v}(v_n(t, x_-), F)]_+ + [\check{w}(v_n(t, x_-), F) - \check{w}(v_n(t, x_+), F)]_+ \right\},\end{aligned}$$

where

$$\begin{aligned}\text{CD}_n &\doteq \left\{ x \in \mathbb{R} : (u_n(t, x_-), u_n(t, x_+)) \text{ is a CD in } x < 0 \text{ such that } \check{w}_n(t, x_-) > \max\{\check{w}_n(t, x_+), w_F\} \right\}, \\ \text{RS}_n &\doteq \left\{ x \in \mathbb{R} : (u_n(t, x_-), u_n(t, x_+)) \text{ is a RS in } x > 0 \text{ such that } v_n(t, x_+) > \max\{v_n(t, x_-), v_F^-\} \right\}.\end{aligned}$$

Let $\varepsilon_n > 0$ be the minimal (v, w) -distance between two points in the grid \mathcal{G}_n , namely

$$\varepsilon_n \doteq \min_{\substack{u^1, u^2 \in \mathcal{G}_n \\ u^1 \neq u^2}} \max\{|v^1 - v^2|, |\check{w}(u^1) - \check{w}(u^2)|\}.$$

The next proposition ensures that the number of discontinuities of u_n is uniformly bounded in time. Moreover, it gives uniform bounds on the total variation of the approximate solution, which allows us to use Helly's Theorem.

Proposition 4.1. *For any fixed $n \in \mathbb{N}$ sufficiently large and $u_n^o \in \mathbf{PC}(\mathbb{R}; \mathcal{G}_n)$, we have that:*

- (a) *the map $t \mapsto \mathcal{T}_n(t)$ is non-increasing and decreases by at least ε_n any time the number of waves increases;*
- (b) *$u_n(t, \cdot) \in \mathbf{PC}(\mathbb{R}; \mathcal{G}_n)$ for all $t > 0$.*

Proof. By construction for $t > 0$ sufficiently small $u_n(t, \cdot)$ belongs to $\mathbf{PC}(\mathbb{R}; \mathcal{G}_n)$, more precisely it is piecewise constant with jumps along a finite number of straight lines. If at time $t > 0$ an interaction occurs, namely two waves meet or a wave reaches $x = 0$, then the involved waves may change speed or strength, while new waves may be created. To prove that $u_n(t, \cdot)$ belongs to $\mathbf{PC}(\mathbb{R}; \mathcal{G}_n)$ we have to provide an a priori upper bound for the number of waves, which follows from (a).

Clearly, if at time $t > 0$ no interaction occurs then $\mathcal{T}_n(t) = \mathcal{T}_n(t_+)$. For this reason we consider below all the possible interactions and distinguish the following main cases:

- a single wave reaches $x = 0$ and no NS is involved;
- a single wave reaches $x = 0$ and a NS is involved;
- two waves interact away from $x = 0$;
- two waves interact at $x = 0$ and no NS is involved;
- two waves interact at $x = 0$ and a NS is involved.

For completeness we estimate

$$\begin{aligned}\Delta \text{TV}_v &\doteq \text{TV}(v_n(t_+, \cdot)) - \text{TV}(v_n(t, \cdot)), & \Delta \hat{Y}_n &\doteq \hat{Y}_n(\check{w}_n(t_+, \cdot)) - \hat{Y}_n(\check{w}_n(t, \cdot)), \\ \Delta \text{TV}_w &\doteq \text{TV}(\check{w}_n(t_+, \cdot)) - \text{TV}(\check{w}_n(t, \cdot)), & \Delta \check{Y}_n &\doteq \check{Y}_n(v_n(t_+, \cdot)) - \check{Y}_n(v_n(t, \cdot)),\end{aligned}$$

and

$$\Delta \# \doteq \#(t_+) - \#(t_-), \quad \Delta \mathcal{T}_n \doteq \mathcal{T}_n(t_+) - \mathcal{T}_n(t_-).$$

For simplicity in the exposition, whenever a NS is involved we consider separately the cases $F \in [f_c^-, f_c^+)$ and $F \in [0, f_c^-)$. Notice that $w^- > w_F$ if and only if $F < f_c^-$, or equivalently $V \neq v_F^+$. Notice also that if $F = f_c^+$ then $\mathcal{D}_2 = \emptyset$, while if $F = 0$ then $\mathcal{D}_1 = \emptyset$. At last, notice that if $F \in [f_c^-, f_c^+)$ and $(u_\ell, u_r) \in \mathcal{D}_2$, then $\hat{w}_\ell = \check{w}_\ell$ and $\check{v}_r = v_r$.

We start with the interaction estimates.

- If a wave (u_ℓ, u_r) reaches $x = 0$, $u_n(t, 0_-) = u_n(t, 0_+)$ and $(u_\ell, u_r) \in \mathcal{D}_1$, then the constraint has no influence on the wave and $0 = \Delta \text{TV}_v = \Delta \text{TV}_w = \Delta \#$. Since any CD has non-negative speed, we have that $\Delta \hat{Y}_n \leq 0$. Since any RS has negative speed, we have that $\Delta \check{Y}_n \leq 0$. As a consequence $\Delta \mathcal{T}_n \leq 0$.
- Assume that a wave (u_ℓ, u_r) reaches $x = 0$, $u_n(t, 0_-) = u_n(t, 0_+)$ and $(u_\ell, u_r) \in \mathcal{D}_2$. If $F \in [f_c^-, f_c^+)$, then one of the following cases occurs:

\mathbf{CD}_F^+ (u_ℓ, u_r) is a CD. In this case $\hat{v}_r \geq v_\ell = v_r = \check{v}_r > \hat{v}_\ell$, $\mathbf{w}_\ell = \hat{\mathbf{w}}_\ell > \check{\mathbf{w}}_r \geq \hat{\mathbf{w}}_r \geq \mathbf{w}_r$ and $f(u_\ell) > F \geq f(u_r)$. $\mathcal{R}_{F,n}[u_\ell, u_r]$ has at most three waves (u_ℓ, \hat{u}_ℓ) , $(\hat{u}_\ell, \check{u}_r)$ and (\check{u}_r, u_r) that are a S, a NS and a possibly null CD, respectively. As a consequence

$$\begin{aligned} \Delta\text{TV}_v &= 2(v_\ell - \hat{v}_\ell) > 0, & \Delta\hat{\Upsilon}_n &= -[\check{v}_r - \hat{v}_\ell]_+ - [\hat{\mathbf{w}}_\ell - \hat{\mathbf{w}}_r]_+ < -(\check{v}_r - \hat{v}_\ell) < 0, \\ \Delta\text{TV}_w &= 0, & \Delta\check{\Upsilon}_n &= 0, \end{aligned}$$

therefore $\Delta\sharp \in \{1, 2\}$ and $\Delta\mathcal{T}_n < -2(\check{v}_r - v_\ell) \leq 0$.

\mathbf{RS}_F^+ (u_ℓ, u_r) is a RS. In this case $v_\ell = \check{v}_\ell < v_r = \check{v}_r$, $\check{\mathbf{w}}_r < \mathbf{w}_\ell = \check{\mathbf{w}}_\ell = \mathbf{w}_r$, $f(u_\ell) = F < f(u_r)$ and $u_\ell, u_r \in \Omega_c$. $\mathcal{R}_{F,n}[u_\ell, u_r]$ has two waves (u_ℓ, \check{u}_r) and (\check{u}_r, u_r) that are a NS and a CD, respectively. As a consequence

$$\begin{aligned} \Delta\text{TV}_v &= 0, & \Delta\hat{\Upsilon}_n &= 0, \\ \Delta\text{TV}_w &= 2(\mathbf{w}_\ell - \check{\mathbf{w}}_r) > 0, & \Delta\check{\Upsilon}_n &= -[\check{v}_r - \check{v}_\ell]_+ - [\check{\mathbf{w}}_\ell - \check{\mathbf{w}}_r]_+ = -(v_r - v_\ell) - (\mathbf{w}_\ell - \check{\mathbf{w}}_r) < 0, \end{aligned}$$

therefore $\Delta\sharp = 1$ and $\Delta\mathcal{T}_n = -2(v_r - v_\ell) < 0$.

If $F \in [0, f_c^-)$, then one of the following cases occurs:

\mathbf{CD}_F^- (u_ℓ, u_r) is a CD. In this case $\hat{v}_r \geq v_\ell = v_r = \check{v}_r > \hat{v}_\ell$, $\hat{\mathbf{w}}_\ell \geq \mathbf{w}_\ell > \check{\mathbf{w}}_r \geq \hat{\mathbf{w}}_r \geq \mathbf{w}_r$ and $f(u_\ell) > F \geq f(u_r)$. $\mathcal{R}_{F,n}[u_\ell, u_r]$ has at most three waves (u_ℓ, \hat{u}_ℓ) , $(\hat{u}_\ell, \check{u}_r)$ and (\check{u}_r, u_r) that are a S or a PT, a NS and a possibly null CD, respectively. As a consequence

$$\begin{aligned} \Delta\text{TV}_v &= 2(v_\ell - \hat{v}_\ell) > 0, & \Delta\hat{\Upsilon}_n &= -[\check{v}_r - \hat{v}_\ell]_+ - [\hat{\mathbf{w}}_\ell - \hat{\mathbf{w}}_r]_+ = -(\hat{v}_r - \hat{v}_\ell) - (\hat{\mathbf{w}}_\ell - \hat{\mathbf{w}}_r) < 0, \\ \Delta\text{TV}_w &= 2(\hat{\mathbf{w}}_\ell - \mathbf{w}_\ell) \geq 0, & \Delta\check{\Upsilon}_n &= 0, \end{aligned}$$

therefore $\Delta\sharp \in \{1, 2\}$ and $\Delta\mathcal{T}_n = -2(\hat{v}_r - v_\ell) - 2(\mathbf{w}_\ell - \hat{\mathbf{w}}_r) < 0$.

\mathbf{RS}_F^- (u_ℓ, u_r) is a RS. In this case $v_\ell < v_r \leq \check{v}_r$, $w_F \leq \check{\mathbf{w}}_r < \mathbf{w}_r = \mathbf{w}_\ell = \check{\mathbf{w}}_\ell$, $f(u_\ell) = F < f(u_r)$ and $u_\ell, u_r \in \Omega_c$. $\mathcal{R}_{F,n}[u_\ell, u_r]$ has two waves (u_ℓ, \check{u}_r) and (\check{u}_r, u_r) that are a NS and a PT or a CD, respectively. As a consequence

$$\begin{aligned} \Delta\text{TV}_v &= 2(\check{v}_r - v_r) \geq 0, & \Delta\hat{\Upsilon}_n &= 0, \\ \Delta\text{TV}_w &= 2(\mathbf{w}_\ell - \check{\mathbf{w}}_r) > 0, & \Delta\check{\Upsilon}_n &= -[\check{v}_r - \check{v}_\ell]_+ - [\check{\mathbf{w}}_\ell - \check{\mathbf{w}}_r]_+ = -(\check{v}_r - v_\ell) - (\mathbf{w}_\ell - \check{\mathbf{w}}_r) < 0, \end{aligned}$$

therefore $\Delta\sharp = 1$ and $\Delta\mathcal{T}_n = -2(v_r - v_\ell) < 0$. Notice that $\check{v}_r > v_r$ if and only if $\mathbf{w}_\ell = \mathbf{w}_r = w^-$ and $v_r > v_\ell = v_F^+$.

Assume that two waves (u_ℓ, u_m) and (u_m, u_r) interact at time $t > 0$. Let $\mathbf{u}_* \doteq \mathbf{u}_*(u_\ell, u_r)$. Notice that $\mathbf{u}_* = u_r$ if and only if (u_ℓ, u_m) is a S or a RS, while $\mathbf{u}_* = u_\ell$ if and only if (u_ℓ, u_m) is a CD.

• If the interaction occurs at $x \neq 0$, then one of the following cases occurs:

CD-S (u_ℓ, u_m) is a CD and (u_m, u_r) is a S. In this case $v_\ell = v_m > v_r = v_*$, $\mathbf{w}_m = \mathbf{w}_r$, \mathbf{w}_* belongs to the closed interval between \mathbf{w}_ℓ and \mathbf{w}_r , $W(u_\ell) = W(\mathbf{u}_*)$, $\mathbf{w}_m = \mathbf{w}_r$, $f(u_m) > f(u_r)$ and $u_m, u_r \in \Omega_c$. $\mathcal{R}_n[u_\ell, u_r]$ has at most two waves (u_ℓ, \mathbf{u}_*) and (\mathbf{u}_*, u_r) that are respectively either a S and a CD, or a PT and a possibly null CD. As a consequence $0 = \Delta\text{TV}_v = \Delta\text{TV}_w = \Delta\hat{\Upsilon}_n = \Delta\check{\Upsilon}_n$, therefore $\Delta\sharp \leq 0$ and $\Delta\mathcal{T}_n = 0$.

CD-RS (u_ℓ, u_m) is a CD and (u_m, u_r) is a RS. In this case $v_\ell = v_m < v_r = v_*$, $\mathbf{w}_\ell = \mathbf{w}_*$, $\mathbf{w}_m = \mathbf{w}_r$, $f(u_\ell) < f(\mathbf{u}_*)$, $f(u_m) < f(u_r)$ and $u_\ell, u_m, \mathbf{u}_*, u_r \in \Omega_c$. $\mathcal{R}_n[u_\ell, u_r]$ has two waves (u_ℓ, \mathbf{u}_*) and (\mathbf{u}_*, u_r) that are a RS and a CD, respectively. As a consequence $0 = \Delta\text{TV}_v = \Delta\text{TV}_w = \Delta\hat{\Upsilon}_n = \Delta\check{\Upsilon}_n$, therefore $\Delta\sharp = 0$ and $\Delta\mathcal{T}_n = 0$.

CD-PT (u_ℓ, u_m) is a CD and (u_m, u_r) is a PT. In this case $v_\ell = v_m = V > v_r = v_*$, $\mathbf{w}_m < w^- \leq \mathbf{w}_r$, \mathbf{w}_* belongs to the closed interval between \mathbf{w}_ℓ and \mathbf{w}_r , $u_\ell \in \Omega_f$, $u_m \in \Omega_f^-$ and $\mathbf{u}_*, u_r \in \Omega_c$. $\mathcal{R}_n[u_\ell, u_r]$ has at most two waves (u_ℓ, \mathbf{u}_*) and (\mathbf{u}_*, u_r) that are either a PT or a S and a possibly null CD, respectively. As a consequence

$$\begin{aligned} \Delta\text{TV}_v &= 0, & \Delta\hat{\Upsilon}_n &\leq 0, \\ \Delta\text{TV}_w &= |\mathbf{w}_\ell - \mathbf{w}_r| - (|\mathbf{w}_\ell - \mathbf{w}_m| + |\mathbf{w}_m - \mathbf{w}_r|) \leq 0, & \Delta\check{\Upsilon}_n &= 0, \end{aligned}$$

therefore $\Delta\sharp \leq 0$ and $\Delta\mathcal{T}_n \leq 0$.

S-S (u_ℓ, u_m) and (u_m, u_r) are Ss. In this case $v_\ell > v_m > v_r$, $\mathbf{w}_\ell = \mathbf{w}_m = \mathbf{w}_r \geq w^-$ and $u_\ell, u_m, u_r \in \Omega_c$. $\mathcal{R}_n[u_\ell, u_r]$ has one wave (u_ℓ, u_r) , which is a S. As a consequence $0 = \Delta\text{TV}_v = \Delta\text{TV}_w = \Delta\hat{\Upsilon}_n = \Delta\check{\Upsilon}_n$, therefore $\Delta\sharp = -1$ and $\Delta\mathcal{T}_n = 0$.

S-RS (u_ℓ, u_m) is a S and (u_m, u_r) is a RS. In this case $v_\ell > v_r > v_m$, $\mathbf{w}_\ell = \mathbf{w}_m = \mathbf{w}_r \geq w^-$ and $u_\ell, u_m, u_r \in \Omega_c$. $\mathcal{R}_n[u_\ell, u_r]$ has one wave (u_ℓ, u_r) , which is a S. As a consequence

$$\begin{aligned}\Delta\text{TV}_v &= -2(v_r - v_m) < 0, & \Delta\hat{\Upsilon}_n &= 0, \\ \Delta\text{TV}_w &= 0, & \Delta\check{\Upsilon}_n &\leq 0,\end{aligned}$$

therefore $\Delta\sharp = -1$ and $\Delta\mathcal{T}_n < 0$.

RS-S (u_ℓ, u_m) is a RS and (u_m, u_r) is a S. In this case $v_m > v_\ell > v_r$, $\mathbf{w}_\ell = \mathbf{w}_m = \mathbf{w}_r \geq w^-$ and $u_\ell, u_m, u_r \in \Omega_c$. $\mathcal{R}_n[u_\ell, u_r]$ has one wave (u_ℓ, u_r) , which is a S. As a consequence

$$\begin{aligned}\Delta\text{TV}_v &= -2(v_m - v_\ell) < 0, & \Delta\hat{\Upsilon}_n &= 0, \\ \Delta\text{TV}_w &= 0, & \Delta\check{\Upsilon}_n &\leq 0,\end{aligned}$$

therefore $\Delta\sharp = -1$ and $\Delta\mathcal{T}_n < 0$.

PT-S (u_ℓ, u_m) is a PT and (u_m, u_r) is a S. In this case $v_\ell = V > v_m > v_r$, $\mathbf{w}_\ell < w^- \leq \mathbf{w}_m = \mathbf{w}_r$, $u_\ell \in \Omega_f^-$ and $u_m, u_r \in \Omega_c$. $\mathcal{R}_n[u_\ell, u_r]$ has one wave (u_ℓ, u_r) , which is a PT. As a consequence $0 = \Delta\text{TV}_v = \Delta\text{TV}_w = \Delta\hat{\Upsilon}_n = \Delta\check{\Upsilon}_n$, therefore $\Delta\sharp = -1$ and $\Delta\mathcal{T}_n = 0$.

PT-RS (u_ℓ, u_m) is a PT and (u_m, u_r) is a RS. In this case $v_\ell = V \geq v_r > v_m$, $\mathbf{w}_\ell < w^- \leq \mathbf{w}_m = \mathbf{w}_r$, $u_\ell \in \Omega_f^-$ and $u_m, u_r \in \Omega_c$. $\mathcal{R}_n[u_\ell, u_r]$ has one wave (u_ℓ, u_r) , which is either a PT or a CD. As a consequence

$$\begin{aligned}\Delta\text{TV}_v &= -2(v_r - v_m) < 0, & \Delta\hat{\Upsilon}_n &= 0, \\ \Delta\text{TV}_w &= 0, & \Delta\check{\Upsilon}_n &\leq 0,\end{aligned}$$

therefore $\Delta\sharp = -1$ and $\Delta\mathcal{T}_n < 0$.

• If the interaction occurs at $x = 0$ and $(u_\ell, u_r) \in \mathcal{D}_1$, then one of the following cases occurs:

CD-S₀ (u_ℓ, u_m) is a CD and (u_m, u_r) is a S. In this case $v_\ell = v_m > v_r = v_*$, $\mathbf{w}_m = \mathbf{w}_r$, \mathbf{w}_* belongs to the closed interval between \mathbf{w}_ℓ and \mathbf{w}_r , $\mathbf{W}(u_\ell) = \mathbf{W}(\mathbf{u}_*)$, $\mathbf{w}_m = \mathbf{w}_r$, $f(u_r) < (u_m) \leq F$, $\min\{f(u_\ell), f(\mathbf{u}_*)\} \leq F$ and $u_m, u_r \in \Omega_c$. $\mathcal{R}_{F,n}[u_\ell, u_r]$ has at most two waves (u_ℓ, \mathbf{u}_*) and (\mathbf{u}_*, u_r) that are respectively either a S and a CD, or a PT and a possibly null CD. As a consequence $\Delta\hat{\Upsilon}_n \leq 0 = \Delta\text{TV}_v = \Delta\text{TV}_w = \Delta\check{\Upsilon}_n$, therefore $\Delta\sharp \leq 0$ and $\Delta\mathcal{T}_n \leq 0$.

CD-RS₀ (u_ℓ, u_m) is a CD and (u_m, u_r) is a RS. In this case $v_\ell = v_m < v_r$, $\mathbf{w}_m = \mathbf{w}_r$, $\mathbf{w}_* = \mathbf{w}_\ell$, $f(u_\ell) < f(\mathbf{u}_*)$, $f(u_m) < f(u_r)$, $\max\{f(u_m), f(\mathbf{u}_*)\} \leq F$ and $u_\ell, u_m, \mathbf{u}_*, u_r \in \Omega_c$. $\mathcal{R}_{F,n}[u_\ell, u_r]$ has two waves (u_ℓ, \mathbf{u}_*) and (\mathbf{u}_*, u_r) that are a RS and a CD, respectively. As a consequence

$$\begin{aligned}\Delta\text{TV}_v &= 0, & \Delta\hat{\Upsilon}_n &\leq 0, \\ \Delta\text{TV}_w &= 0, & \Delta\check{\Upsilon}_n &\leq 0,\end{aligned}$$

therefore $\Delta\sharp = 0$ and $\Delta\mathcal{T}_n \leq 0$.

CD-NS₀ (u_ℓ, u_m) is a CD and (u_m, u_r) is a NS. In this case $v_F^- \leq v_\ell = v_m < v_r \leq v_F^+$, $w^- \leq \mathbf{w}_\ell \leq \mathbf{w}_r < \mathbf{w}_m$, $f(u_\ell) < f(\mathbf{u}_*) \leq F = f(u_m) = f(u_r)$ and $u_\ell, u_m, \mathbf{u}_*, u_r \in \Omega_c$. $\mathcal{R}_{F,n}[u_\ell, u_r]$ has a fan of RSs ranging from u_ℓ to \mathbf{u}_* and a possibly null CD (\mathbf{u}_*, u_r) . As a consequence $\Delta\text{TV}_w = -2(\mathbf{w}_m - \mathbf{w}_r) < 0 = \Delta\text{TV}_v = \Delta\hat{\Upsilon}_n = \Delta\check{\Upsilon}_n$, therefore $\Delta\sharp \in [-1, 2^n - 1]$ and $\Delta\mathcal{T}_n < 0$.

CD-PT₀ (u_ℓ, u_m) is a CD and (u_m, u_r) is a PT. In this case $v_\ell = v_m = V > v_r = v_*$, $\mathbf{w}_m < w^- \leq \mathbf{w}_r$, \mathbf{w}_* belongs to the closed interval between \mathbf{w}_ℓ and \mathbf{w}_r , $\min\{f(u_\ell), f(\mathbf{u}_*)\} \leq F$, $\max\{f(u_m), f(u_r)\} \leq F$, $u_\ell \in \Omega_f^-$, $u_m \in \Omega_f^-$ and $\mathbf{u}_*, u_r \in \Omega_c$. $\mathcal{R}_{F,n}[u_\ell, u_r]$ has at most two waves (u_ℓ, \mathbf{u}_*) and (\mathbf{u}_*, u_r) that are either a PT or a S and a possibly null CD, respectively. As a consequence

$$\begin{aligned}\Delta\text{TV}_v &= 0, & \Delta\hat{\Upsilon}_n &\leq 0, \\ \Delta\text{TV}_w &= |\mathbf{w}_\ell - \mathbf{w}_r| - (|\mathbf{w}_\ell - \mathbf{w}_m| + |\mathbf{w}_m - \mathbf{w}_r|) \leq 0, & \Delta\check{\Upsilon}_n &\leq 0,\end{aligned}$$

therefore $\Delta\sharp \in \{-1, 0\}$ and $\Delta\mathcal{T}_n \leq 0$.

S-S₀ (u_ℓ, u_m) and (u_m, u_r) are Ss. In this case $v_\ell > v_m > v_r$, $\mathbf{w}_\ell = \mathbf{w}_m = \mathbf{w}_r \geq w^-$, $f(u_r) < f(u_\ell) \leq F$ and $u_\ell, u_m, u_r \in \Omega_c$. $\mathcal{R}_{F,n}[u_\ell, u_r]$ has one wave (u_ℓ, u_r) , which is a S. As a consequence $0 = \Delta\text{TV}_v = \Delta\text{TV}_w = \Delta\hat{\Upsilon}_n = \Delta\check{\Upsilon}_n$, therefore $\Delta\sharp = -1$ and $\Delta\mathcal{T}_n = 0$.

S-RS₀ (u_ℓ, u_m) is a S and (u_m, u_r) is a RS. In this case $v_\ell > v_r > v_m$, $\mathbf{w}_\ell = \mathbf{w}_m = \mathbf{w}_r \geq w^-$, $f(u_r) < f(u_\ell) \leq F$ and $u_\ell, u_m, u_r \in \Omega_c$. $\mathcal{R}_{F,n}[u_\ell, u_r]$ has one wave (u_ℓ, u_r) , which is a S. As a consequence

$$\begin{aligned}\Delta\text{TV}_v &= -2(v_r - v_m) < 0, & \Delta\hat{\Upsilon}_n &= 0, \\ \Delta\text{TV}_w &= 0, & \Delta\check{\Upsilon}_n &\leq 0,\end{aligned}$$

therefore $\Delta\sharp = -1$ and $\Delta\mathcal{T}_n < 0$.

RS-S₀ (u_ℓ, u_m) is a RS and (u_m, u_r) is a S. In this case $v_m > v_\ell > v_r$, $\mathbf{w}_\ell = \mathbf{w}_m = \mathbf{w}_r \geq w^-$, $f(u_r) < f(u_\ell) \leq F$ and $u_\ell, u_m, u_r \in \Omega_c$. $\mathcal{R}_{F,n}[u_\ell, u_r]$ has one wave (u_ℓ, u_r) , which is a S. As a consequence

$$\begin{aligned}\Delta\text{TV}_v &= -2(v_m - v_\ell) < 0, & \Delta\hat{\Upsilon}_n &= 0, \\ \Delta\text{TV}_w &= 0, & \Delta\check{\Upsilon}_n &\leq 0,\end{aligned}$$

therefore $\Delta\sharp = -1$ and $\Delta\mathcal{T}_n < 0$.

NS-S₀ (u_ℓ, u_m) is a NS and (u_m, u_r) is a S. In this case $v_m > v_\ell \geq v_r = v_*$, $\mathbf{w}_* = \mathbf{w}_\ell > \mathbf{w}_m = \mathbf{w}_r \geq w^-$, $f(u_\ell) = f(u_m) = F > f(u_r)$ and $u_\ell, u_m, u_r \in \Omega_c$. $\mathcal{R}_{F,n}[u_\ell, u_r]$ has at most two waves (u_ℓ, \mathbf{u}_*) and (\mathbf{u}_*, u_r) that are a possibly null S and CD, respectively. As a consequence $\Delta\text{TV}_v = -2(v_m - v_\ell) < 0 = \Delta\text{TV}_w = \Delta\hat{\Upsilon}_n = \Delta\check{\Upsilon}_n$, therefore $\Delta\sharp \in \{-1, 0\}$ and $\Delta\mathcal{T}_n < 0$.

NS-PT₀ (u_ℓ, u_m) is a NS and (u_m, u_r) is a PT. In this case $v_m = V > v_\ell \geq v_r = v_*$, $\mathbf{w}_\ell = \mathbf{w}_* \geq \mathbf{w}_r = w^- > \mathbf{w}_m$, $f_c^- > f(u_\ell) = f(u_m) = F > f(u_r)$, $f(\mathbf{u}_*) \leq F$, $u_\ell, u_r \in \Omega_c$ and $u_m \in \Omega_f^-$. $\mathcal{R}_{F,n}[u_\ell, u_r]$ has at most two waves (u_ℓ, \mathbf{u}_*) and (\mathbf{u}_*, u_r) that are a possibly null S and a possibly null CD (but not both null), respectively. As a consequence

$$\begin{aligned}\Delta\text{TV}_v &= -2(V - v_\ell) < 0, & \Delta\hat{\Upsilon}_n &= 0, \\ \Delta\text{TV}_w &= -2(w^- - \mathbf{w}_m) < 0, & \Delta\check{\Upsilon}_n &= 0,\end{aligned}$$

therefore $\Delta\sharp \in \{-1, 0\}$ and $\Delta\mathcal{T}_n < 0$.

PT-S₀ (u_ℓ, u_m) is a PT and (u_m, u_r) is a S. In this case $v_\ell = V > v_m > v_r$, $\mathbf{w}_\ell < w^- \leq \mathbf{w}_m = \mathbf{w}_r$, $f(u_r) < f(u_m) \leq \max\{f(u_\ell), f(u_m)\} \leq F$, $u_\ell \in \Omega_f^-$ and $u_m, u_r \in \Omega_c^-$. $\mathcal{R}_{F,n}[u_\ell, u_r]$ has one wave (u_ℓ, u_r) , which is a PT. As a consequence $0 = \Delta\text{TV}_v = \Delta\text{TV}_w = \Delta\hat{\Upsilon}_n = \Delta\check{\Upsilon}_n$, therefore $\Delta\sharp = -1$ and $\Delta\mathcal{T}_n = 0$.

PT-RS₀ (u_ℓ, u_m) is a PT and (u_m, u_r) is a RS. In this case $v_\ell = V \geq v_r > v_m$, $\mathbf{w}_\ell < w^- \leq \mathbf{w}_m = \mathbf{w}_r$, $u_\ell \in \Omega_f^-$ and $u_m, u_r \in \Omega_c$. $\mathcal{R}_{F,n}[u_\ell, u_r]$ has one wave (u_ℓ, u_r) , which is either a PT or a CD. As a consequence

$$\begin{aligned}\Delta\text{TV}_v &= -2(v_r - v_m) < 0, & \Delta\hat{\Upsilon}_n &= 0, \\ \Delta\text{TV}_w &= 0, & \Delta\check{\Upsilon}_n &\leq 0,\end{aligned}$$

therefore $\Delta\sharp = -1$ and $\Delta\mathcal{T}_n < 0$.

PT-NS₀ (u_ℓ, u_m) is a PT and (u_m, u_r) is a NS. In this case $v_\ell = V \geq v_r > v_m$, $\mathbf{w}_\ell < w^- \leq \mathbf{w}_r < \mathbf{w}_m$, $f(u_\ell) < f(u_m) = f(u_r) = F$, $u_\ell \in \Omega_f^-$ and $u_m, u_r \in \Omega_c$. $\mathcal{R}_{F,n}[u_\ell, u_r]$ has one wave (u_ℓ, u_r) , which is either a CD or a PT. As a consequence

$$\begin{aligned}\Delta\text{TV}_v &= -2(v_r - v_m) < 0, & \Delta\hat{\Upsilon}_n &= 0, \\ \Delta\text{TV}_w &= -2(\mathbf{w}_m - \mathbf{w}_r) < 0, & \Delta\check{\Upsilon}_n &= 0,\end{aligned}$$

therefore $\Delta\sharp = -1$ and $\Delta\mathcal{T}_n < 0$.

• Assume that two waves (u_ℓ, u_m) and (u_m, u_r) interact at $x = 0$ and $(u_\ell, u_r) \in \mathcal{D}_2$. If $F \in [f_c^-, f_c^+)$, then one of the following cases occurs:

CD-S_F⁺ (u_ℓ, u_m) is a CD and (u_m, u_r) is a S. In this case $\hat{v}_r \geq v_\ell = v_m > v_r = \check{v}_r > \hat{v}_\ell$, $\mathbf{w}_\ell = \hat{\mathbf{w}}_\ell > \check{\mathbf{w}}_r > \mathbf{w}_m = \mathbf{w}_r$, $\mathbf{w}_\ell > \hat{\mathbf{w}}_r$, $f(u_\ell) > f(\mathbf{u}_*) > F \geq f(u_m) > f(u_r)$ and $u_\ell, u_m, u_r \in \Omega_c$. $\mathcal{R}_{F,n}[u_\ell, u_r]$ has three waves $(u_\ell, \hat{\mathbf{u}}_\ell)$, $(\hat{\mathbf{u}}_\ell, \check{\mathbf{u}}_r)$ and $(\check{\mathbf{u}}_r, u_r)$, which are a S, a NS and a CD, respectively. As a consequence

$$\begin{aligned}\Delta\text{TV}_v &= 2(v_r - \hat{v}_\ell) > 0, & \Delta\hat{\Upsilon}_n &= -(\hat{v}_m - \hat{v}_\ell) - (\mathbf{w}_\ell - \hat{\mathbf{w}}_m) < 0, \\ \Delta\text{TV}_w &= 0, & \Delta\check{\Upsilon}_n &= 0,\end{aligned}$$

therefore $\Delta\sharp = 1$ and $\Delta\mathcal{T}_n = -2(\hat{v}_r - v_r) - 2(\mathbf{w}_\ell - \hat{\mathbf{w}}_r) < 0$.

CD-RS $_F^+$ (u_ℓ, u_m) is a CD and (u_m, u_r) is a RS. In this case $\hat{v}_\ell \leq v_\ell = v_m = \check{v}_m < v_r = \check{v}_r$, $\hat{v}_\ell < \hat{v}_m$, $\mathbf{w}_\ell = \hat{\mathbf{w}}_\ell > \hat{\mathbf{w}}_m \geq \mathbf{w}_m = \mathbf{w}_r$, $\check{\mathbf{w}}_r < \check{\mathbf{w}}_m$, $f(u_m) < f(u_r) \leq f(u_\ell) < f(\mathbf{u}_*)$, $f(u_m) \leq F < f(\mathbf{u}_*)$ and $u_\ell, u_m, u_r \in \Omega_c$. $\mathcal{R}_{F,n}[u_\ell, u_r]$ has at most three waves (u_ℓ, \hat{u}_ℓ), ($\hat{u}_\ell, \check{u}_r$) and (\check{u}_r, u_r), which are a possibly null S, a NS and a possibly null CD, respectively. As a consequence

$$\begin{aligned} \Delta\text{TV}_v &= 2(v_\ell - \hat{v}_\ell) \geq 0, & \Delta\hat{\Upsilon}_n &= -(\hat{v}_m - \hat{v}_\ell) - (\hat{\mathbf{w}}_\ell - \hat{\mathbf{w}}_m) < -(\hat{v}_m - \hat{v}_\ell) < 0, \\ \Delta\text{TV}_w &= \begin{cases} 2(\check{\mathbf{w}}_r - \check{\mathbf{w}}_r) & \text{if } f(u_m) = F \\ 0 & \text{if } f(u_m) < F \end{cases} \geq 0, & \Delta\check{\Upsilon}_n &= -(v_r - v_m) - (\check{\mathbf{w}}_m - \check{\mathbf{w}}_r) < -(\check{\mathbf{w}}_m - \check{\mathbf{w}}_r) < 0, \end{aligned}$$

and therefore

$$\Delta\sharp \in \{-1, 0, 1\}, \quad \Delta\mathcal{T}_n < -2(\hat{v}_m - v_\ell) + \begin{cases} 0 & \text{if } f(u_m) = F \\ -2(\check{\mathbf{w}}_m - \check{\mathbf{w}}_r) & \text{if } f(u_m) < F \end{cases} \leq 0.$$

CD-NS $_F^+$ (u_ℓ, u_m) is a CD and (u_m, u_r) is a NS. In this case $v_\ell = v_m < v_r$, $\mathbf{w}_r < \min\{\mathbf{w}_\ell, \mathbf{w}_m\}$, $\mathbf{w}_\ell \neq \mathbf{w}_m$, $f(u_m) = f(u_r) = F \neq f(u_\ell)$ and $u_\ell, u_m, u_r \in \Omega_c$.

If $f(u_\ell) < F$, then $\mathcal{R}_{F,n}[u_\ell, u_r]$ has a fan of RSs from u_ℓ to \hat{u}_ℓ and a NS (\hat{u}_ℓ, u_r); as a consequence $\Delta\text{TV}_w = -2(\mathbf{w}_m - \mathbf{w}_\ell) < 0 = \Delta\text{TV}_v = \Delta\hat{\Upsilon}_n = \Delta\check{\Upsilon}_n$, therefore $\Delta\sharp \in [0, 2^n - 2]$ and $\Delta\mathcal{T}_n = -2(\mathbf{w}_m - \mathbf{w}_\ell) < 0$.

If $f(u_\ell) > F$, then $\hat{v}_\ell < v_\ell = v_m = \hat{v}_m$, $\hat{\mathbf{w}}_\ell = \mathbf{w}_\ell > \mathbf{w}_m = \hat{\mathbf{w}}_m$, $\mathcal{R}_{F,n}[u_\ell, u_r]$ has a two waves (u_ℓ, \hat{u}_ℓ) and (\hat{u}_ℓ, u_r), that are a S and a NS, respectively; as a consequence

$$\begin{aligned} \Delta\text{TV}_v &= 2(v_\ell - \hat{v}_\ell) > 0, & \Delta\hat{\Upsilon}_n &= -(v_\ell - \hat{v}_\ell) - (\mathbf{w}_\ell - \mathbf{w}_m) < 0, \\ \Delta\text{TV}_w &= 0, & \Delta\check{\Upsilon}_n &= 0, \end{aligned}$$

therefore $\Delta\sharp = 0$ and $\Delta\mathcal{T}_n = -2(\mathbf{w}_\ell - \mathbf{w}_m) < 0$.

CD-PT $_F^+$ (u_ℓ, u_m) is a CD and (u_m, u_r) is a PT. In this case $\mathbf{w}_m < w^- \leq w_F = \hat{\mathbf{w}}_m < \mathbf{w}_\ell$, $v_\ell = v_m = \hat{v}_m = V > v_r = \check{v}_r > \hat{v}_\ell$, $\mathbf{w}_m < w^- \leq \mathbf{w}_r < \check{\mathbf{w}}_r < \mathbf{w}_\ell = \hat{\mathbf{w}}_\ell$, $f(u_\ell) > f(\mathbf{u}_*) > F \geq \max\{f(u_m), f(u_r)\}$, $u_\ell \in \Omega_c^+$, $u_m \in \Omega_c^-$ and $u_r \in \Omega_c$. $\mathcal{R}_{F,n}[u_\ell, u_r]$ has three waves (u_ℓ, \hat{u}_ℓ), ($\hat{u}_\ell, \check{u}_r$) and (\check{u}_r, u_r) that are a S, a NS and a CD, respectively. As a consequence

$$\begin{aligned} \Delta\text{TV}_v &= 2(v_r - \hat{v}_\ell) > 0, & \Delta\hat{\Upsilon}_n &= -(V - \hat{v}_\ell) - (\mathbf{w}_\ell - w_F) < 0, \\ \Delta\text{TV}_w &= -2(\mathbf{w}_r - \mathbf{w}_m) < 0, & \Delta\check{\Upsilon}_n &= 0, \end{aligned}$$

therefore $\Delta\sharp = 1$ and $\Delta\mathcal{T}_n < -2(V - v_r) - 2(\mathbf{w}_\ell - w_F) < 0$.

NS-S $_F^+$ (u_ℓ, u_m) is a NS and (u_m, u_r) is a S. In this case $v_m > v_r = \check{v}_r > v_\ell$, $\mathbf{w}_\ell > \check{\mathbf{w}}_r > \mathbf{w}_m = \mathbf{w}_r \geq w_F \geq w^-$, $f(\mathbf{u}_*) > F = f(u_\ell) = f(u_m) > f(u_r)$ and $u_\ell, u_m, u_r \in \Omega_c$. $\mathcal{R}_{F,n}[u_\ell, u_r]$ has two waves (u_ℓ, \check{u}_r) and (\check{u}_r, u_r) that are a NS and a CD, respectively. As a consequence $\Delta\text{TV}_v = -2(v_m - v_r) < 0 = \Delta\text{TV}_w = \Delta\hat{\Upsilon}_n = \Delta\check{\Upsilon}_n$, therefore $\Delta\sharp = 0$ and $\Delta\mathcal{T}_n = -2(v_m - v_r) < 0$.

NS-RS $_F^+$ (u_ℓ, u_m) is a NS and (u_m, u_r) is a RS. In this case $v_\ell < v_m = \check{v}_m < v_r = \check{v}_r$, $\mathbf{w}_\ell > \mathbf{w}_m = \check{\mathbf{w}}_m = \mathbf{w}_r > \check{\mathbf{w}}_r$, $f(\mathbf{u}_*) > f(u_r) > F = f(u_\ell) = f(u_m)$ and $u_\ell, u_m, u_r \in \Omega_c$. $\mathcal{R}_{F,n}[u_\ell, u_r]$ has two waves (u_ℓ, \check{u}_r) and (\check{u}_r, u_r) that are a NS and a CD, respectively. As a consequence

$$\begin{aligned} \Delta\text{TV}_v &= 0, & \Delta\hat{\Upsilon}_n &= 0, \\ \Delta\text{TV}_w &= 2(\mathbf{w}_r - \check{\mathbf{w}}_r) > 0, & \Delta\check{\Upsilon}_n &= -(v_r - v_m) - (\mathbf{w}_r - \check{\mathbf{w}}_r) < 0, \end{aligned}$$

therefore $\Delta\sharp = 0$ and $\Delta\mathcal{T}_n = -2(v_r - v_m) < 0$.

If $F \in [0, f_c^-)$, then one of the following cases occurs:

CD-S $_F^-$ (u_ℓ, u_m) is a CD and (u_m, u_r) is a S. In this case $\hat{v}_m \geq v_\ell = v_m > v_r = \check{v}_r > \hat{v}_\ell$, $\mathbf{w}_\ell = \hat{\mathbf{w}}_\ell > \check{\mathbf{w}}_r > \mathbf{w}_m = \hat{\mathbf{w}}_m = \mathbf{w}_r \geq w^-$, $f(u_\ell) > f(\mathbf{u}_*) > F \geq f(u_m) > f(u_r)$ and $u_\ell, u_m, u_r \in \Omega_c$. $\mathcal{R}_{F,n}[u_\ell, u_r]$ has three waves (u_ℓ, \hat{u}_ℓ), ($\hat{u}_\ell, \check{u}_r$) and (\check{u}_r, u_r), which are a S, a NS and a CD, respectively. As a consequence

$$\begin{aligned} \Delta\text{TV}_v &= 2(v_r - \hat{v}_\ell) > 0, & \Delta\hat{\Upsilon}_n &= -(\hat{v}_m - \hat{v}_\ell) - (\mathbf{w}_\ell - \mathbf{w}_m) < 0, \\ \Delta\text{TV}_w &= 0, & \Delta\check{\Upsilon}_n &= 0, \end{aligned}$$

therefore $\Delta\sharp = 1$ and $\Delta\mathcal{T}_n = -2(\hat{v}_m - v_r) - 2(\mathbf{w}_\ell - \mathbf{w}_m) < 0$.

CD-RS $\bar{\mathbf{F}}$ (u_ℓ, u_m) is a CD and (u_m, u_r) is a RS. In this case $\hat{v}_\ell \leq v_\ell = v_m \leq \hat{v}_m$, $v_m = \check{v}_m < v_r \leq \check{v}_r$, $\mathbf{w}_\ell = \hat{\mathbf{w}}_\ell \geq \check{\mathbf{w}}_m \geq \mathbf{w}_m = \mathbf{w}_r \geq w^-$, $\mathbf{w}_\ell > \mathbf{w}_m$, $f(u_m) < f(u_r) \leq f(u_\ell) < f(\mathbf{u}_*)$, $f(u_m) \leq F < f(\mathbf{u}_*)$ and $u_\ell, u_m, u_r \in \Omega_c$. $\mathcal{R}_{F,n}[u_\ell, u_r]$ has at most three waves (u_ℓ, \hat{u}_ℓ), ($\hat{u}_\ell, \check{u}_r$) and (\check{u}_r, u_r), which are a possibly null S, a NS and a possibly null CD or PT, respectively. As a consequence

$$\begin{aligned} \Delta \text{TV}_v &= 2(v_\ell - \hat{v}_\ell) + 2(\check{v}_r - v_r) \geq 0, & \Delta \hat{\Upsilon}_n &= -(\hat{v}_m - \hat{v}_\ell) - (\mathbf{w}_\ell - \mathbf{w}_m) < 0, \\ \Delta \text{TV}_w &= \begin{cases} 2(\mathbf{w}_r - \check{\mathbf{w}}_r) & \text{if } f(u_m) = F \\ 0 & \text{if } f(u_m) < F \end{cases} \geq 0, & \Delta \check{\Upsilon}_n &= -(\check{v}_r - v_m) - \begin{cases} (\mathbf{w}_r - \check{\mathbf{w}}_r) & \text{if } f(u_m) = F \\ (\check{\mathbf{w}}_m - \check{\mathbf{w}}_r) & \text{if } f(u_m) < F \end{cases} < 0, \end{aligned}$$

and therefore

$$\Delta \# \in \{-1, 0, 1\}, \quad \Delta \mathcal{T}_n = -2(\hat{v}_m - v_\ell) - 2(v_r - v_m) - 2(\mathbf{w}_\ell - \mathbf{w}_m) - \begin{cases} 0 & \text{if } f(u_m) = F \\ 2(\check{\mathbf{w}}_m - \check{\mathbf{w}}_r) & \text{if } f(u_m) < F \end{cases} < 0.$$

CD-NS $\bar{\mathbf{F}}$ (u_ℓ, u_m) is a CD and (u_m, u_r) is a NS. In this case $v_\ell = v_m = \hat{v}_m < v_r$, $\mathbf{w}_r < \min\{\mathbf{w}_\ell, \mathbf{w}_m\}$, $\mathbf{w}_\ell = \hat{\mathbf{w}}_\ell$, $\mathbf{w}_m = \hat{\mathbf{w}}_m$, $f(u_m) = f(u_r) = F \neq f(u_\ell)$ and $u_\ell, u_m \in \Omega_c$.

If $f(u_\ell) < F$, then $v_r > \hat{v}_\ell > v_\ell = v_m$, $\mathbf{w}_r < \mathbf{w}_\ell < \mathbf{w}_m$ and $\mathcal{R}_{F,n}[u_\ell, u_r]$ has a fan of RSs from u_ℓ to \hat{u}_ℓ and a NS (\hat{u}_ℓ, u_r); as a consequence $\Delta \text{TV}_w = -2(\mathbf{w}_m - \mathbf{w}_\ell) < 0 = \Delta \text{TV}_v = \Delta \hat{\Upsilon}_n = \Delta \check{\Upsilon}_n$, therefore $\Delta \# \in [0, 2^n - 2]$ and $\Delta \mathcal{T}_n = -2(\mathbf{w}_m - \mathbf{w}_\ell) < 0$.

If $f(u_\ell) > F$, then $v_r > v_\ell = v_m = \hat{v}_m > \hat{v}_\ell$, $\mathbf{w}_\ell > \mathbf{w}_m > \mathbf{w}_r$ and $\mathcal{R}_{F,n}[u_\ell, u_r]$ has a two waves (u_ℓ, \hat{u}_ℓ) and (\hat{u}_ℓ, u_r), that are a S and a NS, respectively; as a consequence

$$\begin{aligned} \Delta \text{TV}_v &= 2(v_\ell - \hat{v}_\ell) > 0, & \Delta \hat{\Upsilon}_n &= -(v_\ell - \hat{v}_\ell) - (\mathbf{w}_\ell - \mathbf{w}_m) < 0, \\ \Delta \text{TV}_w &= 0, & \Delta \check{\Upsilon}_n &= 0, \end{aligned}$$

therefore $\Delta \# = 0$ and $\Delta \mathcal{T}_n = -2(\mathbf{w}_\ell - \mathbf{w}_m) < 0$.

CD-PT $\bar{\mathbf{F}}$ (u_ℓ, u_m) is a CD and (u_m, u_r) is a PT. In this case $v_\ell = v_m = \hat{v}_m = V > v_r = \check{v}_r > \hat{v}_\ell$, $\mathbf{w}_\ell = \hat{\mathbf{w}}_\ell > \check{\mathbf{w}}_r \geq \mathbf{w}_r \geq w^- > w_F = \hat{\mathbf{w}}_m \geq \mathbf{w}_m$, $f(u_\ell) > f(\mathbf{u}_*) > F \geq \max\{f(u_m), f(u_r)\}$, $u_\ell \in \Omega_c^+$, $u_m \in \Omega_c^-$ and $u_r \in \Omega_c$. $\mathcal{R}_{F,n}[u_\ell, u_r]$ has at most three waves (u_ℓ, \hat{u}_ℓ), ($\hat{u}_\ell, \check{u}_r$) and (\check{u}_r, u_r) that are a S, a NS and a possibly null CD, respectively. As a consequence

$$\begin{aligned} \Delta \text{TV}_v &= 2(v_r - \hat{v}_\ell) > 0, & \Delta \hat{\Upsilon}_n &= -(v_m - \hat{v}_\ell) - (\mathbf{w}_\ell - w_F) < 0, \\ \Delta \text{TV}_w &= -2(\mathbf{w}_r - \mathbf{w}_m) < 0, & \Delta \check{\Upsilon}_n &= 0, \end{aligned}$$

therefore $\Delta \# \in \{0, 1\}$ and $\Delta \mathcal{T}_n = -2(v_m - v_r) - 2(\mathbf{w}_r - \mathbf{w}_m) - 2(\mathbf{w}_\ell - w_F) < 0$.

NS-S $\bar{\mathbf{F}}$ (u_ℓ, u_m) is a NS and (u_m, u_r) is a S. In this case $v_m > v_r = \check{v}_r > v_\ell$, $\mathbf{w}_\ell > \check{\mathbf{w}}_r > \mathbf{w}_m = \mathbf{w}_r \geq w^-$, $f(u_\ell) = f(u_m) = F > f(u_r)$ and $u_\ell, u_m, u_r \in \Omega_c$. $\mathcal{R}_{F,n}[u_\ell, u_r]$ has two waves (u_ℓ, \check{u}_r) and (\check{u}_r, u_r) that are a NS and a CD, respectively. As a consequence $\Delta \text{TV}_v = -2(v_m - v_r) < 0 = \Delta \text{TV}_w = \Delta \hat{\Upsilon}_n = \Delta \check{\Upsilon}_n$, therefore $\Delta \# = 0$ and $\Delta \mathcal{T}_n = -2(v_m - v_r) < 0$.

NS-RS $\bar{\mathbf{F}}$ (u_ℓ, u_m) is a NS and (u_m, u_r) is a RS. In this case $v_\ell < v_m = \check{v}_m < v_r \leq \check{v}_r$, $\mathbf{w}_\ell > \mathbf{w}_m = \check{\mathbf{w}}_m = \mathbf{w}_r > \check{\mathbf{w}}_r$, $f(u_r) > F = f(u_\ell) = f(u_m)$ and $u_\ell, u_m, u_r \in \Omega_c$. $\mathcal{R}_{F,n}[u_\ell, u_r]$ has two waves (u_ℓ, \check{u}_r) and (\check{u}_r, u_r) that are a NS and either a PT or a CD, respectively. As a consequence

$$\begin{aligned} \Delta \text{TV}_v &= 2(\check{v}_r - v_r) \geq 0, & \Delta \hat{\Upsilon}_n &= 0, \\ \Delta \text{TV}_w &= 2(\mathbf{w}_r - \check{\mathbf{w}}_r) > 0, & \Delta \check{\Upsilon}_n &= -(\check{v}_r - v_m) - (\mathbf{w}_r - \check{\mathbf{w}}_r) < 0, \end{aligned}$$

therefore $\Delta \# = 0$ and $\Delta \mathcal{T}_n = -2(v_r - v_m) < 0$. Notice that $\check{v}_r > v_r$ if and only if $\mathbf{w}_m = \mathbf{w}_r = w^-$ and $\check{v}_r = V > v_r > v_m = v_F^+$.

NS-PT $\bar{\mathbf{F}}$ (u_ℓ, u_m) is a NS and (u_m, u_r) is a PT. In this case $v_m = V > v_r = \check{v}_r > v_\ell$, $\mathbf{w}_\ell > \check{\mathbf{w}}_r > \mathbf{w}_r = w^- > \mathbf{w}_m$, $f(\mathbf{u}_*) > f(u_\ell) = f(u_m) = F > f(u_r)$, $u_\ell, u_r \in \Omega_c$ and $u_m \in \Omega_c^-$. $\mathcal{R}_{F,n}[u_\ell, u_r]$ has two waves (u_ℓ, \check{u}_r) and (\check{u}_r, u_r) that are a NS and a CD, respectively. As a consequence

$$\begin{aligned} \Delta \text{TV}_v &= -2(V - v_r) < 0, & \Delta \hat{\Upsilon}_n &= 0, \\ \Delta \text{TV}_w &= -2(w^- - \mathbf{w}_m) < 0, & \Delta \check{\Upsilon}_n &= 0, \end{aligned}$$

therefore $\Delta \# = 0$ and $\Delta \mathcal{T}_n = -2(V - v_r) - 2(w^- - \mathbf{w}_m) < 0$.

This concludes the proof. \square

Interaction	Result	$\Delta\sharp$	$\Delta\mathcal{T}_n$
CD_F^+	(S,NS,CD), (S,NS)	$\in \{1, 2\}$	< 0
RS_F^+	(NS,CD)	$= 1$	< 0
CD_F^-	(S,NS,CD), (S,NS), (PT,NS,CD), (PT,NS)	$\in \{1, 2\}$	< 0
RS_F^-	(NS,PT), (NS,CD)	$= 1$	< 0
CD-S	(S,CD), (PT,CD), PT	≤ 0	$= 0$
CD-RS	(RS,CD)	$= 0$	$= 0$
CD-PT	(PT,CD), PT, (S,CD), S	≤ 0	≤ 0
S-S	S	< 0	$= 0$
S-RS	S	< 0	< 0
RS-S	S	< 0	< 0
PT-S	PT	< 0	$= 0$
PT-RS	PT, CD	< 0	< 0
CD-S ₀	(S,CD), (PT,CD), PT	≤ 0	≤ 0
CD-RS ₀	(RS,CD)	$= 0$	≤ 0
CD-NS ₀	(RSs,CD), RSs	$\in [-1, 2^n - 1]$	< 0
CD-PT ₀	(PT,CD), PT, (S,CD), S	≤ 0	≤ 0
S-S ₀	S	< 0	$= 0$
S-RS ₀	S	< 0	< 0
RS-S ₀	S	< 0	< 0
NS-S ₀	(S,CD), CD	≤ 0	< 0
NS-PT ₀	(S,CD), S, CD	≤ 0	< 0
PT-S ₀	PT	< 0	$= 0$
PT-RS ₀	PT, CD	< 0	< 0
PT-NS ₀	CD, PT	< 0	< 0
$CD-S_F^+$	(S,NS,CD)	$= 1$	< 0
$CD-RS_F^+$	(S,NS,CD), (NS,CD), (S,NS), NS	$\in \{-1, 0, 1\}$	< 0
$CD-NS_F^+$	(RSs,NS), (S,NS)	$\in [0, 2^n - 2]$	< 0
$CD-PT_F^+$	(S,NS,CD)	$= 1$	< 0
$NS-S_F^+$	(NS,CD)	$= 0$	< 0
$NS-RS_F^+$	(NS,CD)	$= 0$	< 0
$CD-S_F^-$	(S,NS,CD)	$= 1$	< 0
$CD-RS_F^-$	(S,NS,CD), (S,NS,PT), (NS,CD), (NS,PT), (S,NS), NS	$\in \{-1, 0, 1\}$	< 0
$CD-NS_F^-$	(RSs,NS), (S,NS)	$\in [0, 2^n - 2]$	< 0
$CD-PT_F^-$	(S,NS,CD), (S,NS)	$\in \{0, 1\}$	< 0
$NS-S_F^-$	(NS,CD)	$= 0$	< 0
$NS-RS_F^-$	(NS,PT), (NS,CD)	$= 0$	< 0
$NS-PT_F^-$	(NS,CD)	$= 0$	< 0

Table 1: An overview of the interactions considered in the proof of Proposition 4.1.

In Table 1 we collect the most relevant possible interactions considered in the proof of Proposition 4.1 and list the corresponding possible results in terms of wave types, $\Delta\sharp$ and $\Delta\mathcal{T}_n$.

Beside the bound on the number of wave-fronts proved in Proposition 4.1, we need to bound also the number of interactions. This is the aim of the next proposition, which together with Proposition 4.1 ensure the global existence of u_n . We underline that for any interaction $\Delta\sharp \leq 2^n - 1$, see Table 1.

Proposition 4.2. *For any fixed $n \in \mathbb{N}$ sufficiently large and $u_n^o \in \mathbf{PC}(\mathbb{R}; \mathcal{G}_n)$, we have that the number of interactions in $(0, \infty)$ is bounded. In particular u_n is globally defined.*

Proof. From what we already show in the proof of Proposition 4.1, see Table 1, we deduce that

$$t \mapsto 2^n \frac{\mathcal{T}_n(t)}{\varepsilon_n} + \sharp(t)$$

strictly decreases after any interaction, except the following cases.

A CD (u_ℓ, u_m) interacts with (u_m, u_r) and one of the following conditions is satisfied:

- (u_m, u_r) is a S and $\mathfrak{w}_\ell = w^- - 1$;
 - (u_m, u_r) is a S and $w^- - 1 < \mathfrak{w}_\ell \leq w^- = \mathfrak{w}_r$;
 - (u_m, u_r) is a RS;
 - (u_m, u_r) is a PT and $\mathfrak{w}_\ell > w^-$.
- (17)

For this reason it remains to bound the number of only the above type of interactions. We observe that the number of waves of u_n do not change after interactions as in (17). This implies that the number of waves is uniformly bounded. We also observe that any interaction as in (17) has exactly one incoming CD and exactly one outgoing CD. Since no wave can reach any CD from the left (and then possibly have with it an interaction as in (17)), we have that as long as a CD remains a CD (possible further interactions involving it have to be taken into account), it can interact only once with another wave W (or with waves generated by further interactions involving W), moreover in this case W is slower than such CD and is not another CD. Since furthermore we already know that the number of waves is uniformly bounded, there can be only finitely many interactions involving CDs. It is therefore now clear that also the number of the interactions described in (17) is bounded. \square

4.3 Convergence

We first observe that

$$|\rho_\ell - \rho_r| \leq L (|v_\ell - v_r| + |\mathfrak{w}_\ell - \mathfrak{w}_r|)$$

where $L \doteq \max\{\rho^-, \|1/p'\|_{\mathbf{L}^\infty([p^{-1}(w^-), p^{-1}(w^+)]; \mathbb{R})}\}$ because

$$\rho_{\ell,r} = \begin{cases} p^{-1}(\mathfrak{w}_{\ell,r} - v_{\ell,r}) & \text{if } \mathfrak{w}_{\ell,r} \in [w^-, w^+], \\ (\mathfrak{w}_{\ell,r} + 1 - w^-) \rho^- & \text{if } \mathfrak{w}_{\ell,r} \in [w^- - 1, w^-]. \end{cases}$$

As a consequence $\text{TV}(\rho) \leq L(\text{TV}(v) + \text{TV}(\mathfrak{w}))$, hence

$$\text{TV}(u) \leq (1 + L) (\text{TV}(v) + \text{TV}(\mathfrak{w})).$$

Moreover, by Proposition 4.1 and (16) we have that for any $t > 0$

$$\text{TV}(v_n(t, \cdot)) + \text{TV}(\mathfrak{w}_n(t, \cdot)) \leq \mathcal{T}_n(t) \leq \mathcal{T}_n(0) \leq \text{TV}(v^o) + \text{TV}(\mathfrak{w}^o) + 2C \left(\hat{\Upsilon}(u^o) + \check{\Upsilon}(u^o) \right).$$

As a consequence $\text{TV}(u_n)$ is bounded by

$$C_F^o \doteq (1 + L) \left[\text{TV}(v^o) + \text{TV}(\mathfrak{w}^o) + 2C \left(\hat{\Upsilon}(u^o) + \check{\Upsilon}(u^o) \right) \right]. \quad (18)$$

Since u_n^o takes values in Ω , for any $t > 0$ we have that also $u_n(t, \cdot)$ takes values in Ω , hence

$$\|u_n(t, \cdot)\|_{\mathbf{L}^\infty(\mathbb{R}; \Omega)} \leq R + V.$$

Moreover

$$\|u_n(t, \cdot) - u_n(s, \cdot)\|_{\mathbf{L}^1(\mathbb{R}; \Omega)} \leq L_F^o |t - s|, \quad (19)$$

with $L_F^o \doteq C_F^o \max\{V, Rp'(R)\}$. Indeed, if no interaction occurs for times between t and s , then

$$\begin{aligned} \|u_n(t, \cdot) - u_n(s, \cdot)\|_{\mathbf{L}^1(\mathbb{R}; \Omega)} &\leq \sum_{i \in \mathbf{D}(t)} \left| (t - s) \dot{\delta}_n^i(t) \left(\rho_n(t, \delta_n^i(t)_-) - \rho_n(t, \delta_n^i(t)_+) \right) \right| \\ &\quad + \sum_{i \in \mathbf{D}(t)} \left| (t - s) \dot{\delta}_n^i(t) \left(v_n(t, \delta_n^i(t)_-) - v_n(t, \delta_n^i(t)_+) \right) \right| \leq L_F^o |t - s|, \end{aligned}$$

where $\delta_n^i(t) \in \mathbb{R}$, $i \in \mathbf{D}(t) \subset \mathbb{N}$, are the positions of the discontinuities of $u_n(t, \cdot)$. The case when one or more interactions take place for times between t and s is similar, because by the finite speed of propagation of the waves the map $t \mapsto u_n(t, \cdot)$ is \mathbf{L}^1 -continuous across interaction times.

Thus, by applying Helly's Theorem, the approximate solutions $(u_n)_n$ converge (up to a subsequence) in $\mathbf{L}_{\text{loc}}^1(\mathbb{R}_+ \times \mathbb{R}; \Omega)$ to a function $u \in \mathbf{L}^\infty(\mathbb{R}_+; \mathbf{BV}(\mathbb{R}; \Omega)) \cap \mathbf{C}^0(\mathbb{R}_+; \mathbf{L}_{\text{loc}}^1(\mathbb{R}; \Omega))$ and the limit satisfies the estimates in (13).

Proposition 4.3. *Let $u^o \in \mathbf{L}^1 \cap \mathbf{BV}(\mathbb{R}; \Omega)$ and $F \in [0, f_c^+]$ satisfy (H.1) or (H.2). If u is a limit of the approximate solutions $(u_n)_n$ constructed in Section 4.1, then u is a solution to constrained Cauchy problem (3), (4), (5) in the sense of Definition 2.2.*

Proof. We consider separately the conditions listed in Definition 2.2.

(CS.1) The initial condition (4) holds by (13), (19) and the $\mathbf{L}_{\text{loc}}^1$ -convergence of u_n to u .

(CS.2) We prove now (6), that is for any test function $\phi \in \mathbf{C}_c^\infty((0, \infty) \times \mathbb{R}; \mathbb{R})$ we have

$$\int_0^\infty \int_{\mathbb{R}} (\rho \phi_t + f(u) \phi_x) dx dt = 0.$$

Choose $T > 0$ such that $\phi(t, x) = 0$ whenever $t \geq T$. Since u_n is uniformly bounded and f is uniformly continuous on bounded sets, it is sufficient to prove that

$$\int_0^T \int_{\mathbb{R}} (\rho_n \phi_t + f(u_n) \phi_x) dx dt \rightarrow 0. \quad (20)$$

By the Green-Gauss formula the double integral above can be written as

$$\int_0^T \sum_{i \in \mathbf{D}(t)} \left(\dot{\delta}_n^i(t) \Delta \rho_n^i(t) - \Delta f_n^i(t) \right) \phi(t, \delta_n^i(t)) dt,$$

where

$$\Delta \rho_n^i(t) \doteq \rho_n(t, \delta_n^i(t)_+) - \rho_n(t, \delta_n^i(t)_-), \quad \Delta f_n^i(t) \doteq f(u_n(t, \delta_n^i(t)_+)) - f(u_n(t, \delta_n^i(t)_-)).$$

By construction any discontinuity of $u_n(t, \cdot)$ satisfies the first Rankine-Hugoniot condition (9), therefore

$$\dot{\delta}_n^i(t) \Delta \rho_n^i(t) - \Delta f_n^i(t) = 0, \quad i \in \mathbf{D}(t),$$

and (20) is trivial.

The proof of (7) is analogous because by construction any discontinuity of $u_n(t, \cdot)$ away from $x = 0$ satisfies also the second Rankine-Hugoniot condition (10).

(CS.3) We prove now (8), namely that for any $k \in [0, V]$ and $\phi \in \mathbf{C}_c^\infty((0, \infty) \times \mathbb{R}; \mathbb{R})$ such that $\phi(\cdot, 0) \equiv 0$ and $\phi \geq 0$ we have

$$\int_0^\infty \int_{\mathbb{R}} (\mathbf{E}^k(u) \phi_t + \mathbf{Q}^k(u) \phi_x) dx dt \geq 0,$$

where

$$\mathbf{E}^k(u) \doteq \begin{cases} 0 & \text{if } v \geq k, \\ \frac{\rho}{p^{-1}(\mathbb{W}(u) - k)} - 1 & \text{if } v < k, \end{cases} \quad \mathbf{Q}^k(u) \doteq \begin{cases} 0 & \text{if } v \geq k, \\ \frac{f(u)}{p^{-1}(\mathbb{W}(u) - k)} - k & \text{if } v < k. \end{cases}$$

Choose $T > 0$ such that $\phi(t, x) = 0$ whenever $t \geq T$. By the a.e. convergence of u_n to u and the uniform continuity of \mathbf{E}^k and \mathbf{Q}^k , it is sufficient to prove that

$$\liminf_{n \rightarrow \infty} \int_0^T \int_{\mathbb{R}} \left(\mathbf{E}^k(u_n) \phi_t + \mathbf{Q}^k(u_n) \phi_x \right) dx dt \geq 0. \quad (21)$$

By the Green-Gauss formula the double integral above can be written as

$$\int_0^T \sum_{i \in \mathbf{D}(t)} \left(\dot{\delta}_n^i(t) \Delta \mathbf{E}_n^{k,i}(t) - \Delta \mathbf{Q}_n^{k,i}(t) \right) \phi(t, \delta_n^i(t)) dt,$$

where

$$\Delta \mathbf{E}_n^{k,i}(t) \doteq \mathbf{E}^k \left(u_n(t, \delta_n^i(t)_+) \right) - \mathbf{E}^k \left(u_n(t, \delta_n^i(t)_-) \right), \quad \Delta \mathbf{Q}_n^{k,i}(t) \doteq \mathbf{Q}^k \left(u_n(t, \delta_n^i(t)_+) \right) - \mathbf{Q}^k \left(u_n(t, \delta_n^i(t)_-) \right).$$

To estimate the above integral we have to distinguish the following cases.

- If the i th discontinuity is a PT, then we let $x \doteq \delta_n^i(t)$ and observe that

$$\begin{aligned} \rho_n(t, x_-) &< \min \{ \rho_n(t, x_+), p^{-1}(w^- - k) \}, & \dot{\delta}_n^i(t) &= \Lambda(u_n(t, x_-), u_n(t, x_+)), \\ v_n(t, x_-) &= V > v_n(t, x_+), & \mathbf{W}(u_n(t, x_-)) &= w^- \leq \mathbf{w}(u_n(t, x_+)) = \mathbf{W}(u_n(t, x_+)), \end{aligned}$$

hence

$$\begin{aligned} \Delta \mathbf{E}_n^{k,i}(t) &= \begin{cases} \frac{\rho_n(t, x_+)}{\rho_{n,+}^k} - 1 & \text{if } v_n(t, x_+) < k \leq V, \\ 0 & \text{if } k \leq v_n(t, x_+), \end{cases} \\ -\Delta \mathbf{Q}_n^{k,i}(t) &= \begin{cases} k - \frac{f(u_n(t, x_+))}{\rho_{n,+}^k} & \text{if } v_n(t, x_+) < k \leq V, \\ 0 & \text{if } k \leq v_n(t, x_+), \end{cases} \end{aligned}$$

where $\rho_{n,+}^k \doteq p^{-1}(\mathbf{w}(u_n(t, x_+)) - k)$. If $v_n(t, x_+) < k \leq V$, then

$$\begin{aligned} &\dot{\delta}_n^i(t) \Delta \mathbf{E}_n^{k,i}(t) - \Delta \mathbf{Q}_n^{k,i}(t) \\ &= \Lambda(u_n(t, x_-), u_n(t, x_+)) \left[\frac{\rho_n(t, x_+)}{\rho_{n,+}^k} - 1 \right] + k - \frac{f(u_n(t, x_+))}{\rho_{n,+}^k} \\ &= \underbrace{\left[\frac{\rho_n(t, x_+)}{\rho_{n,+}^k} - 1 \right]}_{>0} \underbrace{\left[\Lambda(u_n(t, x_-), u_n(t, x_+)) - \Lambda((\rho_{n,+}^k, k), u_n(t, x_+)) \right]}_{>0} > 0. \end{aligned}$$

- If the i th discontinuity is a CD, then we let $x \doteq \delta_n^i(t)$ and observe that $\dot{\delta}_n^i(t) = v_n(t, x_-) = v_n(t, x_+)$ implies that $\dot{\delta}_n^i(t) \Delta \mathbf{E}_n^{k,i}(t) - \Delta \mathbf{Q}_n^{k,i}(t) = 0$.

- If the i th discontinuity is a S, then we let $x \doteq \delta_n^i(t)$ and observe that

$$\begin{aligned} \rho_n(t, x_-) &< \rho_n(t, x_+), & f(u_n(t, x_-)) &> f(u_n(t, x_+)), & \dot{\delta}_n^i(t) &= \Lambda(u_n(t, x_-), u_n(t, x_+)) < 0, \\ v_n(t, x_-) &> v_n(t, x_+), & \mathbf{w}_{\pm} &\doteq \mathbf{w}(u_n(t, x_-)) = \mathbf{w}(u_n(t, x_+)) \geq w^-, \end{aligned}$$

hence

$$\begin{aligned} \Delta \mathbf{E}_n^{k,i}(t) &= \begin{cases} \frac{\rho_n(t, x_+) - \rho_n(t, x_-)}{p^{-1}(\mathbf{w}_{\pm} - k)} & \text{if } v_n(t, x_+) < v_n(t, x_-) < k, \\ \frac{\rho_n(t, x_+)}{p^{-1}(\mathbf{w}_{\pm} - k)} - 1 & \text{if } v_n(t, x_+) < k \leq v_n(t, x_-), \\ 0 & \text{if } k \leq v_n(t, x_+) < v_n(t, x_-), \end{cases} \\ -\Delta \mathbf{Q}_n^{k,i}(t) &= \begin{cases} \frac{f(u_n(t, x_-)) - f(u_n(t, x_+))}{p^{-1}(\mathbf{w}_{\pm} - k)} & \text{if } v_n(t, x_+) < v_n(t, x_-) < k, \\ k - \frac{f(u_n(t, x_+))}{p^{-1}(\mathbf{w}_{\pm} - k)} & \text{if } v_n(t, x_+) < k \leq v_n(t, x_-), \\ 0 & \text{if } k \leq v_n(t, x_+) < v_n(t, x_-). \end{cases} \end{aligned}$$

If $k > v_n(t, x_-)$ or $k \leq v_n(t, x_+)$, then it is immediate to see that $\delta_n^i(t) \Delta \mathbf{E}_n^{k,i}(t) - \Delta \mathbf{Q}_n^{k,i}(t) = 0$. Furthermore, if $v_n(t, x_+) < k \leq v_n(t, x_-)$, then

$$\begin{aligned} & \delta_n^i(t) \Delta \mathbf{E}_n^{k,i}(t) - \Delta \mathbf{Q}_n^{k,i}(t) \\ &= \Lambda(u_n(t, x_-), u_n(t, x_+)) \left[\frac{\rho_n(t, x_+)}{p^{-1}(\mathbf{w}_\pm - k)} - 1 \right] + k - \frac{f(u_n(t, x_+))}{p^{-1}(\mathbf{w}_\pm - k)} \\ &= \underbrace{\left[\frac{\rho_n(t, x_+)}{p^{-1}(\mathbf{w}_\pm - k)} - 1 \right]}_{>0} \underbrace{\left[\Lambda(u_n(t, x_-), u_n(t, x_+)) - \Lambda\left(p^{-1}(\mathbf{w}_\pm - k), k\right), u_n(t, x_+)\right]}_{>0} > 0. \end{aligned}$$

• If the i th discontinuity is a RS, then we let $x \doteq \delta_n^i(t)$ and observe that

$$\begin{aligned} \rho_n(t, x_-) &> \rho_n(t, x_+), & f(u_n(t, x_-)) &< f(u_n(t, x_+)), & \delta_n^i(t) &= \Lambda(u_n(t, x_-), u_n(t, x_+)) < 0, \\ v_n(t, x_-) &< v_n(t, x_+), & \mathbf{w}_\pm &\doteq \mathbf{w}(u_n(t, x_-)) = \mathbf{w}(u_n(t, x_+)) \geq w^-, \end{aligned}$$

hence

$$\begin{aligned} \Delta \mathbf{E}_n^{k,i}(t) &= \begin{cases} \frac{\rho_n(t, x_+) - \rho_n(t, x_-)}{p^{-1}(\mathbf{w}_\pm - k)} & \text{if } v_n(t, x_-) < v_n(t, x_+) < k, \\ \frac{\rho_n(t, x_-)}{p^{-1}(\mathbf{w}_\pm - k)} - 1 & \text{if } v_n(t, x_-) < k \leq v_n(t, x_+), \\ 0 & \text{if } k \leq v_n(t, x_-) < v_n(t, x_+), \end{cases} \\ -\Delta \mathbf{Q}_n^{k,i}(t) &= \begin{cases} \frac{f(u_n(t, x_-)) - f(u_n(t, x_+))}{p^{-1}(\mathbf{w}_\pm - k)} & \text{if } v_n(t, x_-) < v_n(t, x_+) < k, \\ \frac{f(u_n(t, x_-))}{p^{-1}(\mathbf{w}_\pm - k)} - k & \text{if } v_n(t, x_-) < k \leq v_n(t, x_+), \\ 0 & \text{if } k \leq v_n(t, x_-) < v_n(t, x_+). \end{cases} \end{aligned}$$

If $k > v_n(t, x_+)$ or $k \leq v_n(t, x_-)$, then it is immediate to see that $\delta_n^i(t) \Delta \mathbf{E}_n^{k,i}(t) - \Delta \mathbf{Q}_n^{k,i}(t) = 0$. Furthermore, if $v_n(t, x_-) < k \leq v_n(t, x_+)$, then

$$\begin{aligned} & \delta_n^i(t) \Delta \mathbf{E}_n^{k,i}(t) - \Delta \mathbf{Q}_n^{k,i}(t) \\ &= \Lambda(u_n(t, x_-), u_n(t, x_+)) \left[\frac{\rho_n(t, x_-)}{p^{-1}(\mathbf{w}_\pm - k)} - 1 \right] + \frac{f(u_n(t, x_-))}{p^{-1}(\mathbf{w}_\pm - k)} - k \\ &= \underbrace{\left[\frac{\rho_n(t, x_-)}{p^{-1}(\mathbf{w}_\pm - k)} - 1 \right]}_{>0} \underbrace{\left[\Lambda(u_n(t, x_-), u_n(t, x_+)) + \Lambda\left(u_n(t, x_-), p^{-1}(\mathbf{w}_\pm - k), k\right) \right]}_{<0} \\ &\geq -\frac{2}{\rho^-} p^{-1}(\mathbf{w}_\pm) p'(p^{-1}(\mathbf{w}_\pm)) [\rho_n(t, x_-) - \rho_n(t, x_+)] \end{aligned}$$

because $\rho_n(t, x_-) > p^{-1}(\mathbf{w}_\pm - k) \geq \rho_n(t, x_+) \geq \rho^-$ and because by the concavity of $\mathfrak{L}_{\mathbf{w}_\pm}(\rho) = (\mathbf{w}_\pm - p(\rho)) \rho$ we have

$$\begin{aligned} 0 &> \Lambda(u_n(t, x_-), u_n(t, x_+)) > \Lambda\left(u_n(t, x_-), p^{-1}(\mathbf{w}_\pm - k), k\right) \\ &> \mathfrak{L}'_{\mathbf{w}_\pm}(\rho_n(t, x_-)) = \mathbf{w}_\pm - p(\rho_n(t, x_-)) - \rho_n(t, x_-) p'(\rho_n(t, x_-)) \\ &\geq \mathfrak{L}'_{\mathbf{w}_\pm}(p^{-1}(\mathbf{w}_\pm)) = -p^{-1}(\mathbf{w}_\pm) p'(p^{-1}(\mathbf{w}_\pm)). \end{aligned}$$

The above case by case study shows that

$$\begin{aligned} & \liminf_{n \rightarrow \infty} \int_0^T \int_{\mathbb{R}} \left[\mathbf{E}^k(u_n) \phi_t + \mathbf{Q}^k(u_n) \phi_x \right] dx dt \\ &= \liminf_{n \rightarrow \infty} \int_0^T \sum_{i \in \text{RS}_n(t)} \left[\delta_n^i(t) \Delta \mathbf{E}_n^{k,i}(t) - \Delta \mathbf{Q}_n^{k,i}(t) \right] \phi(t, \delta_n^i(t)) dt \end{aligned}$$

$$\begin{aligned}
&\geq -\frac{2}{\rho^-} \max_{\rho \in [p^{-1}(w^-), R]} |\rho p'(\rho)| \liminf_{n \rightarrow \infty} \int_0^T \sum_{i \in \text{RS}_n(t)} \left[\rho_n(t, \delta_n^i(t)_-) - \rho_n(t, \delta_n^i(t)_+) \right] \phi(t, \delta_n^i(t)) dt \\
&\geq -\frac{2T}{\rho^-} \|\phi\|_{\mathbf{L}^\infty} C_F^o \max_{\rho \in [\rho^-, R]} |\rho p'(\rho)| \doteq -M,
\end{aligned}$$

where $\delta_n^i(t) \in \mathbb{R}$, $i \in \text{RS}_n(t) \subset \mathbb{N}$, are the positions of the RSs of $u_n(t, \cdot)$ and C_F^o is defined in (18).

We claim that for any fixed $h > 0$, there exists a dense set \mathcal{K}_h of values of k in $[0, V]$ such that

$$\liminf_{n \rightarrow \infty} \int_0^T \sum_{i \in \text{RS}_n(t)} \left[\dot{\delta}_n^i(t) \Delta \mathbf{E}_n^{k,i}(t) - \Delta \mathbf{Q}_n^{k,i}(t) \right] \phi(t, \delta_n^i(t)) dt \geq -\frac{1}{h}.$$

To prove it we fix $a, b \in [0, V]$ with $a < b$ and show that there exists $k \in (a, b)$ such that the above estimate is satisfied. Let $l \doteq \lceil 2(Mh + 1)/(b - a) \rceil$ and introduce the set

$$\mathcal{K}_h \doteq \frac{2\mathbb{N} + 1}{l} \cap (a, b).$$

Let $\mathcal{E}_n > 0$ be the maximal (v, w) -distance between two ‘‘consecutive’’ points in the grid \mathcal{G}_n having the same w -coordinate, namely, with a slight abuse of notations, we let

$$\mathcal{E}_n \doteq \max_{\substack{(v^i, w), (v^{i+1}, w) \in \mathcal{G}_n \\ v^i \neq v^{i+1}}} (v^{i+1} - v^i).$$

Let $n_h \in \mathbb{N}$ be sufficiently large so that $\mathcal{E}_{n_h} < 2/l$. Take $n \geq n_h$. We claim that for any $i \in \text{RS}_n(t)$ we have

$$\mathcal{K}_h \cap \left(v_n(t, \delta_n^i(t)_-), v_n(t, \delta_n^i(t)_+) \right)$$

has at most one element. Indeed, if \mathcal{K}_h has more than one element then for any $i \in \text{RS}_n(t)$ we have

$$v_n(t, \delta_n^i(t)_+) - v_n(t, \delta_n^i(t)_-) \leq \mathcal{E}_n < \frac{2}{l} = \min_{\substack{k^1, k^2 \in \mathcal{K}_h \\ k^1 \neq k^2}} |k^1 - k^2|.$$

As a consequence the sum

$$\sum_{k \in \mathcal{K}_h} \left[\dot{\delta}_n^i(t) \Delta \mathbf{E}_n^{k,i}(t) - \Delta \mathbf{Q}_n^{k,i}(t) \right]$$

has at most one nonzero element; moreover

$$-m \left(\rho_n(t, \delta_n^i(t)_-) - \rho_n(t, \delta_n^i(t)_+) \right) \leq \sum_{k \in \mathcal{K}_h} \left[\dot{\delta}_n^i(t) \Delta \mathbf{E}_n^{k,i}(t) - \Delta \mathbf{Q}_n^{k,i}(t) \right],$$

where

$$m \doteq \frac{2}{\rho^-} \max_{\rho \in [\rho^-, R]} |\rho p'(\rho)| = \frac{M}{T C_F^o \|\phi\|_{\mathbf{L}^\infty}}.$$

Therefore we find

$$\sum_{i \in \text{RS}_n(t)} \sum_{k \in \mathcal{K}_h} \left[\dot{\delta}_n^i(t) \Delta \mathbf{E}_n^{k,i}(t) - \Delta \mathbf{Q}_n^{k,i}(t) \right] \geq -m C_F^o.$$

By exchanging the sums, multiplying by the test function and integrating in time we get

$$\sum_{k \in \mathcal{K}_h} \int_0^T \sum_{i \in \text{RS}_n(t)} \left[\dot{\delta}_n^i(t) \Delta \mathbf{E}_n^{k,i}(t) - \Delta \mathbf{Q}_n^{k,i}(t) \right] \phi(t, \delta_n^i(t)) dt \geq -M.$$

Moreover, by construction we have that \mathcal{K}_h is a non-empty set with a finite number of elements (it has at most hM elements), hence

$$hM \max_{k \in \mathcal{K}_h} \left[\int_0^T \sum_{i \in \text{RS}_n(t)} \left[\dot{\delta}_n^i(t) \Delta \mathbf{E}_n^{k,i}(t) - \Delta \mathbf{Q}_n^{k,i}(t) \right] \phi(t, \delta_n^i(t)) dt \right] \geq -M.$$

In conclusion we proved that there exists $k \in \mathcal{K}_h \subseteq (a, b)$ such that the above estimate is satisfied for any $n \geq n_h$; therefore, since \mathcal{K}_h has a finite number of elements, we have

$$\liminf_{n \rightarrow \infty} \int_0^T \sum_{i \in \text{RS}_n(t)} \left[\dot{\delta}_n^i(t) \Delta \mathbf{E}_n^{k,i}(t) - \Delta \mathbf{Q}_n^{k,i}(t) \right] \phi(t, \delta_n^i(t)) dt \geq -\frac{1}{h}.$$

Since a and b are arbitrary, the above estimate holds true for a dense set of values of k in $[0, V]$.

Actually, the above estimate holds for any k in $[0, V]$ because the term in brackets in the above formula is continuous with respect to k . Finally, for the arbitrariness of h , we have that

$$\liminf_{n \rightarrow \infty} \int_0^T \sum_{i \in \text{RS}_n(t)} \left[\dot{\delta}_n^i(t) \Delta \mathbf{E}_n^{k,i}(t) - \Delta \mathbf{Q}_n^{k,i}(t) \right] \phi(t, \delta_n^i(t)) dt \geq 0$$

and this concludes the proof of (21).

(CS.4) We prove now that (5) holds for a.e. $t > 0$, namely

$$f(u(t, 0_{\pm})) \leq F \quad \text{for a.e. } t > 0.$$

By construction $f(u_n(t, 0_{\pm})) \leq F$ for any $t > 0$, namely the approximate solutions satisfy (5). Since weak convergence preserves pointwise inequalities, it is sufficient to prove that $f(u_n(t, 0_{\pm}))$ weakly converges to $f(u(t, 0_{\pm}))$. If ϕ is a smooth test function of time with compact support in $(0, \infty)$ and φ is a smooth test function of space with compact support and such that $\varphi(0) = 1$, then

$$\int_0^\infty f(u_n(t, 0_-)) \phi(t) dt = \int_0^\infty \int_{-\infty}^0 \left[\rho_n(t, x) \dot{\phi}(t) \varphi(x) + f(u_n(t, x)) \phi(t) \dot{\varphi}(x) \right] dx dt.$$

The right-hand side passes to the limit, yielding the analogous expression with u_n replaced by u . By using again the Green-Gauss formula, one finally finds that

$$\lim_{n \rightarrow \infty} \int_0^\infty f(u_n(t, 0_-)) \phi(t) dt = \int_0^\infty f(u(t, 0_-)) \phi(t) dt.$$

As a consequence $f(u_n(t, 0_-))$ weakly converges to $f(u(t, 0_-))$, hence $f(u(t, 0_-)) \leq F$ for a.e. $t > 0$. At last, since we already proved that u satisfies the first Rankine-Hugoniot condition, we have $f(u(t, 0_-)) = f(u(t, 0_+))$, hence $f(u(t, 0_{\pm})) \leq F$ for a.e. $t > 0$. \square

4.4 The density flow through $x = 0$

Let u be the solution of constrained Cauchy problem (3), (4), (5) constructed in the previous section. By Propositions 2.1 and 4.3 we have that non-classical shocks of u can occur only at the constraint location $x = 0$, and in this case the (density) flow at $x = 0$ does not exceed the maximal flow F allowed by the constraint.

In the case of a constrained Riemann problem (3), (5), (15), we know that u coincides with $(t, x) \mapsto \mathcal{R}_F[u_\ell, u_r](x/t)$, moreover if $(u_\ell, u_r) \in \mathcal{D}_2$ then the flow of the non-classical shock of u coincides with F . In the next proposition we show that also for a general constrained Cauchy problem the flow of the non-classical shocks of u coincides with F if the traces at $x = 0$ of the approximate solutions $(u_n)_n$ satisfy a technical condition.

Proposition 4.4. *Let $u^\circ \in \mathbf{L}^1 \cap \mathbf{BV}(\mathbb{R}; \Omega)$, $F \in [0, f_c^+]$ satisfy **(H.1)** or **(H.2)** and u be a limit of the approximate solutions $(u_n)_n$ constructed in Section 4.1. Assume that the traces at $x = 0$ of $(u_n)_n$ and u satisfy (14), that is for any $k \in [0, V]$ and $\phi \in \mathbf{C}_c^\infty((0, \infty) \times \mathbb{R}; \mathbb{R})$ such that $\phi \geq 0$*

$$\lim_{n \rightarrow \infty} \int_0^T \mathbf{N}_F^k(u_n(t, 0_-)) \phi(t, 0) dt = \int_0^T \mathbf{N}_F^k(u(t, 0_-)) \phi(t, 0) dt,$$

with

$$\mathbf{N}_F^k(u) \doteq \begin{cases} f(u) \left[\frac{k}{F} - \frac{1}{p^{-1}(W(u) - k)} \right]_+ & \text{if } F \neq 0, \\ k & \text{if } F = 0. \end{cases}$$

If at time $t_0 > 0$ the limit u has a non-classical discontinuity, then $f(u(t_0, 0_{\pm})) = F$.

Proof. We first prove that for any $k \in [0, V]$ and $\phi \in \mathbf{C}_c^\infty((0, \infty) \times \mathbb{R}; \mathbb{R})$ such that $\phi \geq 0$ we have

$$\int_0^\infty \left[\int_{\mathbb{R}} \left[\mathbf{E}^k(u) \phi_t + \mathbf{Q}^k(u) \phi_x \right] dx + \mathbf{N}_F^k(u(t, 0_-)) \phi(t, 0) \right] dt \geq 0. \quad (22)$$

Notice that (22) differs from (8) not only for an extra term involving $\mathbf{N}_F^k(u(t, 0_+))$, but also because here we do not require that $\phi(\cdot, 0) \equiv 0$.

Choose $T > 0$ such that $\phi(t, x) = 0$ whenever $t \geq T$. By (14), the a.e. convergence of u_n to u and the uniform continuity of \mathbf{E}^k and \mathbf{Q}^k , it is sufficient to prove that

$$\liminf_{n \rightarrow \infty} \int_0^T \left[\int_{\mathbb{R}} \left[\mathbf{E}^k(u_n) \phi_t + \mathbf{Q}^k(u_n) \phi_x \right] dx + \mathbf{N}_F^k(u_n(t, 0_-)) \phi(t, 0) \right] dt \geq 0. \quad (23)$$

As already observed in the proof of Proposition 4.3, by the Green-Gauss formula the double integral above can be written as

$$\int_0^T \sum_{i \in \mathbf{D}(t)} \left[\dot{\delta}_n^i(t) \Delta \mathbf{E}_n^{k,i}(t) - \Delta \mathbf{Q}_n^{k,i}(t) \right] \phi(t, \delta_n^i(t)) dt,$$

where

$$\Delta \mathbf{E}_n^{k,i}(t) \doteq \mathbf{E}^k(u_n(t, \delta_n^i(t)_+)) - \mathbf{E}^k(u_n(t, \delta_n^i(t)_-)), \quad \Delta \mathbf{Q}_n^{k,i}(t) \doteq \mathbf{Q}^k(u_n(t, \delta_n^i(t)_+)) - \mathbf{Q}^k(u_n(t, \delta_n^i(t)_-)).$$

To estimate the above integral we can proceed as in the proof of Proposition 4.3, with the exception that here the i th discontinuity could also be a NS. In this case, that is, if the i th discontinuity is a NS, then




Figure 9: Above $F \in (f_c^-, f_c^+)$, $v_0^\pm \doteq v_n(t, 0_\pm)$ and $v_{0,F}^\pm \doteq F/p^{-1}(\mathbf{W}(u_n(t, 0_\pm)) - k)$. With the first two pictures we show that if $v_0^- < k < v_0^+$, then $v_{0,F}^- < k$. In the last picture we consider the case $v_0^- < v_0^+ < k$ and show that $v_{0,F}^- < v_{0,F}^+ < k$.

$$\begin{aligned} \delta_n^i(t) &= 0, & f(u_n(t, 0_\pm)) &= F, & v_{0,F}^- &\leq v_n(t, 0_-) < v_n(t, 0_+), \\ \dot{\delta}_n^i(t) &= 0, & & & \mathbf{w}(u_n(t, 0_-)) &= \mathbf{W}(u_n(t, 0_-)) \geq \mathbf{W}(u_n(t, 0_+)), \end{aligned}$$

hence

$$\begin{aligned} -\Delta \mathbf{Q}_n^{k,i}(t) &= \begin{cases} \frac{F}{p^{-1}(\mathbf{w}(u_n(t, 0_-)) - k)} - \frac{F}{p^{-1}(\mathbf{W}(u_n(t, 0_+)) - k)} & \text{if } v_n(t, 0_-) < v_n(t, 0_+) < k, \\ \frac{F}{p^{-1}(\mathbf{w}(u_n(t, 0_-)) - k)} - k & \text{if } v_n(t, 0_-) < k \leq v_n(t, 0_+), \\ 0 & \text{if } k \leq v_n(t, 0_-) < v_n(t, 0_+), \end{cases} \\ \mathbf{N}_F^k(u_n(t, 0_-)) &= \begin{cases} \left[k - \frac{F}{p^{-1}(\mathbf{W}(u_n(t, 0_-)) - k)} \right]_+ & \text{if } F \neq 0, \\ k & \text{if } F = 0. \end{cases} \end{aligned}$$




Figure 10: Above $F \in (0, f_c^-)$, $v_0^\pm \doteq v_n(t, 0_\pm)$ and $v_{0,F}^\pm \doteq F/p^{-1}(w(u_n(t, 0_\pm)) - k)$. With the first two pictures we show that if $v_0^- < k < v_0^+$, then $v_{0,F}^- < k$. In the last picture we consider the case $v_0^- < v_0^+ < k$ and show that $v_{0,F}^- < v_{0,F}^+ < k$.

Notice that if $F = 0$, then $u_n(t, 0_+) = (0, V)$ and $u_n(t, 0_-) \in [p^{-1}(w^-), R] \times \{0\}$. We observe, see Figures 9 and 10, that $-\Delta Q_n^{k,i}(t) < 0$ and that $-\Delta Q_n^{k,i}(t) + N_F^k(u_n(t, 0_-)) \geq 0$ and therefore

$$\left[\dot{\delta}_n^i(t) \Delta E_n^{k,i}(t) - \Delta Q_n^{k,i}(t) \right] \phi(t, \delta_n^i(t)) + N_F^k(u(t, 0_-)) \phi(t, 0) = \left[-\Delta Q_n^{k,i}(t) + N_F^k(u_n(t, 0_-)) \right] \phi(t, 0) \geq 0.$$

Thus, by proceeding as in the proof of Proposition 4.3 it is easy to see that (23) holds true. Let us just underline that beside the NSs, the only possible stationary discontinuities at $x = 0$ are PTs and CDs, however in both of these cases we have $f(u_n(t, 0_-)) = 0$ and therefore $N_F^k(u_n(t, 0_-)) = 0$.

We can now prove that if u has a non-classical discontinuity then $f(u(t, 0_\pm)) = F$. This is of course obvious if $F = 0$, due to **(CS.4)** and the fact that $f(u) \geq 0$. We can therefore assume that $F > 0$ and that $x \mapsto u(t_0, x)$ has a (stationary) non-classical shock (u_ℓ, u_r) , with $v_\ell < v_r$ and $f(u_\ell) = f(u_r) \doteq f \leq F$. We want to prove that $f = F$. Consider the test function

$$\phi(t, x) \doteq \left[\int_{|x|-\varepsilon}^{\infty} \varphi_\varepsilon(z) dz \right] \left[\int_{t-t_0+\varepsilon}^{t-t_0+2\varepsilon} \varphi_\varepsilon(z) dz \right],$$

where φ_ε is a smooth approximation of the Dirac mass centred at 0_+ , $\delta_{0_+}^D$, namely

$$\varphi_\varepsilon \in \mathbf{C}_c^\infty(\mathbb{R}; \mathbb{R}_+), \quad \varepsilon > 0, \quad \text{supp}(\delta_\varepsilon) \subseteq [0, \varepsilon], \quad \|\varphi_\varepsilon\|_{\mathbf{L}^1(\mathbb{R}; \mathbb{R})} = 1, \quad \varphi_\varepsilon \rightarrow \delta_{0_+}^D.$$

Observe that as ε goes to zero

$$\begin{aligned} \phi(t_0, x) &\equiv 0 \rightarrow 0, \\ \phi(t, 0) &= \int_{t-t_0+\varepsilon}^{t-t_0+2\varepsilon} \varphi_\varepsilon(z) dz \rightarrow \delta_{t_0-}^D(t), \\ \phi_t(t, x) &= \left[\int_{|x|-\varepsilon}^{\infty} \varphi_\varepsilon(z) dz \right] \left[\varphi_\varepsilon(t-t_0+2\varepsilon) - \varphi_\varepsilon(t-t_0+\varepsilon) \right] \rightarrow 0, \\ \chi_{\mathbb{R}_\pm}(x) \phi_x(t, x) &\rightarrow \mp \delta_{0_\pm}^D(x) \delta_{t_0-}^D(t). \end{aligned}$$

Then by (22) for all k belonging to the interval $(\check{v}(w_\ell, F), \check{v}(w_r, F))$ we have

$$\begin{aligned} &Q^k(u_\ell) - Q^k(u_r) + f \left[\frac{k}{F} - \frac{1}{p^{-1}(w(u_\ell) - k)} \right]_+ \\ &= \left[\frac{f}{p^{-1}(w(u_\ell) - k)} - k \right] + f \left[\frac{k}{F} - \frac{1}{p^{-1}(w(u_\ell) - k)} \right] = \left[\frac{f}{F} - 1 \right] k \geq 0. \end{aligned}$$

Since $f \leq F$, the above estimate implies that $f = F$ and this concludes the proof. \square

We underline that the entropy condition (8) “becomes” (22) if we do not require that the test function ϕ satisfy the condition $\phi(\cdot, 0) \equiv 0$. Even if it is not necessary for the proof of Theorem 2.2, we conclude this section by considering in (7) a test function ϕ which may not satisfy the condition $\phi(\cdot, 0) \equiv 0$.

Proposition 4.5. *Let $u^o \in \mathbf{L}^1 \cap \mathbf{BV}(\mathbb{R}; \Omega)$, $F \in [0, f_c^+]$ satisfy (H.1) or (H.2) and u be a limit of the approximate solutions $(u_n)_n$ constructed in Section 4.1. If the traces at $x = 0$ of $(u_n)_n$ and u satisfy for any $\phi \in \mathbf{C}_c^\infty((0, \infty) \times \mathbb{R}; \mathbb{R})$*

$$\begin{aligned} & \lim_{n \rightarrow \infty} \int_0^T f(u_n(t, 0_-)) \left[\mathbb{W}(u_n(t, 0_-)) - \mathbb{W}(u_n(t, 0_+)) \right]_+ \phi(t, 0) dt \\ &= \int_0^T f(u(t, 0_-)) \left[\mathbb{W}(u(t, 0_-)) - \mathbb{W}(u(t, 0_+)) \right]_+ \phi(t, 0) dt \end{aligned} \quad (24)$$

then u satisfies the following integral condition for any $\phi \in \mathbf{C}_c^\infty((0, \infty) \times \mathbb{R}; \mathbb{R})$

$$\int_0^\infty \left[\int_{\mathbb{R}} [\rho \phi_t + f(u) \phi_x] \mathbb{W}(u) dx - f(u(t, 0_-)) \left[\mathbb{W}(u(t, 0_-)) - \mathbb{W}(u(t, 0_+)) \right]_+ \phi(t, 0) \right] dt = 0.$$

Proof. Choose $T > 0$ such that $\phi(t, x) = 0$ whenever $t \geq T$. By (24), since u_n is uniformly bounded and f is uniformly continuous on bounded sets, it is sufficient to prove that

$$\int_0^T \left[\int_{\mathbb{R}} [\rho_n \phi_t + f(u_n) \phi_x] \mathbb{W}(u_n) dx - f(u_n(t, 0_-)) \left[\mathbb{W}(u_n(t, 0_-)) - \mathbb{W}(u_n(t, 0_+)) \right]_+ \phi(t, 0) \right] dt \rightarrow 0. \quad (25)$$

By the Green-Gauss formula the double integrals above can be written as

$$\int_0^T \sum_{i \in \mathbf{D}(t)} [\dot{\delta}_n^i(t) \Delta Y_n^i(t) - \Delta Q_n^i(t)] \phi(t, \delta_n^i(t)) dt,$$

where

$$\begin{aligned} \Delta Y_n^i(t) &\doteq \rho_n(t, \delta_n^i(t)_+) \mathbb{W}(u_n(t, \delta_n^i(t)_+)) - \rho_n(t, \delta_n^i(t)_-) \mathbb{W}(u_n(t, \delta_n^i(t)_-)), \\ \Delta Q_n^i(t) &\doteq f(u_n(t, \delta_n^i(t)_+)) \mathbb{W}(u_n(t, \delta_n^i(t)_+)) - f(u_n(t, \delta_n^i(t)_-)) \mathbb{W}(u_n(t, \delta_n^i(t)_-)). \end{aligned}$$

If $u_n(t, \cdot)$ does not have a non-classical shock at $\delta_n^i(t)$, then by the Rankine-Hugoniot conditions

$$\dot{\delta}_n^i(t) \Delta Y_n^i(t) - \Delta Q_n^i(t) = 0;$$

moreover, if $\delta_n^i(t) = 0$ and $u_n(t, \cdot)$ has a stationary discontinuity at $x = 0$, namely a phase transition or a contact discontinuity, then $v_n(t, 0_+) = v_n(t, 0) = 0$ and therefore $\text{sign}(v_n(t, 0_+)) = 0$.

On the other hand, if $\delta_n^i(t) = 0$ and $u_n(t, \cdot)$ has a stationary non-classical shock at $x = 0$, then

$$\dot{\delta}_n^i(t) = 0, \quad f(u_n(t, 0_\pm)) = F, \quad \mathbb{W}(u_n(t, 0_-)) \geq \mathbb{W}(u_n(t, 0_+)),$$

and therefore

$$\begin{aligned} \dot{\delta}_n^i(t) \Delta Y_n^i(t) - \Delta Q_n^i(t) &= -F \left[\mathbb{W}(u_n(t, 0_+)) - \mathbb{W}(u_n(t, 0_-)) \right] \\ &= f(u_n(t, 0_-)) \left[\mathbb{W}(u_n(t, 0_-)) - \mathbb{W}(u_n(t, 0_+)) \right]_+. \end{aligned}$$

As a consequence (25) is trivial. \square

Acknowledgements

M. D. Rosini acknowledges the support of Università degli Studi di Ferrara Project 2017 “FIR: Modelli macroscopici per il traffico veicolare o pedonale”. N. Dymski acknowledges the support of the French Government Scholarship (BGF) program for joint PhD thesis of the French Embassy in Poland.

References

- [1] ANDREIANOV, B., DONADELLO, C., AND ROSINI, M. D. A second-order model for vehicular traffics with local point constraints on the flow. *Mathematical Models and Methods in Applied Sciences* 26, 04 (2016), 751–802.
- [2] ANDREIANOV, B., DONADELLO, C., AND ROSINI, M. D. A second-order model for vehicular traffics with local point constraints on the flow. *Mathematical Models and Methods in Applied Sciences* 26, 04 (2016), 751–802.
- [3] ANDREIANOV, B. P., DONADELLO, C., RAZAFISON, U., ROLLAND, J. Y., AND ROSINI, M. D. Solutions of the aw-rascle-zhang system with point constraints. *Networks and Heterogeneous Media* 11, 1 (2016), 29–47.
- [4] AW, A., AND RASCLE, M. Resurrection of “second order” models of traffic flow. *SIAM J. Appl. Math.* 60, 3 (2000), 916–938 (electronic).
- [5] BENYAHIA, M., AND ROSINI, M. D. Entropy solutions for a traffic model with phase transitions. *Nonlinear Analysis: Theory, Methods & Applications* 141 (2016), 167 – 190.
- [6] BENYAHIA, M., AND ROSINI, M. D. A macroscopic traffic model with phase transitions and local point constraints on the flow. *Networks and Heterogeneous Media* 12, 2 (2017), 297–317.
- [7] BLANDIN, S., WORK, D., GOATIN, P., PICCOLI, B., AND BAYEN, A. A general phase transition model for vehicular traffic. *SIAM J. Appl. Math.* 71, 1 (2011), 107–127.
- [8] BRESSAN, A. *Hyperbolic systems of conservation laws*, vol. 20 of *Oxford Lecture Series in Mathematics and its Applications*. Oxford University Press, Oxford, 2000.
- [9] COLOMBO, R. M. Hyperbolic phase transitions in traffic flow. *SIAM J. Appl. Math.* 63, 2 (2002), 708–721 (electronic).
- [10] COLOMBO, R. M., AND GOATIN, P. A well posed conservation law with a variable unilateral constraint. *J. Differential Equations* 234, 2 (2007), 654–675.
- [11] COLOMBO, R. M., GOATIN, P., AND PRIULI, F. S. Global well posedness of traffic flow models with phase transitions. *Nonlinear Anal.* 66, 11 (2007), 2413–2426.
- [12] DAL SANTO, E., ROSINI, M. D., DYMSKI, N., AND BENYAHIA, M. General phase transition models for vehicular traffic with point constraints on the flow. *Mathematical Methods in the Applied Sciences* (2017), 1–19.
- [13] DELLE MONACHE, M. L., AND GOATIN, P. Stability estimates for scalar conservation laws with moving flux constraints. *Netw. Heterog. Media* 12, 2 (2017), 245–258.
- [14] DYMSKI, N., GOATIN, P., AND ROSINI, M. D. Existence of BV solutions for a non-conservative constrained Aw-Rascle-Zhang model for vehicular traffic. working paper or preprint, Feb. 2018.
- [15] GARAVELLO, M., AND GOATIN, P. The Aw-Rascle traffic model with locally constrained flow. *Journal of Mathematical Analysis and Applications* 378, 2 (2011), 634 – 648.
- [16] GOATIN, P. The Aw–Rascle vehicular traffic flow model with phase transitions. *Mathematical and computer modelling* 44, 3 (2006), 287–303.
- [17] HOLDEN, H., AND RISEBRO, N. *Front Tracking for Hyperbolic Conservation Laws*. Applied Mathematical Sciences. Springer Berlin Heidelberg, 2013.
- [18] LIGHTHILL, M. J., AND WHITHAM, G. B. On kinematic waves. II. A theory of traffic flow on long crowded roads. *Proc. Roy. Soc. London. Ser. A.* 229 (1955), 317–345.
- [19] RICHARDS, P. I. Shock waves on the highway. *Operations Res.* 4 (1956), 42–51.
- [20] ROSINI, M. D. *Macroscopic models for vehicular flows and crowd dynamics: theory and applications*. Understanding Complex Systems. Springer, Heidelberg, 2013.
- [21] ZHANG, H. M. A non-equilibrium traffic model devoid of gas-like behavior. *Transportation Research Part B: Methodological* 36, 3 (2002), 275–290.

Mohamed Benyahia, GRAN SASSO SCIENCE INSTITUTE, VIALE F. CRISPI 7, 67100 L’AQUILA, ITALY
E - mail address: benyahia.ramiz@gmail.com

Carlotta Donadello, LABORATOIRE DE MATHÉMATIQUES, CNRS UMR 6623, UNIVERSITÉ DE FRANCHE - COMTÉ, 16 ROUTE DE GRAY, 25030 BESANÇON, FRANCE
E - mail address: carlotta.donadello@univ-fcomte.fr

Nikodem Dymski, UNIWERSYTET MARIJ CURIE-SKŁODOWSKIEJ, PLAC MARIJ CURIE-SKŁODOWSKIEJ 1, 20 - 031 LUBLIN, POLAND, INRIA SOPHIA ANTIPOLIS-MÉDITERRANÉE, UNIVERSITÉ CÔTE D’AZUR, INRIA, CNRS, LJAD, 06902 SOPHIA-ANTIPOLIS, FRANCE
E - mail address: nikodem.dymski@inria.fr

Massimiliano D. Rosini, (Corresponding author), DIPARTIMENTO DI MATEMATICA E INFORMATICA, UNIVERSITÀ DI FERRARA, VIA MACHIAVELLI 35, 44121 FERRARA, ITALY
E - mail address: rsnmsm@unife.it