








## Article

# What Is the Role of Resilience and Coping Strategies on the Mental Health of the General Population during the COVID-19 Pandemic? Results from the Italian Multicentric COMET Study

Gaia Sampogna <sup>1,\*</sup>, Valeria Del Vecchio <sup>1</sup>, Vincenzo Giallonardo <sup>1</sup>, Mario Luciano <sup>1</sup> , Umberto Albert <sup>2,3</sup> , Claudia Carmassi <sup>4</sup> , Giuseppe Carrà <sup>5</sup> , Francesca Cirulli <sup>6</sup> , Bernardo Dell'Osso <sup>7,8</sup>, Giulia Menculini <sup>9</sup>, Mariagiulia Nanni <sup>10</sup>, Maurizio Pompili <sup>11</sup> , Gabriele Sani <sup>12,13</sup> , Umberto Volpe <sup>14</sup>, Valeria Bianchini <sup>15</sup> and Andrea Fiorillo <sup>1</sup>

- <sup>1</sup> Department of Psychiatry, University of Campania Luigi Vanvitelli, Largo Madonna delle Grazie, 80138 Naples, Italy; valeria.delvecchio78@gmail.com (V.D.V.); enzogiallo86@gmail.com (V.G.); mario.luciano@unicampania.it (M.L.); andrea.fiorillo@unicampania.it (A.F.)
- <sup>2</sup> Department of Medicine, Surgery and Health Sciences, University of Trieste, 34128 Trieste, Italy; ualbert@units.it
- <sup>3</sup> Department of Mental Health, Psychiatric Clinic, Azienda Sanitaria Universitaria Giuliano-Isontina—ASUGI, 34128 Trieste, Italy
- <sup>4</sup> Department of Clinical and Experimental Medicine, University of Pisa, 56126 Pisa, Italy; ccarmassi@gmail.com
- <sup>5</sup> Department of Medicine and Surgery, University of Milan Bicocca, 20126 Milano, Italy; giuseppe.carra@unimib.it
- <sup>6</sup> Center for Behavioral Sciences and Mental Health, National Institute of Health, 00161 Rome, Italy; francesca.cirulli@iss.it
- <sup>7</sup> Department of Mental Health, Department of Biomedical and Clinical Sciences “Luigi Sacco”, University of Milan, 20157 Milan, Italy; bernardo.delosso@unimi.it
- <sup>8</sup> Department of Health Sciences, Aldo Ravelli Center for Neurotechnology and Brain Therapeutic, University of Milan, 20142 Milan, Italy
- <sup>9</sup> Department of Psychiatry, University of Perugia, 06123 Perugia, Italy; giuliamenculini@gmail.com
- <sup>10</sup> Department of Biomedical and Specialty Surgical Sciences, Institute of Psychiatry, University of Ferrara, 44121 Ferrara, Italy; nnnmgl@unife.it
- <sup>11</sup> Department of Neurosciences, Mental Health and Sensory Organs, Faculty of Medicine and Psychology, Sapienza University of Rome, 00185 Roma, Italy; maurizio.pompili@uniroma1.it
- <sup>12</sup> Department of Neuroscience, Section of Psychiatry, University Cattolica del Sacro Cuore, 00168 Rome, Italy; gabriele.sani@unicatt.it
- <sup>13</sup> Department of Psychiatry, Fondazione Policlinico Agostino Gemelli IRCCS, 00168 Rome, Italy
- <sup>14</sup> Clinical Psychiatry Unit, Department of Clinical Neurosciences, Università Politecnica delle Marche, 60121 Ancona, Italy; u.volpe@staff.univpm.it
- <sup>15</sup> Department of Life, Health and Environmental Sciences, Psychiatric Unit: Trattamenti Riabilitativi Psicosociali, Interventi Precoci, TRIP, Psychosocial Rehabilitation Treatment, Early Interventions University Unit, University of L'Aquila, 67100 L'Aquila, Italy; valeria.bianchini@univaq.it
- \* Correspondence: gaia.sampogna@gmail.com



**Citation:** Sampogna, G.; Del Vecchio, V.; Giallonardo, V.; Luciano, M.; Albert, U.; Carmassi, C.; Carrà, G.; Cirulli, F.; Dell'Osso, B.; Menculini, G.; et al. What Is the Role of Resilience and Coping Strategies on the Mental Health of the General Population during the COVID-19 Pandemic? Results from the Italian Multicentric COMET Study. *Brain Sci.* **2021**, *11*, 1231. <https://doi.org/10.3390/brainsci11091231>

Academic Editor: Danielle Cha

Received: 16 August 2021

Accepted: 14 September 2021

Published: 17 September 2021

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

**Abstract:** The effects of the COVID-19 pandemic on mental health are now well documented, however, few studies have been focused on the role of coping strategies and resilience in counterbalancing these detrimental effects. Data are derived from the COvid Mental hEalth Trial (COMET), a national multicentric trial carried out in the Italian general population. The final sample consisted of 20,720 participants, 53.1% ( $n = 11,000$ ) of the sample reported low levels of resilience. Adaptive coping strategies and resilience levels did not have any significant protective impact on the levels of depressive, anxiety, and stress symptoms. Only self-distraction was a risk factor for poor mental health (Beta Coefficient,  $B = 0.1$ , 95% Confidence Interval, CI: 0.003 to 0.267 for stress symptoms;  $B = 0.2$ ; 95% CI: 0.077 to 0.324 for anxiety symptoms and  $B = 0.2$ , 95% CI: 0.105 to 0.382 for depressive symptoms). High levels of resilience were predicted by adaptive coping strategies, such as acceptance ( $B = 1.8$ , CI 95% = 1.4–2.7). Exposure to the different weeks of lockdown, being infected by COVID-19, and being a healthcare professional did not influence the levels of resilience. Our findings should be

carefully considered, since the low levels of resilience may represent the missing link between the pandemic and the current increase in mental health problems.

**Keywords:** resilience; coping strategies; trauma; pandemic; mental health

## 1. Introduction

The COVID-19 pandemic is an unprecedented life-threatening event that is affecting the mental and physical health and well-being of the general population worldwide [1–3]. It has been considered as a new traumatic experience, which is completely different from all other natural or man-made disasters [4–6]. The traumatic role of the COVID-19 pandemic is due to its direct and indirect threats to important life resources of the general population, such as safety, health, income, work, housing, and social support [7–10].

The effects of the COVID-19 pandemic on mental health have been well documented. Trials carried out in the last year have identified differential effects in samples including the general population, health professionals, those affected by COVID-19, people with disabilities, or affected by chronic physical and mental health conditions. In particular, the psychiatric and psychological consequences of the pandemic on the general population mainly include high levels of distress, insomnia, depressive and anxiety symptoms [11–19]. Health professionals are at high risk of developing burn-out and insomnia [20–24]. In disabled people and in those with pre-existing mental health problems, an increased risk of treatment interruption has been found, associated with relapses or symptoms worsening [25–29]. Finally, people affected by COVID-19 have experienced high levels of trauma-related disorders, cognitive deficits, and depression [30–34]. Specific risk factors identified for the development of these mental health disturbances include female gender, having previous mental health or physical disorders, loneliness, time spent on the Internet, and unemployment [11,25].

Although the general population as a whole is exposed to the same traumatic event, the perception of the pandemic is highly variable, being mediated by individual psychological and social strengths, including coping strategies and resilience styles.

Coping strategies are defined as the skills needed to manage and adjust to stressful situations, representing an essential element for the adaptation process to stressful and traumatic situations as well as for the recovery process of patients with severe mental disorders [35]. Lazarus and Folkman [36] identified problem-oriented and emotion-focused coping strategies. The former includes practical strategies, such as seeking information and positive communication, in order to deal with the stressful situation and are associated with a better long-term outcome for patients and relatives; the latter are psychologically driven, such as venting or avoidance, and are associated with a worse outcome [22,37].

Resilience is defined as a positive growth or adaptation following periods of homeostatic disruption [38]. This positive adaptation in response to extreme adversities was originally thought to characterize extraordinary individuals; more recently, it has been shown that resilience is relatively common also among children and adolescents exposed to adversity and trauma [39,40].

Several studies have been carried out so far in order to identify the prevalence of personal or social factors protecting people from developing mental disorders during the pandemic [41–46]. In particular, studies have been focused on the perceived levels of stress, resilience, and coping strategies related to COVID-19 in the general population [47,48], in older adults [45,49], in pregnant women [50–53], in college students [54], in children and adolescents [55,56], and in mental health professionals [57,58].

Based on the largest Italian study evaluating the effects of the COVID-19 pandemic on the mental health of the general population [11], in this paper we aimed to: (1) describe the levels of coping strategies and resilience in the Italian general population during the first wave of the pandemic; (2) evaluate the protective role of coping strategies and resilience

on the levels of depressive, anxiety and stress symptoms at DASS-21 scale [59]; (3) assess the relationship between the levels of resilience and respondents' psychiatric symptoms, socio-demographic characteristics and coping strategies.

## 2. Materials and Methods

The COvid Mental hEalth Trial (COMET) is a national trial coordinated by the University of Campania "Luigi Vanvitelli" (Naples) in collaboration with nine university sites: Università Politecnica delle Marche (Ancona), University of Ferrara, University of Milan Bicocca, University of Milan "Statale", University of Perugia, University of Pisa, Sapienza University of Rome, "Catholic" University of Rome, University of Trieste. The Center for Behavioral Sciences and Mental Health of the National Institute of Health in Rome has been involved in the study by supporting the dissemination and implementation of the project according to the clinical guidelines produced by the National Institute of Health for managing the effects of the COVID-19 pandemic.

The COMET was conceived as a cross-sectional observational design using a snowball sampling method for the recruitment of the Italian general population. The full study protocol is available elsewhere [60].

The main outcome measure of the study is the DASS-21, evaluating the general distress on a tripartite model of psychopathology [59]. It consists of 21 items grouped in three subscales: depression, anxiety, and stress. The depression scale assesses dysphoria, hopelessness, devaluation of life, self-deprecation, lack of interest/involvement, anhedonia and inertia. The anxiety scale assesses autonomic arousal, skeletal muscle effects, situational anxiety, and subjective experience of anxious affect. The stress scale evaluates the levels of chronic nonspecific arousal. It assesses difficulty relaxing, nervous arousal, and being easily upset/agitated, irritable/over-reactive and impatient. Each item is rated on a 4-level Likert scale, from 0 (never) to 3 (almost always). The total score is calculated by adding together the response value of each item, with higher scores indicating more severe levels of depressive, anxiety, and stress symptoms. The score at the DASS–depression subscale (e.g., "I felt that I had nothing to look forward to") is divided into normal (0–9), mild (10–12), moderate (13–20), severe (21–27), and extremely severe depression (28–42). The score at the DASS–anxiety subscale (e.g., "I was worried about situations in which I might panic and make a fool of myself") is divided into normal (0–6), mild (7–9), moderate (10–14), severe (15–19), and extremely severe anxiety (20–42). The score at the DASS–stress subscale (e.g., "I tended to over-react to situations") is divided into normal (0–10), mild (11–18), moderate (19–26), severe (27–34), and extremely severe stress (35–42).

The levels of resilience have been evaluated by the Connor Davidson Resilience Scale (CD-RISC), which includes 10 items rated on a 6-level Likert scale. Higher values indicate higher levels of resilience [61]. As reported by Campbell-Sills et al. [62], the levels of resilience can be subdivided into quartiles, from lowest to highest quartiles: 0–29 indicate low resilience, 30–32 median resilience, 33–36 moderate resilience, and 37–40 high resilience.

Coping strategies have been evaluated using the Brief-COPE, consisting of 28 items grouped in 14 subscales [63]. Each item is rated on a 4-level Likert scale from 0 = "I have not been doing this at all" to 3 = "I have been doing this a lot". Coping strategies are grouped into maladaptive strategies, including denial, venting, behavioral disengagement, self-blame, self-distraction, and substance abuse, and adaptive coping strategies, which include emotional support, use of information, positive reframing, planning, and acceptance. Two other subscales include religion and humor.

Respondents' socio-demographic (e.g., gender, age, geographical region, working and housing condition, etc.) and clinical information (e.g., having a previous physical or mental disorder, using illicit drugs or medications, etc.) have been collected through ad-hoc schedules. Other validated and reliable questionnaires included in the study are: the General Health Questionnaire—12 items version (GHQ) [64]; the Obsessive-Compulsive Inventory—Revised version (OCI-R) [65]; the Insomnia Severity Index (ISI) [66]; the Suici-

dal Ideation Attributes Scale (SIDAS) [67]; the Severity of Acute Stress Symptoms Adult Scale (SASS) [68]; the Impact of Event Scale—short version (IES) [69]; the UCLA loneliness scale—short version [70]; the short form of Post-Traumatic Growth Inventory (PTGI) [71]; the Multidimensional Scale of Perceived Social Support (MSPPS) [72] and (only for healthcare professionals) the Maslach Burnout Inventory (MBI) [73].

### *Statistical Analysis*

Descriptive statistics were performed in order to describe the socio-demographic and clinical characteristics of the sample according to the different quartiles of resilience. Chi-square with multiple comparisons and ANOVA with Bonferroni corrections were performed for evaluating differences according to quartiles of resilience in the type of coping strategies adopted, as well as in the levels of depressive, anxiety, and stress symptoms, insomnia, post-traumatic growth, and perceived loneliness.

Multivariate linear regression models were implemented for testing the role of resilience levels and of coping strategies as predictors of the levels of depressive, anxiety, and stress symptoms (primary outcome evaluated with DASS-21). The models were adjusted for the rate of new COVID-19 cases and of COVID-19-related mortality during the study period, as well as for several socio-demographic characteristics, such as gender, age, occupational status, having a physical comorbid condition, having a pre-existing mental disorder, employment status, unemployment due to the pandemic and levels of perceived loneliness. Furthermore, in order to evaluate the impact of the duration of lockdown and of other related containment measures on the primary outcomes, the categorical variable “Week” was also entered in the regression models. Interaction terms have also been created for testing the impact of coping strategies/resilience with main socio-demographic and clinical variables.

In order to identify possible predictors of the levels of resilience, a multivariate linear regression model, weighted for the propensity score, was performed, including as independent variables: adaptive and maladaptive coping strategies, having been infected by COVID-19, having a pre-existing mental disorder, and being a healthcare professional. Furthermore, in order to evaluate the impact of the duration of lockdown and of other related containment measures on the primary outcomes, the categorical variable “Week” was also entered in the regression models. The models were adjusted for the rate of new COVID-19 cases and of COVID-19-related mortality during the study period, as well as for several socio-demographic characteristics, such as gender, age, occupational status, having a physical comorbid condition, hours spent on Internet, levels of perceived loneliness, health status, number of co-habitants, satisfaction with one’s own life, and with housing conditions. All variables have been managed as reported in detail elsewhere [25].

Missing data have been handled using the multiple imputation approach. Statistical analyses were performed using the Statistical Package for Social Sciences (SPSS), version 17.0, and STATA, version 15. For all analyses, the level of statistical significance was set at  $p < 0.05$ .

## **3. Results**

### *3.1. Global Sample*

The final sample consisted of 20,720 participants, 71% female ( $n = 14,720$ ), with a mean age of 40.4 (14.3) years; half of the respondents were in a stable relationship and were living with a partner (Table 1).


More than half of the sample ( $n = 11,000$ ; 53.1%) reported low levels of resilience, which were not associated with age ( $r = 0.008$ ,  $p = 0.235$ ) or gender ( $\chi^2 = 0.860$ ,  $p = 0.301$ ) (Figures 1 and 2); however, higher levels of resilience were found in people with a higher level of education ( $r = 0.017$ ,  $p < 0.005$ ). The levels of resilience did not differ among the general population, patients with pre-existing mental disorders and those infected by COVID-19 (Table 2).


**Table 1.** Socio-demographic and clinical characteristics of the sample ( $n = 20,720$ ).

Age, years, mean $\pm$ SD		40.4 $\pm$ 14.3
Age groups, % ( $n$ )		
	18–24 years old	15.2 (3151)
	25–55 years old	65.2 (13,514)
	55–64 years old	14.0 (2904)
	over 65 years old	5.6 (1151)
Gender, F, % ( $n$ )		71 (14,720)
Living with partner, yes, % ( $n$ )		52.2 (10,808)
University degree, yes, % ( $n$ )		62 (12,844)
Employed, yes, % ( $n$ )		70 (14,518)
Lost job due to the pandemic, yes, % ( $n$ )		6.3 (1302)
Are you practicing smart working, yes, % ( $n$ )		34.2 (7089)
Spending more time on Internet, yes, % ( $n$ )		80.1 (16,598)
Any comorbid physical condition(s), yes, % ( $n$ )		14.5 (3012)
Any mental health problem(s), yes, % ( $n$ )		5.5 (1133)
Have you been infected by COVID-19, yes, % ( $n$ )		1.4 (296)
Have you been isolated due to COVID-19 infection, yes, % ( $n$ )		1.5 (316)
Have you been in contact with someone affected by COVID-19, % ( $n$ )		4.2 (866)

**Table 2.** Differences among groups in the levels of resilience.

	Levels of Resilience M (SD)	$p$ Value
General population	28.3 (10.4)	0.368
People with pre-existing mental disorders	28.1 (10.7)	0.288
People with COVID-19 infection	28.4 (10.4)	0.849
People with pre-existing physical disorders	28.1 (10.4)	0.223

**Figure 1.** Levels of resilience in the global sample ( $n = 20,270$ ).



**Figure 2.** Levels of resilience according to the age group ( $n = 20,270$ ).

### 3.2. Differences in the Sample According to the Levels of Resilience

Low levels of resilience were associated with more severe insomnia (Insomnia Severity Index, ISI mean score:  $6.9 \pm 5.3$ ), depressive symptoms (DASS-Depression mean score:  $12.6 \pm 7.5$ ), and higher levels of perceived loneliness (UCLA mean score:  $12.1 \pm 3.3$ ) (Table 3).

**Table 3.** Differences in clinical characteristics according to the levels of resilience.

	Levels of Resilience				p Value
	Low	Median	Moderate	High	
Insomnia severity index, M (SD)	6.9 (5.3) <sup>a</sup>	6.7 (5.1)	6.6 (5.1)	6.0 (5.2) <sup>a</sup>	0.002
DASS Anxiety subscale, M (SD)	7.6 (6.8)	7.3 (6.8)	7.3 (6.7)	7.6 (6.9)	0.070
DASS Depression subscale, M (SD)	12.6 (7.5) <sup>a</sup>	12.1 (7.6)	12.4 (7.3)	12.0 (7.3) <sup>a</sup>	0.001
DASS Stress subscale, M (SD)	16.4 (7.1)	16.5 (7.2)	16.2 (6.9)	16.3 (7.1)	0.286
GHQ total score, M (SD)	17.6 (3.2)	17.4 (3.0)	17.4 (3.1)	17.4 (3.2)	0.386
IES Avoidance, M (SD)	2.3 (1.9)	2.3 (1.9)	2.4 (2.0)	2.4 (1.9)	0.701
IES Hyperarousal, M (SD)	2.5 (1.9)	2.5 (1.8)	2.6 (1.9)	2.6 (1.9)	0.201
IES Intrusiveness, M (SD)	2.1 (1.9)	2.1 (1.9)	2.1 (2.0)	2.1 (1.9)	0.662
PTGI—Appreciation for life, M (SD)	2.1 (1.3)	2.4 (1.3) <sup>a</sup>	2.5 (1.4)	2.5 (1.4) <sup>a</sup>	0.0001
PTGI—New possibilities, M (SD)	1.4 (1.1) <sup>a</sup>	1.9 (1.1)	2.0 (1.2)	2.2 (1.3) <sup>a</sup>	0.0001
PTGI—Personal strength, M (SD)	1.6 (1.3) <sup>a</sup>	2.2 (1.3)	2.5 (1.4)	2.7 (1.5) <sup>a</sup>	0.0001
PTGI—Relating to others, M (SD)	1.6 (1.3) <sup>a</sup>	1.9 (1.3)	2.1 (1.4)	2.2 (1.4) <sup>a</sup>	0.0001
PTGI—Spiritual help, M (SD)	0.9 (1.1)	1.2 (1.1)	1.3 (1.2)	1.3 (1.1)	0.0001
SASS global score, M (SD)	0.8 (0.7)	0.7 (0.7)	0.8 (0.7)	0.8 (0.7)	0.389
SIDAS global score, M (SD)	4.9 (6.8)	4.6 (6.6)	4.5 (5.8)	4.8 (6.7)	0.490
Support—Significant others, M (SD)	21.1 (7.1) <sup>a</sup>	22.4 (6.3)	23.2 (6.1)	23.4 (6.1) <sup>a</sup>	0.0001
Support—Friends, M (SD)	18.6 (6.8) <sup>a</sup>	20.6 (6.0)	21.2 (5.9)	21.8 (5.8) <sup>a</sup>	0.0001
Support—Family, M (SD)	19.7 (6.9) <sup>a</sup>	21.5 (6.2)	21.9 (6.2)	22.4 (6.2) <sup>a</sup>	0.0001
UCLA global score, M (SD)	2.0 (0.5) <sup>a</sup>	1.9 (0.5)	1.8 (0.5)	1.7 (0.5) <sup>a</sup>	0.0001

DASS, Depression, anxiety, and Stress scale; IES, Impact of Event Scale; GHQ, General Health Questionnaire; PTGI, Post-Traumatic Growth Inventory; SASS, Severity of Acute Stress Symptoms; UCLA, UCLA loneliness scale; M, mean; SD, standard deviation; <sup>a</sup> pairwise comparison,  $p < 0.005$ .

People with low versus high levels of resilience did not differ in intrusive, avoidance, and hyperarousal symptoms at Impact of Event Scale (IES), suicidal ideation, personal possibilities, and sense of closeness with others at Post-Traumatic Growth Inventory (PTGI). On the other hand, highly resilient people reported a significantly higher level of appreciation for life (Table 3).

People with low levels of resilience rarely used positive coping strategies. For example, only 22.5% of those with low levels of resilience used “planning” as a strategy as compared to 54.2% of people identified as having high resilience ( $p < 0.0001$ ). “Acceptance” was used in 28.1% of cases among those with low levels of resilience in contrast with 60.7% in people with high levels of resilience ( $p < 0.0001$ ). On the other hand, maladaptive coping strategies, such as behavioral disengagement was adopted by 63.8% of people in the low resilience group compared to 48.3% in the high resilience group ( $p < 0.0001$ ).

### 3.3. Impact of Coping Strategies and Resilience on Mental Health Status

According to the multivariate regression models, weighted for the propensity score, adaptive coping strategies did not have any influence on the levels of depressive and anxiety symptoms, even controlling for the impact of age, gender, presence of pre-existing mental/physical conditions, as well as the levels of loneliness. Only practical support is associated with lower levels of stress symptoms, with a Beta coefficient (B) of  $-0.186$  (95% CI:  $-0.371$  to  $-0.001$ ) (Table 3). Among the maladaptive coping strategies, only self-distraction was a risk factor for poor mental health, with a Beta coefficient of  $0.162$  (95% CI:  $0.038$  to  $0.286$ ) for anxiety symptoms, and  $B = 0.182$  (95% CI:  $0.044$  to  $0.321$ ) for depressive symptoms (Table 3). The levels of resilience did not have any influence on stress symptoms ( $B = -0.001$ ,  $p < 0.984$ ), depressive symptoms ( $B = -0.008$ ;  $p < 0.230$ ) or anxiety symptoms ( $B = -0.010$ ,  $p < 0.075$ ) (Table 4).

Being female, older, having a pre-existing mental or physical condition were significantly associated with higher levels of depressive, anxiety and stress symptoms.

Interaction terms (i.e., COPE emotional disengagement \* pre-existing mental disorder) have been created and included in the models, but no significant effects were identified (Supplementary Materials Table S1).

### 3.4. Impact of Coping Strategies on Resilience Levels

According to the multivariate regression model, weighted for the propensity score, high levels of resilience were predicted by the presence of adaptive coping strategies, such as acceptance ( $B = 1.8$ , CI 95% =  $1.4$ – $2.7$ ), planning ( $B = 2.1$ , CI 95% =  $1.7$ – $2.5$ ), and positive reframing ( $B = 1.3$ , CI 95% =  $0.9$ – $1.6$ ). Interestingly, the levels of resilience were reduced by other adaptive coping strategies, such as the search for information ( $B = -1.1$ , 95% CI =  $-1.5$  to  $-0.6$ ) and emotional support ( $B = -1.2$ , 95% CI =  $-1.6$  to  $-0.7$ ).

On the contrary, low levels of resilience were predicted by maladaptive coping strategies, including self-blame ( $B = -0.6$ , CI 95% =  $-0.9$  to  $-0.3$ ), emotional disengagement ( $B = -1.0$ , CI 95% =  $-1.4$  to  $-0.6$ ), venting ( $B = -0.3$ , CI 95% =  $-0.7$  to  $-0.3$ ) and self-distraction ( $B = -0.7$ , CI 95% =  $-1$  to  $-0.4$ ), and by higher levels of perceived loneliness ( $B = -0.3$ , 95% CI =  $-0.39$  to  $-0.24$ ).

The exposure to the different weeks of lockdown, being infected by COVID-19, and being a healthcare professional did not influence the levels of resilience, even after controlling for gender, age, the presence of physical or psychiatric comorbidities.

The levels of depressive, anxiety, or stress symptoms, insomnia, post-traumatic symptoms, and suicidal ideation did not have any influence on the levels of resilience (Table 3).

Finally, high levels of post-traumatic growth, such as identifying new possibilities ( $B = 1.5$ , 95% CI =  $1.2$ – $1.8$ ) and improving own personal strengths ( $B = 1.5$ , 95% CI =  $1.3$ – $1.8$ ) were significant protective factors for high levels of resilience (Table 5).

Table 4. Impact of coping strategies and resilience on mental health status.

	DASS Stress				DASS Anxiety				DASS Depression			
	B	Sign.	95% Confidence Interval		B	Sign.	95% Confidence Interval		B	Sign.	95% Confidence Interval	
			Lower Bound	Upper Bound			Lower Bound	Upper Bound			Lower Bound	Upper Bound
Intercept	18.292	0.000	16.740	19.844	8.871	0.000	7.427	10.316	15.352	0.000	13.733	16.971
COPE Active coping	−0.002	0.977	−0.159	0.154	0.030	0.683	−0.115	0.176	−0.056	0.500	−0.219	0.107
COPE Denial	0.037	0.644	−0.119	0.193	0.076	0.304	−0.069	0.221	0.008	0.920	−0.154	0.171
COPE Substance abuse	0.048	0.608	−0.135	0.231	0.097	0.266	−0.074	0.267	0.061	0.534	−0.130	0.251
COPE Emotional support	0.094	0.311	−0.088	0.277	−0.055	0.530	−0.225	0.116	0.047	0.626	−0.143	0.238
COPE Practical support	<b>−0.186</b>	<b>0.049</b>	<b>−0.371</b>	<b>−0.001</b>	0.002	0.985	−0.171	0.174	−0.082	0.403	−0.275	0.111
COPE Emotional disengagement	0.134	0.120	−0.035	0.302	0.070	0.381	−0.087	0.227	0.103	0.252	−0.073	0.278
COPE Venting	0.095	0.233	−0.061	0.250	0.087	0.239	−0.058	0.232	0.088	0.287	−0.074	0.250
COPE Reframing	−0.095	0.203	−0.241	0.051	0.126	0.069	−0.010	0.262	−0.055	0.481	−0.207	0.098
COPE Planning	<b>0.216</b>	<b>0.014</b>	<b>0.044</b>	<b>0.387</b>	0.118	0.149	−0.042	0.278	0.112	0.221	−0.067	0.291
COPE Humor	0.084	0.243	−0.057	0.225	0.125	0.063	−0.007	0.256	−0.019	0.801	−0.166	0.128
COPE Acceptance	−0.077	0.341	−0.235	0.081	−0.139	0.065	−0.286	0.009	−0.002	0.978	−0.168	0.163
COPE Religion	0.031	0.625	−0.094	0.157	0.001	0.988	−0.116	0.118	−0.020	0.770	−0.151	0.112
COPE Self-blame	−0.079	0.281	−0.221	0.064	0.013	0.847	−0.120	0.146	−0.027	0.727	−0.175	0.122
COPE Self-distraction	0.101	0.138	−0.032	0.234	<b>0.162</b>	<b>0.010</b>	<b>0.038</b>	<b>0.286</b>	<b>0.182</b>	<b>0.010</b>	<b>0.044</b>	<b>0.321</b>
Resilience levels	0.000	0.984	−0.012	0.012	−0.010	0.075	−0.021	0.001	−0.008	0.230	−0.020	0.005
Gender, ref. male	<b>2.061</b>	<b>0.000</b>	<b>1.849</b>	<b>2.272</b>	<b>2.086</b>	<b>0.000</b>	<b>1.889</b>	<b>2.283</b>	<b>1.693</b>	<b>0.000</b>	<b>1.471</b>	<b>1.914</b>
Age	<b>0.056</b>	<b>0.000</b>	<b>−0.063</b>	<b>−0.049</b>	<b>−0.062</b>	<b>0.000</b>	<b>−0.069</b>	<b>−0.055</b>	<b>−0.052</b>	<b>0.000</b>	<b>−0.059</b>	<b>−0.044</b>
Quarantine, yes	−0.238	0.546	−1.010	0.534	−0.226	0.538	−0.945	0.493	−1.072	<b>0.009</b>	<b>−1.878</b>	<b>−0.267</b>
Being infected by COVID, yes	0.262	0.438	−0.400	0.924	<b>1.552</b>	<b>0.000</b>	<b>0.935</b>	<b>2.169</b>	<b>1.452</b>	<b>0.000</b>	<b>0.761</b>	<b>2.143</b>
Healthcare professional, yes	0.217	0.561	−0.514	0.947	0.051	0.883	−0.629	0.731	−0.180	0.644	−0.942	0.582
Mental disorder, yes	0.672	0.093	−0.112	1.456	<b>4.534</b>	<b>0.000</b>	<b>3.804</b>	<b>5.265</b>	<b>3.871</b>	<b>0.000</b>	<b>3.053</b>	<b>4.689</b>
Pre-existing physical condition, yes	<b>0.787</b>	<b>0.000</b>	<b>0.504</b>	<b>1.707</b>	<b>1.490</b>	<b>0.000</b>	<b>1.227</b>	<b>1.754</b>	<b>0.995</b>	<b>0.000</b>	<b>0.699</b>	<b>1.269</b>
Employed, yes	−0.269	<b>0.018</b>	−0.491	−0.047	0.611	0.000	0.404	0.818	1.324	0.000	1.092	1.556
Lost job, yes	0.696	<b>0.001</b>	0.299	1.093	<b>1.219</b>	<b>0.000</b>	<b>0.850</b>	<b>1.589</b>	<b>2.231</b>	<b>0.000</b>	<b>1.817</b>	<b>2.645</b>



Table 4. Cont.

	DASS Stress				DASS Anxiety				DASS Depression			
	B	Sign.	95% Confidence Interval		B	Sign.	95% Confidence Interval		B	Sign.	95% Confidence Interval	
			Lower Bound	Upper Bound			Lower Bound	Upper Bound			Lower Bound	Upper Bound
Time to exposure, ref. week												
March 30–April 8												
Week April 15–April 9	<b>1.642</b>	<b>00.000</b>	<b>0.866</b>	<b>2.418</b>	<b>2.444</b>	<b>00.000</b>	<b>1.721</b>	<b>3.166</b>	<b>1.645</b>	<b>00.000</b>	<b>0.836</b>	<b>2.455</b>
Week April 16–April 22	<b>1.520</b>	<b>00.000</b>	<b>0.937</b>	<b>2.102</b>	<b>1.842</b>	<b>00.000</b>	<b>1.299</b>	<b>2.384</b>	<b>1.481</b>	<b>00.000</b>	<b>0.874</b>	<b>2.089</b>
Week April 23–April 29	<b>0.935</b>	<b>00.000</b>	<b>0.430</b>	<b>1.439</b>	<b>1.009</b>	<b>00.000</b>	<b>0.539</b>	<b>1.478</b>	<b>0.810</b>	<b>0.003</b>	<b>0.283</b>	<b>1.336</b>
Week April 30–May 4	<b>0.349</b>	<b>0.041</b>	<b>0.015</b>	<b>0.684</b>	<b>0.341</b>	<b>0.032</b>	<b>0.029</b>	<b>0.653</b>	<b>0.387</b>	<b>0.030</b>	<b>0.038</b>	<b>0.736</b>
PTGI—Relating to others	0.080	0.123	−0.022	0.181	0.085	0.078	−0.009	0.180	0.050	0.353	−0.056	0.156
PTGI—New possibilities	−0.005	0.933	−0.113	0.104	−0.099	0.054	−0.200	0.002	−0.058	0.316	−0.171	0.055
PTGI—Personal strength	−0.068	0.304	−0.197	0.062	−0.127	0.039	−0.248	−0.007	−0.065	0.349	−0.200	0.071
PTGI—Spiritual help	−0.014	0.768	−0.107	0.079	0.024	0.594	−0.063	0.110	0.023	0.643	−0.074	0.120
PTGI—Appreciation life	0.005	0.938	−0.123	0.133	<b>0.198</b>	<b>0.001</b>	<b>0.079</b>	<b>0.318</b>	<b>0.216</b>	<b>0.002</b>	<b>0.083</b>	<b>0.350</b>
Support from others	0.011	0.238	−0.007	0.030	0.011	0.221	−0.007	0.028	−0.010	0.322	−0.029	0.010
Support from friends	−0.011	0.258	−0.030	0.008	−0.017	0.058	−0.034	0.001	<b>−0.022</b>	<b>0.029</b>	<b>−0.041</b>	<b>−0.002</b>
Support from family	−0.009	0.318	−0.027	0.009	−0.008	0.358	−0.025	0.009	−0.007	0.480	−0.026	0.012
UCLA Loneliness	0.011	0.518	−0.023	0.045	0.010	0.523	−0.021	0.042	<b>0.045</b>	<b>0.013</b>	<b>0.009</b>	<b>0.080</b>
Cases COVID-19	00.000	0.101	−00.000	0.001	00.000	0.108	−6.967	00.000	00.000	0.132	−2.086	00.000
Death COVID-19	00.000	0.621	−0.002	0.001	0.002	0.109	−00.000	0.003	0.001	0.385	−0.001	0.002
Model statistics												
			20.300 (39), <i>p</i> < 0.001				48.315 (39), <i>p</i> < 00.000				34.732 (39), <i>p</i> < 0.001	
F (df), <i>p</i> -value			0.035				0.082				0.060	
Adjusted R <sup>2</sup>												

PTGI—Post Traumatic Growth Inventory; DASS, Depression, Anxiety, Stress scale; COPE, Brief—COPE. Significant *p* values are highlighted in bold characters.

Table 5. Predictors of resilience levels.

	B	p-Value	95% Confidence Interval	
			Lower Bound	Upper Bound
Intercept	13.221	0.000	9.786	16.656
Time to exposure, ref. week March 30–April 8				
Week April 15–April 9	1.197	0.177	−0.540	2.934
Week April 16–April 22	0.614	0.358	−0.695	1.924
Week April 23–April 29	0.941	0.105	−0.196	2.078
Week April 30–May 4	0.236	0.534	−0.508	0.980
Age, ref. >34 ys				
Age 18–25 ys	0.204	0.421	−0.293	0.701
Age 25–28 ys	0.072	0.840	−0.629	0.773
Gender female, ref. Male	−0.171	0.500	−0.669	0.326
Having a pre-existing mental health problem	0.141	0.775	−0.827	1.110
Being health care professional	−0.384	0.241	−1.026	0.257
Having been infected by COVID	−0.031	0.952	−1.031	0.970
Being in one of the most affected Italian regions	−0.478	0.046	−0.948	−0.008
Adaptive coping strategies				
Acceptance	<b>1.826</b>	<b>0.000</b>	<b>1.474</b>	<b>2.179</b>
Emotional support	<b>−1.171</b>	<b>0.000</b>	<b>−1.580</b>	<b>−0.762</b>
Planning	<b>2.091</b>	<b>0.000</b>	<b>1.711</b>	<b>2.470</b>
Active	<b>0.622</b>	<b>0.000</b>	<b>0.273</b>	<b>0.970</b>
Positive reframing	<b>1.258</b>	<b>0.000</b>	<b>0.934</b>	<b>1.582</b>
Use of information	<b>−1.088</b>	<b>0.000</b>	<b>−1.499</b>	<b>−0.676</b>
<i>Maladaptive coping strategies</i>				
Self-blame	<b>−0.591</b>	<b>0.000</b>	<b>−0.912</b>	<b>−0.270</b>
Denial	−0.046	0.802	−0.402	0.311
Substance abuse	0.333	0.120	−0.087	0.752
Emotional disengagement	<b>−1.030</b>	<b>0.000</b>	<b>−1.406</b>	<b>−0.654</b>
Venting	−0.332	0.067	−0.689	0.024
Self-distraction	<b>−0.706</b>	<b>0.000</b>	<b>−1.006</b>	<b>−0.406</b>
<i>Other coping strategies</i>				
Humor	<b>2.284</b>	<b>00.000</b>	<b>1.970</b>	<b>2.597</b>
Religion	<b>−0.816</b>	<b>0.000</b>	<b>−1.097</b>	<b>−0.535</b>
PTGI—Appreciation life	<b>−0.449</b>	<b>0.000</b>	<b>−0.658</b>	<b>−0.239</b>
PTGI—New possibilities	<b>1.464</b>	<b>0.000</b>	<b>1.173</b>	<b>1.755</b>
PTGI—Personal strength	<b>1.540</b>	<b>0.000</b>	<b>1.297</b>	<b>1.784</b>
PTGI—Relating to others	<b>−0.264</b>	<b>0.023</b>	<b>−0.491</b>	<b>−0.036</b>
PTGI—Spiritual help	−0.032	0.829	−0.323	0.259
DASS_Stress	−0.014	0.426	−0.049	0.021
DASS Depression	0.020	0.256	−0.015	0.056
DASS Anxiety	−0.031	0.126	−0.070	0.009
IES global score	0.030	0.152	−0.011	0.070
ISI global score	−0.024	0.360	−0.074	0.027
SASS global score	0.252	0.280	−0.205	0.710
GHQ global score	0.013	0.704	−0.055	0.081
SIDAS global score	0.005	0.761	−0.029	0.040
OCI global score	−0.022	0.225	−0.056	0.013
UCLA global score	<b>−0.321</b>	<b>0.000</b>	<b>−0.398</b>	<b>−0.243</b>
Support from others	0.018	0.407	−0.025	0.061
Support from friends	<b>0.211</b>	<b>0.000</b>	<b>0.169</b>	<b>0.253</b>
Support from family	<b>0.091</b>	<b>0.000</b>	<b>0.051</b>	<b>0.132</b>
Cases COVID-19	0.010	0.285	0.000	0.001
Death COVID-19	−9.224	0.956	−0.003	0.003

Table 5. Cont.

		<b>B</b>	<b><i>p</i>-Value</b>	<b>95% Confidence Interval</b>	
				<b>Lower Bound</b>	<b>Upper Bound</b>
Model statistics					
	F (df), <i>p</i> -value		81.704 (41), <i>p</i> < 0.000		
Adjusted R <sup>2</sup>			0.388		

DASS, Depression, Anxiety and Stress scale; IES, Impact of Event Scale; GHQ, General Health Questionnaire; PTGI, Post-Traumatic Growth Inventory; SASS, Severity of Acute Stress Symptoms; UCLA, UCLA loneliness scale; OCI—obsessive-compulsive inventory; ISI—Insomnia severity index; B, Beta coefficient. Significant *p*-values are highlighted in bold characters; Italics and underline indicate the different subscales.

#### 4. Discussion

The study herein describes resilience levels in a large sample of the Italian general population during the initial phase of the pandemic. This data provides valuable insight into how the pandemic affected a Westernized country. Our findings should be carefully considered in order to develop ad hoc supportive and preventive psychosocial interventions for limiting the long-term detrimental effects of the pandemic on mental health.

The most striking finding is the presence of low levels of resilience, as reported by the majority of the sample. Since the pandemic is leading to increased levels of mental health concerns among the general population [11,25], this finding suggests that there may be a link between psychiatric symptoms and low levels of resilience [39,74].

We did not find any difference in the levels of resilience according to gender and age, although previous studies found that women are more resilient than men, and that older people are more resilient than younger ones [75–78]. This finding may suggest that this pandemic has the same effects on the whole population, regardless of gender and age. We did find a relationship between resilience and education, confirming that the development of resilience is influenced by education and by culture. However, the mean years of education of our sample were quite high and cannot be considered fully representative of the Italian general population.

During crises, people adopt different coping strategies [79], as confirmed in our sample from the general population adopting a variety of coping strategies to overcome the situation. In particular, people using self-distraction or behavioral disengagement more frequently were less likely to have high levels of resilience. This is in line with data from another survey carried out in Australia, showing that maladaptive coping strategies are associated with high levels of anxiety and distress symptoms [80].

The main hypothesis of our study, i.e., the protective role of adaptive coping strategies and resilience in the presence of depressive, anxiety, and stress symptoms, was not confirmed by our results. In fact, the most relevant factors associated with poor mental health were female gender, become unemployed due to the pandemic, the duration of lockdown, being infected by COVID-19, and having a pre-existing mental health problem [25]. These findings should be interpreted with caution as the results were likely influenced by the severe impact of the pandemic and the rapid containment measures that were implemented which is historically unprecedented.

One of the main aims of the study was to identify the predictors of the levels of resilience in the general population. People using positive coping strategies, such as planning, acceptance and reframing, have higher levels of resilience. This is in line with a study carried out on health care professionals, which showed that having positive attitudes in the workplace is associated with low levels of distress [57], greater psychological wellbeing, and better quality of life [81,82].

An interesting finding is related to the use of searching for information as an adaptive coping strategy, which appears to be a negative predictor of resilience. This finding highlights the negative role held by media during the pandemic, with sensationalistic (and sometimes biased) news rapidly spread through social media, and the tendency of the

general population to look for information more frequently than in the past [83]. For this tendency, the WHO has coined the term “infodemic”, suggesting its maladaptive role as a coping strategy [84,85]. Furthermore, there is the need for a better collaboration between health professionals and journalists in order to provide the general population with correct, balanced, and unambiguous information on the pandemic [86].

The presence of maladaptive coping strategies is a significant risk factor for low levels of resilience. This finding confirms data from previous epidemics, where people adopting avoidance or denial more frequently reported an increase in the levels of distress and a reduction in resilience [87,88]. The predictive role of coping strategies on the levels of resilience should be considered in psychiatric routine clinical practice, since the adoption of adaptive coping strategies can be improved through ad hoc psychoeducational interventions [89,90].

We also found that the levels of resilience were not influenced by the duration of lockdown, having been infected by the COVID-19 and being a healthcare professional. An earlier study carried out in Italy on healthcare professionals found that resilience was lower in healthcare professionals than in the general population after the first COVID-19 wave [91], but the small sample size may have influenced the predictive power of those findings. In our sample, we did not find any association between the role of healthcare professionals and the levels of resilience.

It is interesting that the levels of resilience were not influenced by geographical factors, infection, and mortality rates by COVID-19, nor by the duration of exposure to lockdown. In previous analyses based on the COMET study, we found that the duration of lockdown was a significant risk factor for the levels of depressive, anxiety and stress symptoms in the general population [25]. However, while the duration of lockdown and related containment measures is an evolving process, resilience is more stable as it is related to the individual’s cognitive style, personality traits, and temperament [92–96]. These aspects, which have not been considered in the current paper, will be analyzed in forthcoming analyses in order to verify the role of personality traits as possible mediators of coping strategies and resilience.

The present study has some limitations which are hereby acknowledged. First, the snowball sampling methodology could have led to a selection bias, with only those interested in the psychological consequences of the pandemic willing to participate [97]. Second, the cross-sectional design of the survey prevents us to delineate any causal relationship between the selected variables. Finally, several variables which could have an impact on coping strategies and on resilience levels, such as family functioning, family communication styles, and acceptance of restrictive measures related to the pandemic, have not been collected in our study.

## 5. Conclusions

The assessment of the levels of resilience and of coping strategies adopted by the general population in the context of different types of adverse situations is of great importance for the development of ad hoc supportive and preventive psychosocial interventions. It is well established that the pandemic will have longstanding, and far-reaching, consequences on global mental health and wellness. Therefore, the identification of those factors which may explain the causality between the pandemic and its impact on mental health will be crucial from a public health perspective. The presence of low levels of resilience in the general population may be the missing link between the pandemic and increasing concerns surrounding mental health.

**Supplementary Materials:** Supplementary materials are available online at <https://www.mdpi.com/article/10.3390/brainsci11091231/s1>, Table S1: Impact of coping strategies and resilience on mental health status, adjusted for interaction terms.

**Author Contributions:** Conceptualization, V.G., G.S. (Gaia Sampogna) and A.F.; methodology, M.L. and V.D.V.; formal analysis, G.S. (Gabriele Sani); investigation, U.A., C.C., G.S. (Gabriele Sani), G.C., M.P., U.V., B.D., M.N. and F.C.; writing—original draft preparation, G.S. (Gabriele Sani), V.B. and

G.M.; writing—review and editing, A.F., U.A., C.C., G.S. (Gabriele Sani), M.P., U.V., B.D., M.N. and F.C. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research received no external funding.

**Institutional Review Board Statement:** The study was conducted according to the guidelines of the Declaration of Helsinki, and approved by the Institutional Review Board (of University of Campania “L. Vanvitelli” (protocol code Protocol number: 0007593/I, date of approval 30 March 2020).

**Informed Consent Statement:** Informed consent was obtained from all subjects involved in the study.

**Data Availability Statement:** The data presented in this study are available on request from the corresponding author.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

1. Ghebreyesus, T.A. Addressing mental health needs: An integral part of COVID-19 response. *World Psychiatry* **2020**, *19*, 129–130. [[CrossRef](#)] [[PubMed](#)]
2. Kaufman, K.; Petkova, E.; Bhui, K.; Schulze, T. A global needs assessment in times of a global crisis: World psychiatry response to the COVID-19 pandemic. *BJPsych Open* **2020**, *6*, E48. [[CrossRef](#)] [[PubMed](#)]
3. Knapp, M.; Wong, G. Economics and mental health: The current scenario. *World Psychiatry* **2020**, *19*, 3–14. [[CrossRef](#)]
4. Solomou, I.; Constantinidou, F. Prevalence and Predictors of Anxiety and Depression Symptoms during the COVID-19 Pandemic and Compliance with Precautionary Measures: Age and Sex Matter. *Int. J. Environ. Res. Public Health* **2020**, *17*, 4924. [[CrossRef](#)]
5. Twenge, J.M.; Joiner, T.E. Mental distress among U.S. adults during the COVID-19 pandemic. *J. Clin. Psychol.* **2020**, *76*, 2170–2182. [[CrossRef](#)]
6. Marazziti, D.; Stahl, S.M. The relevance of COVID-19 pandemic to psychiatry. *World Psychiatry* **2020**, *19*, 261. [[CrossRef](#)]
7. Unützer, J.; Kimmel, R.J.; Snowden, M. Psychiatry in the age of COVID-19. *World Psychiatry* **2020**, *19*, 130–131. [[CrossRef](#)]
8. Gorwood, P.; Fiorillo, A. One year after the COVID-19: What have we learnt, what shall we do next? *Eur. Psychiatry* **2021**, *64*, e15. [[CrossRef](#)]
9. Green, M.F.; Lee, J.; Wynn, J.K. Experimental approaches to social disconnection in the general community: Can we learn from schizophrenia research? *World Psychiatry* **2020**, *19*, 177–178. [[CrossRef](#)]
10. Luciano, M.; De Rosa, C.; Del Vecchio, V.; Sampogna, G.; Sbordone, D.; Atti, A.R.; Bardicchia, F.; Bertossi, F.; Calò, S.; Cava, L.; et al. Perceived insecurity, mental health and urbanization: Results from a multicentric study. *Int. J. Soc. Psychiatry* **2016**, *62*, 252–261. [[CrossRef](#)]
11. Fiorillo, A.; Gorwood, P. The consequences of the COVID-19 pandemic on mental health and implications for clinical practice. *Eur. Psychiatry* **2020**, *63*, e32. [[CrossRef](#)]
12. McIntyre, R.S.; Lee, Y. Preventing suicide in the context of the COVID-19 pandemic. *World Psychiatry* **2020**, *19*, 250–251. [[CrossRef](#)]
13. Wasserman, D.; Iosue, M.; Wuestefeld, A.; Carli, V. Adaptation of evidence-based suicide prevention strategies during and after the COVID-19 pandemic. *World Psychiatry* **2020**, *19*, 294–306. [[CrossRef](#)]
14. Krystal, A.D.; Prather, A.A.; Ashbrook, L.H. The assessment and management of insomnia: An update. *World Psychiatry* **2019**, *18*, 337–352. [[CrossRef](#)] [[PubMed](#)]
15. McCracken, L.M.; Badinlou, F.; Buhrman, M.; Brocki, K.C. Psychological impact of COVID-19 in the Swedish population: Depression, anxiety, and insomnia and their associations to risk and vulnerability factors. *Eur. Psychiatry* **2020**, *63*, e81. [[CrossRef](#)] [[PubMed](#)]
16. Squeglia, L.M. Alcohol and the developing adolescent brain. *World Psychiatry* **2020**, *19*, 393–394. [[CrossRef](#)] [[PubMed](#)]
17. De Hert, M.; Mazereel, V.; Detraux, J.; Van Assche, K. Prioritizing COVID-19 vaccination for people with severe mental illness. *World Psychiatry* **2021**, *20*, 54–55. [[CrossRef](#)]
18. Wang, Q.; Xu, R.; Volkow, N.D. Increased risk of COVID-19 infection and mortality in people with mental disorders: Analysis from electronic health records in the United States. *World Psychiatry* **2021**, *20*, 124–130. [[CrossRef](#)]
19. Tyrer, P. COVID-19 health anxiety. *World Psychiatry* **2020**, *19*, 307–308. [[CrossRef](#)]
20. Morin, C.M.; Carrier, J. The Acute Effects of the COVID-19 Pandemic on Insomnia and Psychological Symptoms. *Sleep Med.* **2020**, *77*, 346–347. [[CrossRef](#)]
21. Al-Humadi, S.; Bronson, B.; Muhrad, S.; Paulus, M.; Hong, H.; Cáceda, R. Depression, Suicidal Thoughts, and Burnout Among Physicians During the COVID-19 Pandemic: A Survey-Based Cross-Sectional Study. *Acad. Psychiatry* **2021**, *14*, 1–9.
22. Carmassi, C.; Bertelloni, C.A.; Gesi, C.; Conversano, C.; Stratta, P.; Massimetti, G.; Rossi, A.; Dell’Osso, L. New DSM-5 PTSD guilt and shame symptoms among Italian earthquake survivors: Impact on maladaptive behaviors. *Psychiatry Res.* **2017**, *251*, 142–147. [[CrossRef](#)]

23. Carmassi, C.; Gesi, C.; Corsi, M.; Cremone, I.M.; Bertelloni, C.A.; Massimetti, E.; Olivieri, M.C.; Conversano, C.; Santini, M.; Dell'Osso, L. Exploring PTSD in emergency operators of a major University Hospital in Italy: A preliminary report on the role of gender, age, and education. *Ann. Gen. Psychiatry* **2018**, *17*, 17. [[CrossRef](#)]
24. Dong, Z.Q.; Ma, J.; Hao, Y.N.; Shen, X.L.; Liu, F.; Gao, Y.; Zhang, L. The social psychological impact of the COVID-19 pandemic on medical staff in China: A cross-sectional study. *Eur. Psychiatry* **2020**, *63*, e65. [[CrossRef](#)]
25. Fiorillo, A.; Sampogna, G.; Giallonardo, V.; Del Vecchio, V.; Luciano, M.; Albert, U.; Carmassi, C.; Carrà, G.; Cirulli, F.; Dell'Osso, B.; et al. Effects of the lockdown on the mental health of the general population during the COVID-19 pandemic in Italy: Results from the COMET collaborative network. *Eur. Psychiatry* **2020**, *63*, e87. [[CrossRef](#)]
26. Storch, E.A.; Sheu, J.C.; Guzick, A.G.; Schneider, S.C.; Cepeda, S.L.; Rombado, B.R.; Gupta, R.; Hoch, C.T.; Goodman, W.K. Impact of the COVID-19 pandemic on exposure and response prevention outcomes in adults and youth with obsessive-compulsive disorder. *Psychiatry Res.* **2021**, *295*, 113597. [[CrossRef](#)]
27. Chatterjee, S.S.; Barikar, C.M.; Mukherjee, A. Impact of COVID-19 pandemic on pre-existing mental health problems. *Asian J. Psychiatr.* **2020**, *51*, 102071. [[CrossRef](#)] [[PubMed](#)]
28. Alonzi, S.; La Torre, A.; Silverstein, M.W. The psychological impact of preexisting mental and physical health conditions during the COVID-19 pandemic. *Psychol. Trauma* **2020**, *12*, S236–S238. [[CrossRef](#)]
29. Li, J.; Yang, Z.; Qiu, H.; Wang, Y.; Jian, L.; Ji, J.; Li, K. Anxiety and depression among general population in China at the peak of the COVID-19 epidemic. *World Psychiatry* **2020**, *19*, 249–250. [[CrossRef](#)]
30. Janiri, D.; Carfi, A.; Kotzalidis, G.D.; Bernabei, R.; Landi, F.; Sani, G. Gemelli Against COVID-19 Post-Acute Care Study Group. Posttraumatic Stress Disorder in Patients After Severe COVID-19 Infection. *JAMA Psychiatry* **2021**, *78*, 567–569. [[CrossRef](#)]
31. Bryant, R.A. Post-traumatic stress disorder: A state-of-the-art review of evidence and challenges. *World Psychiatry* **2019**, *18*, 259–269. [[CrossRef](#)]
32. Qiu, D.; Li, Y.; Li, L.; He, J.; Ouyang, F.; Xiao, S. Prevalence of post-traumatic stress symptoms among people influenced by coronavirus disease 2019 outbreak: A meta-analysis. *Eur. Psychiatry* **2021**, *64*, e30. [[CrossRef](#)] [[PubMed](#)]
33. Rooksby, M.; Furuhashi, T.; McLeod, H.J. Hikikomori: A hidden mental health need following the COVID-19 pandemic. *World Psychiatry* **2020**, *19*, 399–400. [[CrossRef](#)] [[PubMed](#)]
34. Reed, G.M.; First, M.B.; Kogan, C.S.; Hyman, S.E.; Gureje, O.; Gaebel, W.; Maj, M.; Stein, D.J.; Maercker, A.; Tyrer, P.; et al. Innovations and changes in the ICD-11 classification of mental, behavioural and neurodevelopmental disorders. *World Psychiatry* **2019**, *18*, 3–19. [[CrossRef](#)] [[PubMed](#)]
35. Tintori, A.; Cerbara, L.; Ciancimino, G.; Crescimbene, M.; La Longa, F.; Versari, A. Adaptive behavioural coping strategies as reaction to COVID-19 social distancing in Italy. *Eur. Rev. Med. Pharmacol. Sci.* **2020**, *24*, 10860–10866.
36. Lazarus, R.S.; Folkman, S. *Stress, Appraisal, and Coping*; Springer: New York, NY, USA, 1984.
37. Morina, N.; Sterr, T.N. Lack of evidence for the efficacy of psychotherapies for PTSD and depression in child and adolescent refugees. *World Psychiatry* **2019**, *18*, 107–108. [[CrossRef](#)] [[PubMed](#)]
38. Richardson, G.E. The metatheory of resilience and resiliency. *J. Clin. Psychol.* **2002**, *58*, 307–321. [[CrossRef](#)]
39. Masten, A.S. Ordinary magic: Resilience processes in development. *Am. Psychol.* **2001**, *56*, 227–238. [[CrossRef](#)]
40. Feldman, R. What is resilience: An affiliative neuroscience approach. *World Psychiatry* **2020**, *19*, 132–150. [[CrossRef](#)]
41. Magson, N.R.; Freeman, J.Y.A.; Rapee, R.M.; Richardson, C.E.; Oar, E.L.; Fardouly, J. Risk and Protective Factors for Prospective Changes in Adolescent Mental Health during the COVID-19 Pandemic. *J. Youth Adolesc.* **2021**, *50*, 44–57. [[CrossRef](#)]
42. Coulombe, S.; Pacheco, T.; Cox, E.; Khalil, C.; Doucerain, M.M.; Auger, E.; Meunier, S. Risk and Resilience Factors During the COVID-19 Pandemic: A Snapshot of the Experiences of Canadian Workers Early on in the Crisis. *Front. Psychol.* **2020**, *11*, 580702. [[CrossRef](#)]
43. Karatzias, T.; Shevlin, M.; Hyland, P.; Ben-Ezra, M.; Cloitre, M.; Owkzarek, M.; McElroy, E. The network structure of ICD-11 complex post-traumatic stress disorder across different traumatic life events. *World Psychiatry* **2020**, *19*, 400–401. [[CrossRef](#)]
44. McElroy, E.; Shevlin, M.; Murphy, S.; Roberts, B.; Makhshvili, N.; Javakhishvili, J.; Bisson, J.; Ben-Ezra, M.; Hyland, P. ICD-11 PTSD and complex PTSD: Structural validation using network analysis. *World Psychiatry* **2019**, *18*, 236–237. [[CrossRef](#)]
45. Vannini, P.; Gagliardi, G.P.; Kuppe, M.; Dossett, M.L.; Donovan, N.J.; Gatchel, J.R.; Quiroz, Y.T.; Premnath, P.Y.; Amariglio, R.; Sperling, R.A.; et al. Stress, resilience, and coping strategies in a sample of community-dwelling older adults during COVID-19. *J. Psychiatr. Res.* **2021**, *138*, 176–185. [[CrossRef](#)]
46. Park, C.L.; Finkelstein-Fox, L.; Russell, B.S.; Fendrich, M.; Hutchison, M.; Becker, J. Psychological resilience early in the COVID-19 pandemic: Stressors, resources, and coping strategies in a national sample of Americans. *Am. Psychol.* **2021**, *3*. [[CrossRef](#)]
47. Petzold, M.B.; Bendau, A.; Plag, J.; Pyrkosch, L.; Mascarell Maricic, L.; Betzler, F.; Rogoll, J.; Große, J.; Ströhle, A. Risk, resilience, psychological distress, and anxiety at the beginning of the COVID-19 pandemic in Germany. *Brain Behav.* **2020**, *10*, e01745. [[CrossRef](#)]
48. Alegria, M.; Falgas-Bague, I.; Fong, H.F. Engagement of ethnic minorities in mental health care. *World Psychiatry* **2020**, *19*, 35–36. [[CrossRef](#)] [[PubMed](#)]
49. Backhaus, I.; Sisenop, F.; Begotaraj, E.; Cachia, J.; Capolongo, S.; Carta, M.G.; Jakubauskiene, M.; Jevtic, M.; Nakov, V.; Pirlog, M.C.; et al. Resilience and Coping With COVID-19: The COPERS Study. *Int. J. Public Health* **2021**, *66*, 1604007. [[CrossRef](#)] [[PubMed](#)]

50. Kinser, P.A.; Jallo, N.; Amstadter, A.B.; Thacker, L.R.; Jones, E.; Moyer, S.; Rider, A.; Karjane, N.; Salisbury, A.L. Depression, Anxiety, Resilience, and Coping: The Experience of Pregnant and New Mothers During the First Few Months of the COVID-19 Pandemic. *J. Womens Health* **2021**, *30*, 654–664. [[CrossRef](#)]
51. Brown, S. Perinatal mental health and the COVID-19 pandemic. *World Psychiatry* **2020**, *19*, 333–334. [[CrossRef](#)]
52. Chandra, P.S.; Nanjundaswamy, M.H. Pregnancy specific anxiety: An under-recognized problem. *World Psychiatry* **2020**, *19*, 336–337. [[CrossRef](#)] [[PubMed](#)]
53. Glover, V. Prenatal mental health and the effects of stress on the foetus and the child. Should psychiatrists look beyond mental disorders? *World Psychiatry* **2020**, *19*, 331–332. [[CrossRef](#)]
54. Ye, Z.; Yang, X.; Zeng, C.; Wang, Y.; Shen, Z.; Li, X.; Lin, D. Resilience, Social Support, and Coping as Mediators between COVID-19-related Stressful Experiences and Acute Stress Disorder among College Students in China. *Appl. Psychol. Health Well Being* **2020**, *12*, 1074–1094. [[CrossRef](#)] [[PubMed](#)]
55. Zainel, A.A.; Qotba, H.; Al-Maadeed, A.; Al-Kohji, S.; Al Mujalli, H.; Ali, A.; Al Mannai, L.; Aladab, A.; AlSaadi, H.; AlKarbi, K.A.; et al. Psychological and Coping Strategies Related to Home Isolation and Social Distancing in Children and Adolescents During the COVID-19 Pandemic: Cross-sectional Study. *JMIR Form. Res.* **2021**, *5*, e24760. [[CrossRef](#)]
56. Duarte, C.S.; Monk, C.; Weissman, M.M.; Posner, J. Intergenerational psychiatry: A new look at a powerful perspective. *World Psychiatry* **2020**, *19*, 175–176. [[CrossRef](#)] [[PubMed](#)]
57. Babore, A.; Lombardi, L.; Viceconti, M.L.; Pignataro, S.; Marino, V.; Crudele, M.; Candelori, C.; Bramanti, S.M.; Trumello, C. Psychological effects of the COVID-2019 pandemic: Perceived stress and coping strategies among healthcare professionals. *Psychiatry Res.* **2020**, *293*, 113366. [[CrossRef](#)] [[PubMed](#)]
58. Franck, E.; Haegdorens, F.; Goossens, E.; van Gils, Y.; Portzky, M.; Somville, F.; Abuawad, M.; Sloomans, S.; Van Bogaert, P. The Role of Coping Behavior in Healthcare Workers' Distress and Somatization During the COVID-19 Pandemic. *Front. Psychol.* **2021**, *12*, 684618. [[CrossRef](#)]
59. Lovibond, S.H.; Lovibond, P.F. *Manual for the Depression Anxiety & Stress Scales*, 2nd ed.; Psychology Foundation: Sydney, Australia, 1995.
60. Giallonardo, V.; Sampogna, G.; Del Vecchio, V.; Luciano, M.; Albert, U.; Carmassi, C.; Carrà, G.; Cirulli, F.; Dell'Osso, B.; Nanni, M.G.; et al. The Impact of Quarantine and Physical Distancing Following COVID-19 on Mental Health: Study Protocol of a Multicentric Italian Population Trial. *Front. Psychiatry* **2020**, *11*, 533. [[CrossRef](#)]
61. Connor, K.M.; Davidson, J.R.T. Development of a new resilience scale: The Connor-Davidson Resilience Scale (CD-RISC). *Depress. Anxiety* **2003**, *18*, 71–82. [[CrossRef](#)]
62. Campbell-Sills, L.; Forde, D.R.; Stein, M.B. Demographic and childhood environmental predictors of resilience in a community sample. *J. Psychiatr. Res.* **2009**, *43*, 1007–1012. [[CrossRef](#)]
63. Carver, C.S. You want to measure coping but your protocol' too long: Consider the brief cope. *Int. J. Behav. Med.* **1997**, *4*, 92–100. [[CrossRef](#)] [[PubMed](#)]
64. Goldberg, D.P.; Gater, R.; Sartorius, N.; Ustun, T.B.; Piccinelli, M.; Gureje, O.; Rutter, C. The validity of two versions of the GHQ in the WHO study of mental illness in general health care. *Psychol. Med.* **1997**, *27*, 191–197. [[CrossRef](#)] [[PubMed](#)]
65. Foa, E.B.; Huppert, J.D.; Leiberg, S.; Langner, R.; Kichic, R.; Hajcak, G.; Salkovskis, P.M. The Obsessive-Compulsive Inventory: Development and validation of a short version. *Psychol. Assess.* **2002**, *14*, 485–496. [[CrossRef](#)] [[PubMed](#)]
66. Morin, C.M.; Belleville, G.; Bélanger, L.; Ivers, H. The Insomnia Severity Index: Psychometric indicators to detect insomnia cases and evaluate treatment response. *Sleep* **2011**, *34*, 601–608. [[CrossRef](#)] [[PubMed](#)]
67. van Spijker, B.A.; Batterham, P.J.; Calear, A.L.; Farrer, L.; Christensen, H.; Reynolds, J.; Kerkhof, A.J. The suicidal ideation attributes scale (SIDAS): Community-based validation study of a new scale for the measurement of suicidal ideation. *Suicide Life Threat. Behav.* **2014**, *44*, 408–419. [[CrossRef](#)]
68. Kilpatrick, D.G.; Resnick, H.S.; Friedman, M.J. *Severity of Acute Stress Symptoms—Adult (National Stressful Events Survey Acute Stress Disorder Short Scale [NSESSS])*; American Psychiatric Association: Washington, DC, USA, 2013.
69. Thoresen, S.; Tambs, K.; Hussain, A.; Heir, T.; Johansen, V.A.; Bisson, J.I. Brief measure of posttraumatic stress reactions: Impact of Event Scale-6. *Soc. Psychiatry Psychiatr. Epidemiol.* **2010**, *45*, 405–412. [[CrossRef](#)]
70. Hays, R.D.; Di Matteo, M.R. A short-form measure of loneliness. *J. Personal. Assess.* **1987**, *51*, 69–81. [[CrossRef](#)]
71. Cann, A.; Calhoun, L.G.; Tedeschi, R.G.; Taku, K.; Vishnevsky, T.; Triplett, K.N.; Danhauer, S.C. A short form of the Posttraumatic Growth Inventory. *Anxiety Stress Coping* **2010**, *23*, 127–137. [[CrossRef](#)]
72. Zimet, G.D.; Dahlem, N.W.; Zimet, S.G.; Farley, G.K. The Multidimensional Scale of Perceived Social Support. *J. Personal. Assess.* **1988**, *1*, 30–41. [[CrossRef](#)]
73. Maslach, C.; Jackson, S.E. *MBI: Maslach Burnout Inventory*; Consulting Psychologists Press: Palo Alto, CA, USA, 1981.
74. MacLeod, A. Euthymia: Why it really does matter. *World Psychiatry* **2020**, *19*, 1. [[CrossRef](#)]
75. Netuveli, G.; Wiggins, R.D.; Montgomery, S.M.; Hildon, Z.; Blane, D. Mental health and resilience at older ages: Bouncing back after adversity in the British Household Panel Survey. *J. Epidemiol. Community Health* **2008**, *62*, 987–991. [[CrossRef](#)]
76. Whitley, R.; Shepherd, G.; Slade, M. Recovery colleges as a mental health innovation. *World Psychiatry* **2019**, *18*, 141–142. [[CrossRef](#)] [[PubMed](#)]

77. Portnoy, G.A.; Relyea, M.R.; Decker, S.; Shamaskin-Garroway, A.; Driscoll, M.; Brandt, C.A.; Haskell, S.G. Understanding Gender Differences in Resilience Among Veterans: Trauma History and Social Ecology. *J. Trauma. Stress* **2018**, *31*, 845–855. [[CrossRef](#)] [[PubMed](#)]
78. Sinha, M.; Collins, P.; Herrman, H. Collective action for young people's mental health: The citiesRISE experience. *World Psychiatry* **2019**, *18*, 114–115. [[CrossRef](#)] [[PubMed](#)]
79. Kar, N.; Kar, B.; Kar, S. Stress and coping during COVID-19 pandemic: Result of an online survey. *Psychiatry Res.* **2021**, *295*, 113598. [[CrossRef](#)] [[PubMed](#)]
80. Rahman, M.A.; Hoque, N.; Alif, S.M.; Salehin, M.; Islam, S.M.S.; Banik, B.; Sharif, A.; Nazim, N.B.; Sultana, F.; Cross, W. Factors associated with psychological distress, fear and coping strategies during the COVID-19 pandemic in Australia. *Glob. Health* **2020**, *16*, 95. [[CrossRef](#)] [[PubMed](#)]
81. Cai, Y. *Renaissance of Resilience: A Buzzword or a New Ideal? Management and Organization Review*; Cambridge University Press: Cambridge, UK, 2020; Volume 16, pp. 976–980.
82. Khalid, I.; Khalid, T.J.; Qabajah, M.R.; Barnard, A.G.; Qushmaq, I.A. Healthcare workers emotions, perceived stressors and coping strategies during a MERS-CoV outbreak. *Clin. Med. Res.* **2016**, *14*, 7–14. [[CrossRef](#)]
83. Crocamo, C.; Viviani, M.; Famigliani, L.; Bartoli, F.; Pasi, G.; Carrà, G. Surveilling COVID-19 Emotional Contagion on Twitter by Sentiment Analysis. *Eur. Psychiatry* **2021**, *64*, E17. [[CrossRef](#)]
84. Rovetta, A.; Bhagavathula, A.S. COVID-19-Related Web Search Behaviors and Infodemic Attitudes in Italy: Infodemiological Study. *JMIR Public Health Surveill.* **2020**, *6*, e19374. [[CrossRef](#)]
85. Gao, J.; Zheng, P.; Jia, Y.; Chen, H.; Mao, Y.; Chen, S.; Wang, Y.; Fu, H.; Dai, J. Mental health problems and social media exposure during COVID-19 outbreak. *PLoS ONE* **2020**, *15*, e0231924. [[CrossRef](#)]
86. Maiorano, A.; Lasalvia, A.; Sampogna, G.; Pocai, B.; Ruggeri, M.; Henderson, C. Reducing Stigma in Media Professionals: Is there Room for Improvement? Results from a Systematic Review. *Can. J. Psychiatry* **2017**, *62*, 702–715. [[CrossRef](#)]
87. Phua, D.H.; Tang, H.K.; Tham, K.Y. Coping responses of emergency physicians and nurses to the 2003 severe acute respiratory syndrome outbreak. *Acad. Emerg. Med.* **2005**, *12*, 322–328. [[CrossRef](#)]
88. Teasdale, E.; Yardley, L.; Schlotz, W.; Michie, S. The importance of coping appraisal in behavioural responses to pandemic flu. *Br. J. Health Psychol.* **2012**, *17*, 44–59. [[CrossRef](#)]
89. Stewart, D.E.; Appelbaum, P.S. COVID-19 and psychiatrists' responsibilities: A WPA position paper. *World Psychiatry* **2020**, *19*, 406–407. [[CrossRef](#)]
90. Jorm, A.F.; Kitchener, B.A.; Reavley, N.J. Mental Health First Aid training: Lessons learned from the global spread of a community education program. *World Psychiatry* **2019**, *18*, 142–143. [[CrossRef](#)] [[PubMed](#)]
91. Lisi, L.; Ciaffi, J.; Bruni, A.; Mancarella, L.; Brusi, V.; Gramegna, P.; Ripamonti, C.; Quaranta, E.; Borlandelli, E.; Gallo, G.; et al. Levels and Factors Associated with Resilience in Italian Healthcare Professionals during the COVID-19 Pandemic: A Web-Based Survey. *Behav. Sci.* **2020**, *10*, 183. [[CrossRef](#)] [[PubMed](#)]
92. Carver, C.S.; Connor-Smith, J. Personality and coping. *Annu. Rev. Psychol.* **2010**, *61*, 679–704. [[CrossRef](#)] [[PubMed](#)]
93. Oshio, A.; Taku, K.; Hirano, M.; Saeed, G. Resilience and Big Five personality traits: A meta-analysis. *Personal. Individ. Differ.* **2018**, *127*, 54–60. [[CrossRef](#)]
94. Ormel, J.; Cuijpers, P.; Jorm, A.F.; Schoevers, R. Prevention of depression will only succeed when it is structurally embedded and targets big determinants. *World Psychiatry* **2019**, *18*, 111–112. [[CrossRef](#)] [[PubMed](#)]
95. Reynolds, C.F. Optimizing personalized management of depression: The importance of real-world contexts and the need for a new convergence paradigm in mental health. *World Psychiatry* **2020**, *19*, 266–268. [[CrossRef](#)]
96. Maj, M.; Stein, D.J.; Parker, G.; Zimmerman, M.; Fava, G.A.; De Hert, M.; Demyttenaere, K.; McIntyre, R.S.; Wittchen, H.U. The clinical characterization of the adult patient with depression aimed at personalization of management. *World Psychiatry* **2020**, *19*, 269–293. [[CrossRef](#)] [[PubMed](#)]
97. Baltar, F.; Brunet, I. Social research 2.0: Virtual snowball sampling method using Facebook. *Internet Res.* **2012**, *22*, 57–74. [[CrossRef](#)]