

BRIEF REPORT

Open Access



# Physiological effects of lung-protective ventilation in patients with lung fibrosis and usual interstitial pneumonia pattern versus primary ARDS: a matched-control study

Roberto Tonelli<sup>1,2,3†</sup>, Salvatore Grasso<sup>4†</sup>, Andrea Cortegiani<sup>5,6</sup>, Lorenzo Ball<sup>7,8</sup>, Ivana Castaniere<sup>1,3</sup>, Luca Tabbi<sup>1</sup>, Riccardo Fantini<sup>1</sup>, Dario Andrisani<sup>1,2,3</sup>, Filippo Gozzi<sup>1,2,3</sup>, Antonio Moretti<sup>1,3</sup>, Giulia Bruzzi<sup>1,3</sup>, Linda Manicardi<sup>1,3</sup>, Stefania Cerri<sup>1,3</sup>, Anna Valeria Samarelli<sup>1,3</sup>, Giulia Raineri<sup>1,3</sup>, Francesco Murgolo<sup>4</sup>, Andrea Carzoli<sup>1</sup>, Rossella Di Mussi<sup>4</sup>, Stefano Busani<sup>9</sup>, Raffaella Rizzoni<sup>10</sup>, Giacomo Grasselli<sup>11,12</sup>, Enrico Clini<sup>1,3,13\*†</sup> and Alessandro Marchioni<sup>1,3†</sup>

## Abstract

**Background** Although patients with interstitial pneumonia pattern (ILD-UIP) and acute exacerbation (AE) leading to severe acute respiratory failure may require invasive mechanical ventilation (MV), physiological data on lung mechanics during MV are lacking. We aimed at describing the physiological effect of lung-protective ventilation in patients with AE-ILD-UIP compared with primary ARDS.

**Methods** Partitioned lung and chest wall mechanics were assessed in a series of AE-ILD-UIP patients matched 1:1 with primary ARDS as controls (based on BMI and PaO<sub>2</sub>/FiO<sub>2</sub> ratio). Three PEEP levels (zero = ZEEP, 4–8 cmH<sub>2</sub>O = PEEP<sub>LOW</sub>, and titrated to achieve positive end-expiratory transpulmonary pressure  $P_{L,EE} = PEEP_{TITRATED}$ ) were used for measurements.

**Results** Ten AE-ILD-UIP patients and 10 matched ARDS were included. In AE-ILD-UIP median  $P_{L,EE}$  at ZEEP was  $-4.3$  [ $-7.6$ – $-2.3$ ] cmH<sub>2</sub>O and lung elastance ( $E_L$ ) 44 [40–51] cmH<sub>2</sub>O/L. At PEEP<sub>LOW</sub>,  $P_{L,EE}$  remained negative and  $E_L$  did not change ( $p = 0.995$ ) versus ZEEP. At PEEP<sub>TITRATED</sub>,  $P_{L,EE}$  increased to 0.8 [0.3–1.5] cmH<sub>2</sub>O and  $E_L$  to 49 [43–59] ( $p = 0.004$  and  $p < 0.001$  compared to ZEEP and PEEP<sub>LOW</sub>, respectively).  $\Delta P_L$  decreased at PEEP<sub>LOW</sub> ( $p = 0.018$ ) and increased at PEEP<sub>TITRATED</sub> ( $p = 0.003$ ). In matched ARDS control PEEP titration to obtain a positive  $P_{L,EE}$  did not result in significant changes in  $E_L$  and  $\Delta P_L$ .

**Conclusions** In mechanically ventilated AE-ILD-UIP patients, differently than in patients with primary ARDS, PEEP titrated to obtain a positive  $P_{L,EE}$  significantly worsened lung mechanics.

<sup>†</sup>Roberto Tonelli and Salvatore Grasso: Shared first authorship.

<sup>†</sup>Enrico Clini and Alessandro Marchioni: Shared senior authorship.

\*Correspondence:

Enrico Clini

enrico.clini@unimore.it

Full list of author information is available at the end of the article



**Keywords** Interstitial lung disease, Pulmonary fibrosis, Usual interstitial pneumonia, Acute respiratory failure, ARDS, Lung elastance, Lung elastance, Respiratory mechanics, End-inspiratory transpulmonary pressure, End-expiratory transpulmonary pressure, Invasive mechanical ventilation, VILI, Transpulmonary pressure

## Background

Patients with interstitial lung disease and usual interstitial pneumonia pattern (ILD-UIP) may experience severe acute hypoxic respiratory failure (AHRF) during acute exacerbations (AE-ILD-UIP) [1], requiring invasive respiratory support (MV) [2]; nevertheless, the mortality following MV exceeds 80% [3]. Patho-physiologically, AE-ILD resembles acute respiratory distress syndrome (ARDS) with diffuse alveolar damage (DAD), superimposed on a background of lung fibrosis [4].

In ARDS, lung-protective MV strategies contributed to mitigate ventilatory induced lung injury (VILI), thus decreasing mortality [5, 6]. Talmor and coworkers showed that an esophageal pressure ( $P_{es}$ )-guided positive end-expiratory pressure (PEEP) titration to obtain a positive end-expiratory transpulmonary pressure ( $P_{L,EE}$ ) is useful to recruit dependent lung regions, improve lung mechanics and minimize atelectrauma in these patients [7].

Retrospective data suggest that patients with AE-ILD are particularly susceptible to stress and strain, and hence at higher risk of VILI [8]. Thus, it seems straightforward to use lung-protective ventilatory strategies in these patients. However, little is known on  $P_{L,EE}$  in patients with AE-ILD-UIP and even less on the potential impact of lung-protective strategies aimed at maintaining positive  $P_{L,EE}$ .

We studied the impact of different PEEP settings (zero PEEP [ZEEP], PEEP<sub>LOW</sub> and PEEP<sub>TITRATED</sub> to obtain positive  $P_{L,EE}$ ) in patients with AE-ILD-UIP, compared it with matched primary ARDS controls. We hypothesized that the impact of PEEP titration on partitioned respiratory mechanics could be different between the groups.

## Methods

### Study setting and population

The study (ClinicalTrials.gov ID NCT05098717) was carried out at the Respiratory Intensive Care Unit (RICU) of the University Hospital of Modena (Italy) in accordance with the Ethics Committee “Area Vasta Emilia Nord” approval (registered protocol number 327/2022). Informed consent to divulgate data was obtained from participants or their relatives, as appropriate.

Patients with AE-ILD-UIP developing AHRF and consecutively admitted to the RICU (August 1st, 2016, to July 1st, 2022) were eligible for enrollment. Inclusion criteria were age > 18 years; established diagnosis of ILD with a UIP pattern on a high-resolution computed tomography scan; invasive MV in volume-controlled mode. Patients suffering from chronic obstructive pulmonary disease, neuromuscular disease and chest wall deformities were excluded. AE-ILD-UIP were then matched 1:1 by body mass index, PaO<sub>2</sub>/FIO<sub>2</sub> and acute physiology and chronic health evaluation (APACHE) II score at admission, to a group of patients with primary ARDS under MV extracted from our dataset over the same period.

### Study procedures and aim

According to our institutional protocol, patients with AE-ILD-UIP or ARDS requiring MV were submitted to a partitioned respiratory mechanics measurements within 24 h from admission during three different lung-protective strategies including low  $V_T$  (6 ml/Kg/PBW) and three consecutive PEEP levels, i.e., 0 cmH<sub>2</sub>O (ZEEP), 4–8 cmH<sub>2</sub>O (PEEP<sub>LOW</sub>), and  $P_{es}$ -guided titration to obtain positive  $P_{L,EE}$  (PEEP<sub>TITRATED</sub>). At each phase, PEEP level was maintained for 30 min before recording all respiratory parameters and arterial blood sampling (see details in Additional file 1: Supplement [9, 10]).

The aim was to report measures of partitioned respiratory mechanics under lung-protective MV at different PEEP levels in patients with AE-ILD-UIP compared with ARDS.

### Data collection and analysis plan

Demographics, clinical characteristics, available pulmonary functions tests within 12 months before AE-ILD and partitioned respiratory mechanics were collected.

Data were displayed as median and interquartile range for continuous variables and numbers and percentages for dichotomous variables. Group comparison was built using a one-to-one propensity score matching procedure with the nearest-neighbor method without replacement (caliper = 0.2). Comparison between continuous variables was performed with Wilcoxon

and Wilcoxon signed-rank tests. Dichotomous variables were compared using the  $\chi^2$  test. Kruskal–Wallis was used to test as an interaction for whether the change in respiratory mechanics and physiological variables according to PEEP settings was different between groups. Statistics was performed using SPSS version 25.0 with PSMATCHING3 R Extension command (IBM Corp., Armonk, NY, USA) and GraphPad Prism version 8.0 (GraphPad Software, Inc., La Jolla, Ca, USA) unless otherwise indicated.

**Results**

**Respiratory mechanics of AE-ILD-UIP**

Over the study period a total of 21 patients with AE-ILD-UIP underwent MV. Of these, 10 patients were analyzed according to inclusion criteria (see Additional file 1: Supplement). All of them died while on MV.

Respiratory mechanics of AE-ILD-UIP at different PEEP levels are reported in Table 1, while changes in respiratory mechanics at different levels are shown in Fig. 1 (and Additional file 1: eFigure 2, Supplement). At ZEEP the median lung elastance ( $E_L$ ) was 44.4 cmH<sub>2</sub>O/L, transpulmonary driving pressure ( $\Delta P_L$ ) was 21.1 cmH<sub>2</sub>O,  $P_{L,EE}$  was -4.3 cmH<sub>2</sub>O and end-inspiratory transpulmonary pressure ( $P_{L,EI}$ ) was 16.7 cmH<sub>2</sub>O (Table 1). During the PEEP<sub>LOW</sub> phase  $P_{L,EE}$  remained below 0 cmH<sub>2</sub>O (Table 1), median  $E_L$  and  $P_{L,EI}$  did not change (Fig. 1, panel A and E) while  $\Delta P_L$  significantly decreased from baseline ( $p=0.018$ ). During the PEEP<sub>TITRATED</sub> phase  $P_{L,EE}$  was 0.8 cmH<sub>2</sub>O (Table 1) and  $E_L$  significantly increased as compared to both ZEEP and PEEP<sub>LOW</sub> ( $p=0.04$  and  $p<0.0001$ , respectively, Fig. 1, panel A), while  $P_{L,EI}$  and  $\Delta P_L$  were higher as compared to PEEP<sub>LOW</sub> ( $p<0.001$  and  $p=0.003$ , respectively, Fig. 1, panel E and G).

**AE-ILD-UIP as compared with historical, matched, ARDS controls**

AE-ILD-UIP and matched ARDS groups were similar for SAPS II score (Additional file 1: eTable 1, Supplement). Lung infection was the cause for developing ARDS in all patients.

PEEP<sub>TITRATED</sub>, but not PEEP<sub>LOW</sub> setting, resulted in higher PEEP in ARDS () compared with AE-ILD-UIP (14

**Table 1** Blood gas analyses and partitioned respiratory mechanics of the AE-ILD-UIP and the ARDS population at different PEEP levels. Data are presented as median value and IQR

| Variable                              | AE-ILD-UIP       | ARDS             | p-value |
|---------------------------------------|------------------|------------------|---------|
| <b>ZEEP phase</b>                     |                  |                  |         |
| $E_L$ , cmH <sub>2</sub> O/L          | 44.4 (39.7–50.7) | 17.9 (9.9–23.3)  | <0.0001 |
| $E_{cww}$ , cmH <sub>2</sub> O/L      | 3.2 (2.5–5.7)    | 5.4 (4–7.4)      | 0.12    |
| $E_{tot}$ , cmH <sub>2</sub> O/L      | 49 (43.9–54.7)   | 22 (16.8–28)     | <0.0001 |
| $P_{L,EI}$ , cmH <sub>2</sub> O       | 16.7 (14.8–19)   | 4.4 (2.9–6.3)    | <0.0001 |
| $P_{L,EE}$ , cmH <sub>2</sub> O       | -4.3 (-7.6--2.3) | -4.1 (-7.6--2.9) | 0.66    |
| $\Delta P_{aw}$ , cmH <sub>2</sub> O  | 16.8 (13.8–19.3) | 14.4 (11.5–21.2) | 0.56    |
| $\Delta P_L$ , cmH <sub>2</sub> O     | 21.1 (17.8–23.6) | 9.3 (7–11.5)     | <0.0001 |
| pH, value                             | 7.42 (7.41–7.42) | 7.4 (7.37–7.41)  | 0.07    |
| pO <sub>2</sub> , cmH <sub>2</sub> O  | 74 (66.5–80)     | 83 (75–89)       | 0.3     |
| pCO <sub>2</sub> , cmH <sub>2</sub> O | 39.5 (38–45)     | 39 (36.8–40)     | 0.06    |
| <b>PEEP<sub>LOW</sub> phase</b>       |                  |                  |         |
| $E_L$ , cmH <sub>2</sub> O/L          | 43.3 (36.8–53)   | 14.6 (12.2–19.1) | <0.0001 |
| $E_{cww}$ , cmH <sub>2</sub> O/L      | 3.4 (2.3–5.6)    | 5.7 (4.3–8.3)    | 0.09    |
| $E_{tot}$ , cmH <sub>2</sub> O/L      | 48.5 (40–56.8)   | 22.1 (19.1–25.2) | <0.0001 |
| $P_{L,EI}$ , cmH <sub>2</sub> O       | 15.3 (11.3–18.7) | 10.5 (5–14)      | 0.01    |
| $P_{L,EE}$ , cmH <sub>2</sub> O       | -2.6 (-4.3--1.2) | -2.5 (-4.6--0.5) | 0.75    |
| $\Delta P_{aw}$ , cmH <sub>2</sub> O  | 16.8 (14.3–18.6) | 15.1 (11.9–21.4) | 0.9     |
| $\Delta P_L$ , cmH <sub>2</sub> O     | 18.4 (15.6–21.8) | 12.3 (8.5–16.6)  | 0.02    |
| PEEP, cmH <sub>2</sub> O              | 4 (4–4)          | 4 (4–5)          | 0.2     |
| pH, value                             | 7.41 (7.38–7.42) | 7.4 (7.36–7.42)  | 0.3     |
| pO <sub>2</sub> , cmH <sub>2</sub> O  | 93.5 (73.3–107)  | 74.5 (68.3–80.5) | 0.1     |
| pCO <sub>2</sub> , cmH <sub>2</sub> O | 41 (38.5–42)     | 42 (39.8–47)     | 0.18    |
| <b>PEEP<sub>TITRATED</sub> phase</b>  |                  |                  |         |
| $E_L$ , cmH <sub>2</sub> O/L          | 48.8 (59–42.8)   | 15.2 (12.4–19.7) | <0.0001 |
| $E_{cww}$ , cmH <sub>2</sub> O/L      | 3.7 (3.2–5.9)    | 5.7 (4.7–7.2)    | 0.1     |
| $E_{tot}$ , cmH <sub>2</sub> O/L      | 55.3 (45.9–62.5) | 20.6 (19–24.5)   | <0.0001 |
| $P_{L,EI}$ , cmH <sub>2</sub> O       | 23.3 (21.3–26.7) | 16.9 (13.5–19.2) | 0.001   |
| $P_{L,EE}$ , cmH <sub>2</sub> O       | 0.8 (0.3–1.5)    | 2.4 (0.6–4.9)    | 0.04    |
| $\Delta P_{aw}$ , cmH <sub>2</sub> O  | 19.1 (16.1–21.6) | 15.3 (9.4–17)    | 0.01    |
| $\Delta P_L$ , cmH <sub>2</sub> O     | 22.6 (20.8–25.8) | 13.9 (6.6–16.5)  | 0.0001  |
| PEEP, cmH <sub>2</sub> O              | 12 (10–14)       | 14 (12--17.5)    | 0.03    |
| pH, value                             | 7.38 (7.35–7.4)  | 7.37 (7.34–7.4)  | 0.34    |
| pO <sub>2</sub> , cmH <sub>2</sub> O  | 80 (58–93)       | 105 (80–134)     | 0.01    |
| pCO <sub>2</sub> , cmH <sub>2</sub> O | 42 (39.5–43.3)   | 45 (40–48)       | 0.08    |

AE-ILD-UIP, acute exacerbation of interstitial lung disease with usual interstitial pneumonia pattern; ARDS, acute respiratory distress syndrome; IQR, interquartile range;  $\Delta P_L$ , transpulmonary driving pressure;  $P_{L,EI}$ , end-inspiratory transpulmonary pressure;  $P_{L,EE}$ , end-expiratory transpulmonary pressure;  $\Delta P_{aw}$ , driving pressure;  $E_{tot}$ , respiratory system elastance;  $E_{cww}$ , chest wall elastance;  $E_L$ , lung elastance; PEEP, positive end-expiratory pressure

VS 12 cmH<sub>2</sub>O,  $p < 0.001$ ). At ZEEP, ARDS patients had lower  $E_L$  (17.9 cmH<sub>2</sub>O/L,  $p \leq 0.0001$ ),  $P_{L, EI}$  (4.4 cmH<sub>2</sub>O,  $p < 0.0001$ ), and  $\Delta P_L$  (9.3 cmH<sub>2</sub>O,  $p < 0.0001$ ) compared with AE-ILD-UIP. During the PEEP<sub>LOW</sub> and the PEEP<sub>TITRATED</sub> phases, ARDS patients still had lower  $E_L$  (14.6 cmH<sub>2</sub>O/L,  $p < 0.0001$  and 15.2 cmH<sub>2</sub>O/L,  $p < 0.0001$  respectively),  $P_{L, EI}$  (10.5 cmH<sub>2</sub>O,  $p < 0.0001$  and 16.9 cmH<sub>2</sub>O,  $p = 0.001$  respectively), and  $\Delta P_L$  (12.3 cmH<sub>2</sub>O,  $p = 0.02$  and 13.9 cmH<sub>2</sub>O,  $p = 0.0001$  respectively) as compared to AE-ILD-UIP.

Figure 1 shows that during the PEEP trial  $E_L$ ,  $P_{L, EI}$  and  $\Delta P_L$  were different in AE-ILD-UIP and ARDS patients.  $E_L$  remained unchanged at PEEP<sub>LOW</sub> and worsened at PEEP<sub>TITRATED</sub> in AE-ILD-UIP, whereas it did not change in patients with ARDS (Fig. 1, panel A and B).

## Discussion

With this study, we report for the first time that AE-ILD-UIP patients under lung-protective MV strategy respond favorably in terms of respiratory mechanics to low PEEP levels, whereas respond unfavorably (and rather uniformly) to a  $P_{es}$ -guided PEEP strategy to obtain positive  $P_{L, EE}$ . The mechanical behavior of AE-ILD-UIP was different from that of matched “pulmonary” ARDS controls.

In our AE-ILD-UIP patients a low PEEP strategy resulted in reduction of  $\Delta P_L$  indirectly suggesting alveolar recruitment, probably occurring in the areas of DAD superimposed to UIP. Indeed, despite we did not measure alveolar recruitment, we assume that in AE-ILD-UIP patients the lung regions spared by fibrosis but affected by DAD were likely de-recruited at ZEEP. Thus, it seems that the low PEEP strategy could be wise in patients with AE-UIP-ILD at least in terms of lung mechanics. These results are novel and referred to a cohort of patients rarely studied in the intensive care context. A previous study by Nava et al. assessed the respiratory mechanics during MV in seven patients with end-stage idiopathic pulmonary fibrosis [11] and reported values of lung elastance (46.1 cmH<sub>2</sub>O/L) similar to those found in our work. However, in that study lung mechanics were only measured at ZEEP.

Tailored PEEP titration in the context of lung-protective ventilation is still under debate [12]. During controlled MV,  $P_{L, EE}$  may be negative at ZEEP, indicating

that the dependent lung regions are compressed [13]. This condition predisposes to tidal alveolar collapse and re-opening, resulting in high local shear forces that enhance VILI (atelectrauma) [14]. Negative  $P_{L, EE}$  is common in ARDS patients ventilated with lower PEEP levels in supine position and this largely explains the beneficial physiological effects of PEEP titration to achieve a positive  $P_{L, EE}$  reported in preclinical and clinical studies [15, 16]. Notwithstanding, the EPVent-2 trial showed that these positive physiological effects have to deal with the potential PEEP-induced lung injury caused by overdistension in the non-dependent lung regions [17]. We hypothesized the “Talmor” PEEP titration protocol in AE-ILD-UIP could lead to beneficial physiological effects also in patients with AE-ILD-UIP; however we found a rather sharp increase in  $E_L$  and  $\Delta P_L$  in all of them (Fig. 1). It is tempting to speculate that  $P_{es}$ -guided PEEP titration resulted in squishing among the patchy fibrotic tissue of the non-fibrotic lung regions (so called “squishy ball lung” phenomenon) [18] and that this effect invalidated the potential benefits of alveolar recruitment in the dependent lung regions.


Our study suffers from limitations: the small sample, the lack of quantitative analysis of hyper-inflated lung tissue [19] during PEEP titration and no end-expiratory lung volume assessment [20] allow only preliminary pathophysiological insights. Moreover, we did not assess the role of fluid balance as confounding factor. Finally, a selection bias should be acknowledged, as patients with AE-ILD are not usually placed on MV given the poor prognosis.

## Conclusions

In AE-ILD-UIP mechanically ventilated patients, low PEEP strategy may improve respiratory mechanics and, at difference with primary ARDS, PEEP titrated to obtain a positive  $P_{L, EE}$  significantly worsened lung mechanics. This paves the way to larger studies to clarify the best physiological response to PEEP in these patients. However, we feel that our findings could have practical implications when managing patients with AE-ILD-UIP under MV, suggesting that low PEEP strategy may be preferable to prevent lung injury.

(See figure on next page.)

**Fig. 1** Measured individual values of  $E_L$ ,  $P_{L, EE}$ ,  $P_{L, EI}$  and  $\Delta P_L$  the matched study groups at ZEEP, PEEP<sub>LOW</sub> and PEEP<sub>TITRATED</sub> phase. When testing as an interaction for whether the change in physiological variables at different PEEP levels was different between AE-ILD-UIP and ARDS (dotted  $p$ -values line), statistical difference was found for  $E_L$  ( $p < 0.001$ , panel **A** and **B**),  $P_{L, EI}$  ( $p < 0.001$ , panel **E** and **F**,  $p < 0.00$ ) and  $\Delta P_L$  ( $< 0.001$ , panel **G** and **H**).  $E_L$ , lung elastance;  $P_{L, EI}$ , end-inspiratory transpulmonary pressure;  $P_{L, EE}$ , end-expiratory transpulmonary pressure;  $P_L$ , transpulmonary driving pressure; ZEEP, zero positive end-expiratory pressure; PEEP, positive end-expiratory pressure; AE-ILD-UIP, acute exacerbation of interstitial lung disease with usual interstitial pneumonia pattern; ARDS, acute respiratory distress syndrome



**Fig. 1** (See legend on previous page.)

## Abbreviations

|                 |   |
|-----------------|---|
| ILD             | Interstitial lung disease                         |
| AE-ILD          | Acute exacerbation of ILD                         |
| UIP             | Usual interstitial pneumonia                      |
| ARDS            | Acute respiratory distress syndrome               |
| AHRF            | Acute hypoxic respiratory failure                 |
| bpm             | Breaths per minute                                |
| MV              | Invasive mechanical ventilation                   |
| ETI             | Endotracheal intubation                           |
| NIV             | Noninvasive mechanical ventilation                |
| PEEP            | Positive end-expiratory pressure                  |
| PBW             | Predicted body weight                             |
| PSV             | Pressure support                                  |
| APACHE II       | Acute physiology and chronic health evaluation II |
| SAPS II         | Simplified Acute Physiology Score                 |
| IPF             | Idiopathic pulmonary fibrosis                     |
| RICU            | Respiratory intensive care unit                   |
| ICU             | Intensive care unit                               |
| $P_{es}$        | Esophageal pressure                               |
| $P_{es,El}$     | End-inspiratory esophageal pressure               |
| $P_{es,EE}$     | End-expiratory esophageal pressure                |
| $P_L$           | Transpulmonary pressure                           |
| $\Delta P_L$    | Transpulmonary driving pressure                   |
| $P_{L,El}$      | End-inspiratory transpulmonary pressure           |
| $P_{L,EE}$      | End-expiratory transpulmonary pressure            |
| $P_{plat}$      | End-inspiratory plateau pressure                  |
| $\Delta P_{aw}$ | Driving pressure                                  |
| $E_{tot}$       | Respiratory system elastance                      |
| $E_{cw}$        | Chest wall elastance                              |
| $E_L$           | Lung elastance                                    |
| Vt              | Tidal volume                                      |
| IQR             | Interquartile range                               |

## Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s13054-023-04682-5>.

**Additional file 1.** Respiratory mechanics assessment protocol. Study algorithm and clinical and additional mechanical characteristics of the study population.

## Acknowledgements

None.

## Author contributions

RT and SG designed the study, enrolled the patients, analyzed the data, and wrote the paper and should be considered as first authors. IC, LT, RF, DA, FG, GB, LM, A Moretti and A Carzoli made substantial contributions to the literature review, data collection, and paper writing. AVS, GR, and SB reviewed the literature, wrote the manuscript, and produced the figures. A Cortegiani and LB, analyzed data and produced figures. GG designed the study. RR elaborated the analysis and wrote the paper. A Marchioni and EC designed the study, wrote, reviewed, and edited the manuscript, and share senior authorship. All authors have read and approved the final version of the manuscript. RT and SG share first authorship. AM and EC share senior authorship.

## Funding

No funding available.

## Availability of data and materials

Data are available at the Respiratory Disease Unit of the University Hospital of Modena, Italy, upon request.

## Declarations

### Ethics approval and consent to participate

This study was conducted in accordance with Ethics Committee "Area Vasta Emilia Nord" approval (registered protocol number 327/2022). Informed

consent to participate in the study and to allow their clinical data to be analyzed and published were obtained from participants, as appropriate.

### Consent for publication

Informed consent was waived because of the retrospective nature of the study and the analysis used anonymous clinical data.

### Competing interests

Authors have no competing interests with any organization or entity with a financial interest in competition with the subject, matter or materials discussed in the manuscript.

### Author details

<sup>1</sup>Respiratory Diseases Unit, Department of Medical and Surgical Sciences, University Hospital of Modena, University of Modena Reggio Emilia, Modena, Italy. <sup>2</sup>Clinical and Experimental Medicine PhD Program, University of Modena Reggio Emilia, Modena, Italy. <sup>3</sup>Laboratory of Cell Therapies and Respiratory Medicine, Department of Medical and Surgical Sciences for Children and Adults, University Hospital of Modena, Modena, Italy. <sup>4</sup>Dipartimento di Medicina di Precisione e Rigenerativa e Area Ionica (DiMePre-J) Sezione di Anestesiologia e Rianimazione, Università degli Studi di Bari "Aldo Moro", Ospedale Policlinico, Bari, Italy. <sup>5</sup>Department of Surgical, Oncological and Oral Science (Di.Chir.On.S.), University of Palermo, Palermo, Italy. <sup>6</sup>Department of Anesthesia, Intensive Care and Emergency, Policlinico Paolo Giaccone, Palermo, Italy. <sup>7</sup>Department of Surgical Sciences and Integrated Diagnostics, University of Genoa, Genoa, Italy. <sup>8</sup>Anesthesia and Critical Care, San Martino Policlinico Hospital, IRCCS for Oncology and Neurosciences, Genoa, Italy. <sup>9</sup>University Hospital of Modena, Modena, Italy. <sup>10</sup>Department of Engineering, University of Ferrara, Via Saragat 1, Ferrara, Italy. <sup>11</sup>Dipartimento di Anestesia, Rianimazione e Emergenza-Urgenza, Fondazione IRCCS (Istituto di Ricovero e Cura a Carattere Scientifico) Ca' Granda Ospedale Maggiore Policlinico, Milan, Italy. <sup>12</sup>Department of Pathophysiology and Transplantation, University of Milan, Milan, Italy. <sup>13</sup>Laboratory of Experimental Pneumology, University Hospital of Modena, Policlinico, UNIMORE, Università degli Studi di Modena e Reggio Emilia, Modena, Italy.

Received: 3 August 2023 Accepted: 7 October 2023

Published online: 18 October 2023

## References

1. Tonelli R, Castaniere I, Cortegiani A, et al. Inspiratory effort and respiratory mechanics in patients with acute exacerbation of idiopathic pulmonary fibrosis: a preliminary matched control study. *Pulmonology*. Published online September 2022:S2531043722002045. <https://doi.org/10.1016/j.pulmoe.2022.08.004>
2. Kershaw CD, Batra K, Torrealba JR, Terada LS. Characteristics and evaluation of acute exacerbations in chronic interstitial lung diseases. *Respir Med*. 2021;183:106400. <https://doi.org/10.1016/j.rmed.2021.106400>.
3. Fumeaux T, Rothmeier C, Jolliet P. Outcome of mechanical ventilation for acute respiratory failure in patients with pulmonary fibrosis. *Intensive Care Med*. 2001;27(12):1868–74. <https://doi.org/10.1007/s00134-001-1150-0>.
4. Marchioni A, Tonelli R, Ball L, et al. Acute exacerbation of idiopathic pulmonary fibrosis: Lessons learned from acute respiratory distress syndrome? *Crit Care*. 2018;22(1):80. <https://doi.org/10.1186/s13054-018-2002-4>.
5. Fan E, Del Sorbo L, Goligher EC, et al. An Official American Thoracic Society/European Society of Intensive Care Medicine/Society of Critical Care Medicine Clinical Practice Guideline: Mechanical Ventilation in Adult Patients with Acute Respiratory Distress Syndrome. *Am J Respir Crit Care Med*. 2017;195(9):1253–63. <https://doi.org/10.1164/rccm.201703-0548ST>.
6. Acute Respiratory Distress Syndrome Network, Brower RG, Matthay MA, et al. Ventilation with lower tidal volumes as compared with traditional tidal volumes for acute lung injury and the acute respiratory distress syndrome. *N Engl J Med*. 2000;342(18):1301–1308. <https://doi.org/10.1056/NEJM200005043421801>
7. Talmor D, Sarge T, Malhotra A, et al. Mechanical ventilation guided by esophageal pressure in acute lung injury. *N Engl J Med*. 2008;359(20):2095–104. <https://doi.org/10.1056/NEJMoa0708638>.

8. Mollica C, Paone G, Conti V, et al. Mechanical ventilation in patients with end-stage idiopathic pulmonary fibrosis. *Respiration*. 2010;79(3):209–15. <https://doi.org/10.1159/000225932>.
9. the PLeUral pressure working Group (PLUG—Acute Respiratory Failure section of the European Society of Intensive Care Medicine), Mauri T, Yoshida T, et al. Esophageal and transpulmonary pressure in the clinical setting: meaning, usefulness and perspectives. *Intensive Care Med*. 2016;42(9):1360–1373. <https://doi.org/10.1007/s00134-016-4400-x>
10. Chiumello D, Consonni D, Coppola S, Froio S, Crimella F, Colombo A. The occlusion tests and end-expiratory esophageal pressure: measurements and comparison in controlled and assisted ventilation. *Ann Intensive Care*. 2016;6(1):13. <https://doi.org/10.1186/s13613-016-0112-1>.
11. Nava S, Rubini F. Lung and chest wall mechanics in ventilated patients with end stage idiopathic pulmonary fibrosis. *Thorax*. 1999;54(5):390–5. <https://doi.org/10.1136/thx.54.5.390>.
12. Constantin JM, Jabaudon M, Lefrant JY, et al. Personalised mechanical ventilation tailored to lung morphology versus low positive end-expiratory pressure for patients with acute respiratory distress syndrome in France (the LIVE study): a multicentre, single-blind, randomised controlled trial. *Lancet Respir Med*. 2019;7(10):870–80. [https://doi.org/10.1016/S2213-2600\(19\)30138-9](https://doi.org/10.1016/S2213-2600(19)30138-9).
13. Kacmarek RM, Villar J, Sulemanji D, et al. Open lung approach for the acute respiratory distress syndrome: a pilot, randomized controlled trial\*. *Crit Care Med*. 2016;44(1):32–42. <https://doi.org/10.1097/CCM.0000000000001383>.
14. Bilek AM, Dee KC, Gaver DP. Mechanisms of surface-tension-induced epithelial cell damage in a model of pulmonary airway reopening. *J Appl Physiol*. 2003;94(2):770–83. <https://doi.org/10.1152/jappphysiol.00764.2002>.
15. Spieth PM, Güldner A, Carvalho AR, et al. Open lung approach vs acute respiratory distress syndrome network ventilation in experimental acute lung injury. *Br J Anaesth*. 2011;107(3):388–97. <https://doi.org/10.1093/bja/aer144>.
16. Staffieri F, Stripoli T, De Monte V, et al. Physiological effects of an open lung ventilatory strategy titrated on elastance-derived end-inspiratory transpulmonary pressure: Study in a pig model\*. *Crit Care Med*. 2012;40(7):2124–31. <https://doi.org/10.1097/CCM.0b013e31824e1b65>.
17. Beitler JR, Sarge T, Banner-Goodspeed VM, et al. Effect of titrating positive end-expiratory pressure (PEEP) with an esophageal pressure-guided strategy vs an empirical high PEEP-F<sub>IO2</sub> strategy on death and days free from mechanical ventilation among patients with acute respiratory distress syndrome: a randomized clinical trial. *JAMA*. 2019;321(9):846. <https://doi.org/10.1001/jama.2019.0555>.
18. Marchioni A, Tonelli R, Rossi G, et al. Ventilatory support and mechanical properties of the fibrotic lung acting as a “squishy ball.” *Ann Intensive Care*. 2020;10(1):13. <https://doi.org/10.1186/s13613-020-0632-6>.
19. GECOVID (GEnoa COVID-19) group, Ball L, Robba C, et al. Computed tomography assessment of PEEP-induced alveolar recruitment in patients with severe COVID-19 pneumonia. *Crit Care*. 2021;25(1):81. <https://doi.org/10.1186/s13054-021-03477-w>
20. Chiumello D, Carlesso E, Cadringer P, et al. Lung stress and strain during mechanical ventilation for acute respiratory distress syndrome. *Am J Respir Crit Care Med*. 2008;178(4):346–55. <https://doi.org/10.1164/rccm.200710-1589OC>.

## Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.